

Table of Content

Acknowledgement	iii
Abstract (English)	v
Abstract (Thai)	xi
List of Tables	xxiv
List of Figures	xxxii
Introduction	1
Chapter 1	
Literature review	4
1.1 Manganese in soil	4
1.2 Manganese uptake and translocation in the rice plant	7
1.2.1 Short distance transport	7
1.2.2 Long distance transport	10
1.2.2.1 Xylem transport	10
1.2.2.2 Phloem transport	11
1.2.3 Mobility in the phloem	12
1.2.4 Transport between the xylem and phloem	13
1.2.5 Manganese remobilization	14
Seed germination	14
Vegetative stage	14
Reproductive stage	15
1.3 Manganese deficiency	16
1.3.1 Function of manganese in plant	16

ลิขสิทธิ์มหาวิทยาลัยเชียงใหม่
 Copyright © by Chiang Mai University
 All rights reserved

Photosynthesis and oxygen evolution	16
Manganese containing enzymes	17
Manganese dependent or activated enzymes	17
Proteins, Carbohydrates and lipids	19
Cell division and extension	20
1.3.2 Response by rice plant to low manganese in soil	20
1.4 Nutrient efficiency	22
1.5 Genotypic variation of manganese deficiency	23
1.6 Genetics of Mn efficiency	25
Chapter 2 Genotypic variation in Mn efficiency in Thai rice genotypes	27
2.1 Introduction	27
2.2 Materials and Methods	29
2.2.1 Experiment 2.2.1 Genotypic variation among improved Thai rice varieties	29
2.2.2 Experiment 2.2.2 Genotypic variation among upland rice from calcareous soil area in northern Thailand	31
<i>Sub experiment 1 Variation between upland rice genotypes and seed lots with the same name</i>	31
<i>Sub experiment 2 Characterization in seed morphology between and within some of upland rice genotypes</i>	32
<i>Sub experiment 3 Responses of 3 upland rice genotypes to Mn levels</i>	32

2.2.3	Statistic analysis	36
2.3	Results	37
2.3.1	Experiment 2.2.1 Genotypic variation among improved Thai rice varieties	37
2.3.2	Experiment 2.2.2 Genotypic variation among upland rice from calcareous soil area in northern Thailand	51
	<i>Sub experiment 1 Variation between upland rice genotypes and seed lots with the same name</i>	51
	<i>Sub experiment 2 Characterization in seed morphology between and within some of upland rice genotypes</i>	60
	<i>Sub experiment 3 Responses of upland rice genotype to Mn levels</i>	63
2.4	Discussion	76
Chapter 3	Manganese acquisition in Mn efficient and inefficient rice genotypes	80
3.1	Introduction	80
3.2	Materials and Methods	82
3.2.1	Experiment 3.2.1 Manganese acquisition of rice in sand culture	82
	<i>Sub experiment 1 The influence of a Mn efficient genotype Mn nutrition of a Mn inefficient plants growing nearby</i>	82
	<i>Sub experiment 2 Ability to use different forms of Mn in</i>	83

Mn efficient and Mn inefficient genotypes

3.2.2 Experiment 3.2.2 Manganese acquisition of rice in 84
solution culture

Sub experiment 1 Comparing the responses of rice to 84
manganese deficiency with and without aeration of the
nutrient solution

Sub experiment 2 Acidification and reduction power of 85
root exudate from Mn efficient and inefficient rice
genotypes

3.2.3 Statistic analysis 86

3.3 Results 87

3.3.1 Experiment 3.3.1 Manganese acquisition of rice in 87
sand culture

Sub experiment 1 The influence of a Mn efficient genotype 87
Mn nutrition of a Mn inefficient plants growing nearby

Sub experiment 2 Ability to use different forms of Mn in 106
Mn efficient and Mn inefficient genotypes

3.3.2 Experiment 3.3.2 Manganese acquisition of rice in 119
solution culture

Sub experiment 1 Comparing the responses of rice to 119
manganese deficiency with and without aeration of the
nutrient solution

	<i>Sub experiment 2 Acidification and reduction power of root exudate from Mn efficient and inefficient rice genotypes</i>	123
	3.4 Discussion	130
Chapter 4	Response to external manganese levels in Mn efficient and inefficient rice genotypes	134
	4.1 Introduction	134
	4.2 Materials and Methods	136
	4.3 Results	138
	4.4 Discussion	155
Chapter 5	Manganese utilization in Mn efficient and inefficient rice genotypes	157
	5.1 Introduction	157
	5.2 Materials and Methods	159
	5.2.1 Experiment 5.2.1 Genotypic differences in the production and partitioning of carbohydrates between root and shoot of rice grown under Mn deficiency	159
	5.2.2 Experiment 5.2.2 Genotypic differences in the producing of grain yield of rice grown under Mn deficiency	161
	5.2.3 Statistic analysis	162

ลิขสิทธิ์มหาวิทยาลัยเชียงใหม่

Copyright © by Chiang Mai University

All rights reserved

5.3 Results	163
5.3.1 Experiment 5.3.1 Genotypic differences in the production and partitioning of carbohydrates between root and shoot of rice grown under Mn deficiency	163
5.3.2 Experiment 5.3.2 Genotypic differences in the producing of grain yield of rice grown under Mn deficiency	175
5.4 Discussion	187
Chapter 6 General Discussion	190
6.1 Genotypic variation in Mn efficiency in rice	190
6.2 Physiological response to Mn deficiency in Mn-efficient and inefficient rice genotypes	192
<i>Acquisition efficiency</i>	193
<i>Uptake efficiency</i>	194
<i>Utilization efficiency</i>	194
6.3 General conclusion	195
6.4 Further research	196
References	197
Curriculum vitae	216

ลิขสิทธิ์มหาวิทยาลัยเชียงใหม่

Copyright © by Chiang Mai University

All rights reserved

LIST OF TABLES

Table	Page
1.1 Characteristic differences in mobility of mineral nutrients in the phloem	13
2.1 The composition of nutrient solution for rice grown in solution culture (modified from Yoshida <i>et al.</i> , 1976).	30
2.2 Seed samples from Pang-Ma-Pha district, Mae Hong Son province in year 2005.	34
2.3 The composition of nutrient solution for rice grown in solution culture (modified by Insalud, 2006).	35
2.4 Mn concentration in YEB (mg Mn g ⁻¹) of five rice genotypes grown in sand culture with two levels of Mn at 8 weeks after transplanting.	44
2.5 Mn concentration in shoot (mg Mn g ⁻¹) of five rice genotypes grown in sand culture with two levels of Mn at 8 weeks after transplanting.	45
2.6 Mn concentration in root (mg Mn g ⁻¹) of five rice genotypes grown in sand culture with two levels of Mn at 8 weeks after transplanting.	46
2.7 Mn content in shoot (mg Mn plant ⁻¹) of five rice genotypes grown in sand culture with two levels of Mn at 8 weeks after transplanting.	47
2.8 Mn content in root (mg Mn plant ⁻¹) of five rice genotypes grown in sand culture at 8 weeks after transplanting.	48
2.9 Mn content in whole plant (mg Mn plant ⁻¹) of five rice genotypes grown in sand culture with two levels of Mn at 8 weeks after transplanting.	49
2.10 YEB-1 chlorophyll content of 11 seed accessions and 2 rice local checks	54

	(KDML105 and PSL1) at 30 days after transplanting (DAT) in Mn ₀ .	
2.11	YEB-1 chlorophyll content of 11 seed accessions and 2 rice local checks (KDML105 and PSL1) at 60 days after transplanting (DAT) in Mn ₀ .	55
2.12	Number of leaves of 11 seed accessions and 2 rice local checks (KDML105 and PSL1) at 30 days after transplanting (DAT) in Mn ₀ .	56
2.13	Number of leaves of 11 seed accessions and 2 rice local checks (KDML105 and PSL1) at 60 days after transplanting (DAT) in Mn ₀ .	57
2.14	Number of tillers of 11 seed accessions and 2 rice local checks (KDML105 and PSL1) at 30 days after transplanting (DAT) in Mn ₀ .	58
2.15	Number of tillers of 11 seed accessions and 2 rice local checks (KDML105 and PSL1) at 60 days after transplanting (DAT) in Mn ₀ .	59
2.16	Seed characters and Shannon's Index of 3 upland rice genotypes.	61
2.17	Grain length, width, thickness and weight between and within 3 upland rice genotypes.	62
2.18	YEB-1 chlorophyll content (SPAD unit) of 4 upland rice and 2 rice local checks (KDML105 and PSL1) grown in solution culture at 30 days after transplanting.	66
2.19	Concentration in YEB (mg Mn kg ⁻¹) of 4 upland rice and 2 rice local checks (KDML105 and PSL1) grown in solution culture at 30 days after transplanting.	70
2.20	Concentration in shoot (mg Mn kg ⁻¹) of 4 upland rice and 2 rice local checks (KDML105 and PSL1) grown in solution culture at 30 days after	71

	transplanting.	
2.21	Concentration in root (mg Mn kg ⁻¹) of 4 upland rice and 2 rice local checks (KDML105 and PSL1) grown in solution culture at 30 days after transplanting.	72
2.22	Content in shoot (mg Mn plant ⁻¹) of 4 upland rice and 2 rice local checks (KDML105 and PSL1) grown in solution culture at 30 days after transplanting.	73
2.23	Content in root (mg Mn plant ⁻¹) of 4 upland rice and 2 rice local checks (KDML105 and PSL1) grown in solution culture at 30 days after transplanting.	74
3.1	Shoot dry weight (g plant ⁻¹) of two rice genotypes grown together in pairs in the same pot and separately in sand culture at 4 and 8 weeks after transplanting.	96
3.2	Root dry weight (g plant ⁻¹) of two rice genotypes grown together in pairs in the same pot and separately in sand culture at 4 and 8 weeks after transplanting.	97
3.3	Mn concentration in YEB (mg Mn kg ⁻¹) of two rice genotypes grown together in pairs in the same pot and separately in sand culture at 4 and 8 weeks after transplanting.	99
3.4	Mn concentration in shoot (mg Mn kg ⁻¹) of two rice genotypes grown together in pairs in the same pot and separately in sand culture at 4 and 8 weeks after transplanting.	100

3.5	Mn concentration in root (mg Mn kg ⁻¹) of two rice genotypes grown together in pairs in the same pot and separately in sand culture at 4 and 8 weeks after transplanting.	101
3.6	Mn content in shoot (mg Mn plant ⁻¹) of two rice genotypes grown together in pairs in the same pot and separately in sand culture at 4 and 8 weeks after transplanting.	102
3.7	Mn content in root (mg Mn plant ⁻¹) of two rice genotypes grown together in pairs in the same pot and separately in sand culture at 4 and 8 weeks after transplanting.	103
3.8	Mn content in whole plant (mg Mn plant ⁻¹) of two rice genotypes grown together in pairs in the same pot and separately in sand culture at 4 and 8 weeks after transplanting.	104
3.9	Response to Mn (II) and Mn (IV) of YEB-1 chlorophyll content (SPAD unit) in two rice genotypes grown at 30 days after transplanting.	108
3.10	Response to Mn (II) and Mn (IV) of number of leaves (plant ⁻¹) in two rice genotypes grown at 30 days after transplanting.	109
3.11	Response to Mn (II) and Mn (IV) of number of tillers (plant ⁻¹) in two rice genotypes grown at 30 days after transplanting.	110
3.12	Response to Mn(II) and Mn (IV) of shoot dry weight (g plant ⁻¹) in two rice genotypes grown at 30 days after transplanting.	111
3.13	Responded to Mn (II) and Mn (IV) of root dry weight (g plant ⁻¹) in two rice genotypes grown at 30 day after transplanting.	112

3.14	Response to Mn (II) and Mn (IV) of Mn concentration in YEB (mg Mn Kg ⁻¹) in two rice genotypes grown at 30 days after transplanting.	113
3.15	Response to Mn (II) and Mn (IV) of shoot Mn concentration (mg Mn Kg ⁻¹) in two rice genotypes grown at 30 days after transplanting.	114
3.16	Response to Mn (II) and Mn (IV) of root Mn concentration (mg Mn Kg ⁻¹) in two rice genotypes grown at 30 days after transplanting.	115
3.17	Response to Mn(II) and Mn (IV) of shoot Mn content (mg Mn plant ⁻¹) in two rice genotypes grown at 30 days after transplanting.	116
3.18	Response to Mn (II) and Mn (IV) of root Mn content (mg Mn plant ⁻¹) in two rice genotypes grown at 30 days after transplanting.	117
3.19	Response to Mn (II) and Mn (IV) of Mn uptake efficiency (mg Mn g ⁻¹ root DW) in two rice genotypes grown at 30 days after transplanting.	118
3.20	Response to Mn deficiency of root length (cm) of KDML105 and PSL1 grown in nutrient solution with and without oxygen supply(+O ₂ and – O ₂) at 15 and 30 day after transplanting.	121
3.21	Response to Mn deficiency of shoot length (cm) of KDML105 and PSL1 grown in nutrient solution with and without oxygen supply(+O ₂ and – O ₂) at 15 and 30 days after transplanting.	122
3.22	Response to Mn of YEB-1 chlorophyll content (SPAD unit) of two rice genotypes grown at 2 – 4 weeks after transplanting.	125
3.23	Response to Mn of number of leaves (plant ⁻¹) of two rice genotypes grown at 2 -4 weeks after transplanting.	126

3.24	Response to Mn of number of tillers (plant^{-1}) of two rice genotypes grown at 2 -4 weeks after transplanting.	127
3.25	Response to Mn of number of root exudate by reduce pH in rhizosphere solution of two rice genotypes grown at 4 weeks after transplanting.	129
4.1	Response to Mn levels of YEB-1 chlorophyll content (SPAD unit) at 30 and 60 days after transplanting.	142
4.2	Response to Mn of number of leaves (plant^{-1}) at 30 and 60 days after transplanting.	143
4.3	Response to Mn of number of tillers (plant^{-1}) at 30 and 60 days after transplanting.	144
4.4	Response to Mn of shoot dry weight (g plant^{-1}) at 30 and 60 days after transplanting.	145
4.5	Response to Mn of root dry weight (g plant^{-1}) at 30 and 60 days after transplanting.	146
4.6	Response to Mn of YEB Mn concentration (mg Mn kg^{-1}) at 30 and 60 days after transplanting.	147
4.7	Response to Mn of shoot Mn concentration (mg Mn kg^{-1}) at 30 and 60 days after transplanting.	148
4.8	Response to Mn of root Mn concentration (mg Mn kg^{-1}) at 30 and 60 days after transplanting.	149
4.9	Response to Mn of shoot Mn content (mg Mn plant^{-1}) at 30 and 60 days after transplanting.	150

4.10	Response to Mn of root Mn content (mg Mn plant ⁻¹) at 30 and 60 days after transplanting.	151
4.11	Response to Mn of whole plant Mn content (mg Mn plant ⁻¹) at 30 and 60 days after transplanting.	152
4.12	Response to Mn of Mn uptake efficiency (mg Mn g ⁻¹ root dry weight) at 30 and 60 days after transplanting.	153
5.1	Response to Mn of YEB-1 chlorophyll content (SPAD unit) and relative chlorophyll content in YEB-1 at 15 and 30 days after transplanting.	166
5.2	Response to Mn of number of leaves (plant ⁻¹) and relative number of leaves at 15 and 30 days after transplanting.	167
5.3	Response to Mn of number of tillers (plant ⁻¹) and relative number of tillers at 15 and 30 days after transplanting.	168
5.4	Response to Mn of shoot and root dry weight (g plant ⁻¹) at 30 days after transplanting.	169
5.5	Response to Mn of relative shoot and root dry weight at 30 days after transplanting.	170
5.6	Response to Mn of TNC concentration (mg glucose equivalent g ⁻¹ dry weight) at 30 days after transplanting.	171
5.7	Response to Mn of relative TNC concentration at 30 days after transplanting.	172
5.8	Response to Mn Root / Shoot ratio of TNC concentration at 30 days after transplanting.	173

5.9	Response to Mn of YEB-1 chlorophyll content (SPAD unit) and relative YEB-1 chlorophyll content at 30, 60 days after transplanting and maturity.	178
5.10	Response to Mn of number of leaves (plant^{-1}) and relative number of leaves at 30, 60 days after transplanting and maturity.	179
5.11	Response to Mn of number of tillers (plant^{-1}) and relative number of tillers at 30, 60 day after transplanting and maturity.	180
5.12	Response to Mn of shoot, root and total dry weight (g plant^{-1}) at maturity.	181
5.13	Response to Mn of grain yield and relative grain yield (g plant^{-1}) at maturity.	182
5.14	Response to Mn of Mn concentration in YEB, shoot and root (mg Mn kg^{-1}) at maturity.	183
5.15	Response to Mn of Mn concentration in seed (mg Mn kg^{-1}) at maturity.	184
5.16	Response to Mn of Mn content in YEB, shoot and root (mg Mn plant^{-1}) at maturity.	185
5.17	Response to Mn of Mn content in seed (mg Mn plant^{-1}) at maturity.	186
5.18	Response to Mn of Mn uptake efficiency (mg Mn g^{-1} root DW) at maturity.	186

LIST OF FIGURES


Figure	Page
1.1 Mn oxidation-reduction cycle in the soil (Dion and Mann, 1946).	6
1.2 Model for root responses to iron deficiency in graminaceous species; Strategy II: (E) enhanced synthesis and release of phytosiderophores; (TR) translocator for Fe (III) phytosiderophores in the plasma membrane; structure of the phytosiderophore muginic acid and its corresponding Fe (III) chelate. (Marschner, 1995)	22
2.1 Relative number of leaves (number of leaves in $Mn_0/Mn_{0.5}$) of five rice genotypes.	40
2.2 Relative number of tillers (number of tillers in $Mn_0/Mn_{0.5}$) of five rice genotypes.	40
2.3 Relative YEB chlorophyll content (YEB chlorophyll in $Mn_0/Mn_{0.5}$) of five rice genotypes.	41
2.4 Relative shoot dry weight (dry weight in $Mn_0/Mn_{0.5}$) of five rice genotypes at 8 week after transplanting.	42
2.5 Relative root dry weight (dry weight in $Mn_0/Mn_{0.5}$) of five rice genotypes at 8 week after transplanting.	42
2.6 Relative total dry weight (dry weight in $Mn_0/Mn_{0.5}$) of five rice genotypes at 8 week after transplanting.	43
2.7 Relative Mn uptake efficiency (Mn uptake efficiency in $Mn_0/Mn_{0.5}$) at Mn_0 compared with $Mn_{0.5}$ in sand culture at 8 weeks after transplanting.	50

2.8	Relative Mn uptake (whole plant content in $Mn_0/Mn_{0.5}$) at Mn_0 compared with $Mn_{0.5}$ in sand culture at 8 weeks after transplanting.	50
2.9	Relative chlorophyll content in YEB-1 of 4 upland rice genotypes and 2 local rice checks (KDML105 and PSL1) grown in solution culture at 30 days after transplanting.	67
2.10	Relative shoot dry weight of 4 upland rice and 2 rice local checks (KDML105 and PSL1) grown in solution culture at 30 days after transplanting.	68
2.11	Relative root dry weight of 4 upland rice and 2 rice local checks (KDML105 and PSL1) grown in solution culture at 30 days after transplanting.	68
2.12	Relative total dry weight of 4 upland rice and 2 rice local checks (KDML105 and PSL1) grown in solution culture at 30 days after transplanting.	69
2.13	Relative Mn uptake efficiency of 4 upland rice and 2 rice local checks (KDML105 and PSL1) grown in solution culture at 30 days after transplanting.	75
3.1	Model of PSL1 and KDML105 were grown together in pairs in the same pot and separately.	83
3.2	Relative $Mn_0 / Mn_{0.5}$ of chlorophyll content in YEB-1 at 1 - 8 weeks after transplanting. ($Mn_0 = 0$ ppm and $Mn_{0.5} = 0.5$ ppm)	93
3.3	Relative $Mn_0 / Mn_{0.5}$ of number of tillers at 1 - 8 weeks after	94

	transplanting.	
3.4	Relative $Mn_0 / Mn_{0.5}$ of number of leaves at 1 - 8 weeks after transplanting.	95
3.5	Relative total dry weight of two rice genotypes at Mn_0 compared with $Mn_{0.5}$ in sand culture at (A) 4 weeks and (B) 8 weeks after transplanting.	98
3.6	Relative Mn uptake efficiency at Mn_0 compared with $Mn_{0.5}$ of rice genotypes grown in sand culture at (A) 4 weeks and (B) 8 weeks after transplanting.	105
3.7	Response to Mn deficiency of chlorophyll content in YEB-1 (SPAD unit) of KDML105 and PSL1 grown in nutrient solution with and without oxygen supply(+O ₂ and -O ₂) at 15 days after transplanting.	120
3.8	Response to Mn deficiency of chlorophyll content in YEB-1 (SPAD unit) of KDML105 and PSL1 grown in nutrient solution with and without oxygen supply(+O ₂ and -O ₂) at 30 days after transplanting.	120
3.9	Response to Mn deficient of root exudate by reduction of MnIV-oxide applied on filter paper of two rice genotypes grown at 4 weeks after transplanting.	128
3.10	Response to Mn sufficient of root exudate by reduction of MnIV-oxide applied on filter paper of two rice genotypes grown at 4 weeks after transplanting.	128
4.1	Relative Mn uptake efficiency ((Mn uptake efficiency in Mn_0 , $Mn_{0.25}/Mn_{0.5}$) x 100) of KDML105 and PSL1 in 3 levels of Mn (0, 0.25	154

and 0.5 ppm; Mn₀, Mn_{0.25} and Mn_{0.5}, respectively) at 30 days after transplanting (a) and 60 day after transplanting (b). At Mn_{0.5}=100%.

- 5.1 Relative dry weight and TNC concentration in shoot and root at 30 days after transplanting. 174
- 5.2 Diagrammatic representation of Mn response modified from Gerloff (1977). 189



ลิขสิทธิ์มหาวิทยาลัยเชียงใหม่

Copyright© by Chiang Mai University

All rights reserved