# TABLE OF CONTENTS

	Page	
ACKNOWLEDGEMENTS	iii	
ABSTRACT (ENGLISH)	V	
ABSTRACT (THAI)	vii	
LIST OF TABLES	xii	
LIST OF FIGURES	xiv	
ABBREVIATIONS AND SYMBOLS	xvi	
CHAPTER 1: INTRODUCTION	1	
1.1 Rational and/or hypothesis		
1.2 Theory and literature review		
1.2.1 Aflatoxins	2	
1.2.1.1 Physical and chemical properties of aflatoxins	3	
1.2.1.2 Occurrence of aflatoxins	6	
1.2.1.3 Toxicity of aflatoxins	8	
1.2.1.4 Detoxification of aflatoxins	9	
00 vngh 1.2.2 Adsorption studies ang Mai Universit	15	
1.2.2.1 Adsorption isotherm	19	
1.2.2.2. Isotherm equations	21	
1.2.3 Determination of aflatoxins	24	
1.2.3.1 UV/VIS spectrophotometry	24	
1.2.3.2 High performance liquid chromatography	25	

	1.3 Research objectives	26
СНА	PTER 2: EXPERIMENTAL	28
	2.1 Instruments and apparatus	28
	2.2 Chemicals	28
	2.3 Preparations of the standard aflatoxin B <sub>1</sub>	29
	2.3.1 Stock standard aflatoxin B <sub>1</sub> solution	29
	2.3.2 Standard aflatoxin B <sub>1</sub> solution	29
	2.4 Preparations of other chemical solutions	30
	2.4.1 Potassium hydroxide solution 10%	30
	2.4.2 Sodium hypochlorite solution 1%	30
	2.5 Adsorption studies	30
	2.5.1 Construction of the aflatoxin $B_1$ calibration curves	30
	2.5.2 Study of adsorption isotherms	30
	2.5.3 Study on the temperature effect	31
	2.6 Isotherm fitting by 2D Table Curve program	31
	2.7 Characterizations of the adsorbents	32
	2.7.1 Determination of particle size	32
	2.7.2 Determination of total surface area and pore size	
	distribution S Person Construction	32
	2.7.3 Determination of total carbon content	32
	2.7.4 Determination of pH, conductivity and cation exchange	
	capacity (CEC)	33

Page

	Pag
CHAPTER 3: RESULTS AND DISCUSSION	34
3.1 Construction of the aflatoxin $B_1$ calibration curves	34
3.2 Study of adsorption isotherms	36
3.3 Study on the temperature effect	41
3.4 Isotherm fitting by 2D Table curve	46
3.5 Characterizations of the adsorbents	51
3.5.1 Determination of particle size	51
3.5.2 Determination of total surface area and pore size	
distribution	5 52
3.5.3 Determination of total carbon content	53
3.5.4 Determination of pH, conductivity and cation exchange	
capacity (CEC)	55
CHAPTER 4: CONCLUSIONS	57
THE RELEVANCE OF THE RESEARCH WORK TO THAILAND	60
REFERENCES	61
APPENDICES JM 19191311820	67
curriculum vitae by Chiang Mai Unive	

# LIST OF TABLES

Ta	ble of the	Page
1.1	Physical and chemical properties of aflatoxins	5
1.2	Molecular size of aflatoxins	5
1.3	Light absorption and fluorescence properties of aflatoxins	6
1.4	Characteristics associated with physical and chemical adsorption	18
3.1	Absorbance at 362 nm obtained from a series of standard aflatoxin $B_1$ solutions at 25 °C	35
3.2	Comparison fitting characteristics of isotherm data obtained from the	
	linearized Langmuir and Freundlich models at 25 °C	40
3.3	The linearized Langmuir and Freundlich fitting characteristics of	
	isotherm data obtained from adsorption of aflatoxin $B_1$ onto six	
	adsorbents at different temperatures	44
3.4	Isotherm equations used to fit the experimental data of a latoxin $B_1$	
	adsorption onto different adsorbents by 2D Table Curve program	47
3.5	User-defined used to fit in 2D Table Curve program	47
3.6	Fitting characteristics for the best-fit isotherms	49
3.7	The particle size of different adsorbents	51
3.8	The total surface area and pore size distribution of different adsorbents	52
3.9	The total carbon content of different adsorbents	54
3.1	0 The values of pH, conductivity and cation exchange capacity	
	(CEC) of different adsorbents	55

Table	Page
A1 Adsorption data of the commercial toxin binder at 25 $^{\circ}$ C	68
A2 Adsorption data of the commercial toxin binder at 37 °C	69
A3 Adsorption data of the commercial toxin binder at 45 °C	69
A4 Adsorption data of the commercial bentonite at 25 °C	70
A5 Adsorption data of the commercial bentonite at 37 °C	70
A6 Adsorption data of the commercial bentonite at 45 °C	71
A7 Adsorption data of activated carbon at 25 °C	71
A8 Adsorption data of activated carbon at 37 °C	72
A9 Adsorption data of activated carbon at 45 °C	72
A10 Adsorption data of the commercial zeolite at 25 °C	73
A11 Adsorption data of the commercial zeolite at 37 °C	73
A12 Adsorption data of the commercial zeolite at 45 °C	74
A13 Adsorption data of synthetic Na-X zeolite at 25 °C	74
A14 Adsorption data of synthetic Na-X zeolite at 37 °C	75
A15 Adsorption data of synthetic Na-X zeolite at 45 °C	75
A16 Adsorption data of synthetic sodalite zeolite at 25 °C	76
A17 Adsorption data of synthetic sodalite zeolite at 37 °C	76
A18 Adsorption data of synthetic sodalite zeolite at 45 °C	77 SITY

## LIST OF FIGURES

Fi	gure	Page
1.1	Chemical structures of aflatoxins	4
1.2	2 The six types of adsorption isotherms according to the classification of	
	IUPAC	19
1.3	3 Characteristics of linearized Langmuir isotherm plot	22
1.4	Characteristics of linearized Freundlich isotherm plot	24
3.1	Calibration curve of the standard aflatoxin $B_1$ solution at 25 °C	34
3.2	2 Adsorption isotherms for aflatoxin $B_1$ on six adsorbents at 25 °C	36
3.3	<sup>3</sup> The linearized Langmuir plots for aflatoxin $B_1$ adsorption onto: (a) the	
	commercial toxin binder, (b) the commercial bentonite, (c) activated	
	carbon, (d) the commercial zeolite, (e) synthetic Na-X zeolite, and (f)	
	synthetic sodalite zeolite at 25 °C	38
3.4	The linearized Freundlich plots for aflatoxin $B_1$ adsorption onto: (a) the	
	commercial toxin binder, (b) the commercial bentonite, (c) activated	
	carbon, (d) the commercial zeolite, (e) synthetic Na-X zeolite, and (f)	
	synthetic sodalite zeolite at 25 °C	39
3.5	Adsorption isotherms of aflatoxin $\mathbf{B}_1$ onto: (a) the commercial toxin	
	binder, (b) the commercial bentonite, (c) activated carbon, (d) the	
	commercial zeolite, (e) synthetic Na-X zeolite, and (f) synthetic	
	sodalite zeolite at different temperatures	42

### Figure

- 3.6 The best-fit isotherm for the adsorption of aflatoxin B<sub>1</sub> onto: (a) the commercial toxin binder, (b) the commercial bentonite, (c) activated carbon, (d) the commercial zeolite, (e) synthetic Na-X zeolite, and (f) synthetic sodalite zeolite
- B The character of adsorbents (a) The commercial toxin binder, (b) the commercial bentonite, (c) activated carbon, (d) the commercial zeolite,
  (e) synthetic Na-X zeolite, and (f) synthetic sodalite zeolite

**ลิขสิทธิ์มหาวิทยาลัยเชียงใหม่** Copyright<sup>©</sup> by Chiang Mai University All rights reserved

48

78

# ABBREVIATIONS AND SYMBOLS

Å	angstrom
С	concentration
°C	degree Celsius
CEC	cation exchange capacity
e.g.	for example
ELISA	enzymelinked immunosorbent assay
et. al.	and coworkers
°F	degree Fahrenheit
g	gram
GFM	generalized Freundlich model
$\Delta G$	Gibbs free energy
HPLC	high performance liquid chromatography
ΔH	enthalpy of adsorption
hr	hour
i.e.	that is han an a
kcal	kilocalorie Chiang Mai University
kg	kilogram
$K_d$	the distribution coefficient
$K_{f}$	Freundlich's constant related to adsorption capacity
LC	liquid chromatography
GLM	generalized Langmuir model

meq	miliequivalence
MFM	modified Freundlich model
mg	milligram
min	minute
mL	milliliter
mol	molar
MW	molecular weight
n	the heterogeneity factor
n <sup>f</sup>	the affinity constant
nm	nanometer
ppm	part per million (mg/kg)
ppb	part per billion (µg/kg)
P	pressure
$P^{o}$	normal vapour pressure
<i>Q</i> <sub>max</sub>	the estimated maximum capacity (mol/kg)
rpm	round per minute
$\Delta S$	entropy of adsorption
S.D	standard deviation
<b>T</b> vrigh	temperature by Chiang Mai University
TLC	thin layer chromatography
%	percentage n t S r e s e r v e o
μm	micrometer