

REFERENCES

- Adak, G. K., Long, S. M., and O'Brian, S. J. (2002). Trends in indigenous food borne disease and deaths, *England and Wales. Gut*, 51:832–841.
- Aguirre, M., and Collins, M.D. (1993). Lactic acid bacteria and human clinical infection, *J. Appl. Bacteriol.*, 75:95-107.
- Axelsson, L.T., Chung, T.C., Dobrogosz, W.J., Lindgren, S.E. (1989). Production of a broad spectrum antimicrobial substance by *Lactobacillus reuteri*. *Microbial Ecology of Health and Disease* 2:131–136.
- Axelsson, L.T. (1990). *Lactobacillus reuteri*, a member of the gut bacterial flora. Studies on antagonism, metabolism and genetics. Dissertation report 44. Uppsala, Sweden, University of Agriculture.
- Axelsson, L.T. (1993). Lactic Acid Bacteria: Classification and physiology. In S. Salminen, and A.V. Wright(eds). Lactic Acid Bacteria, pp. 1-64 New York: Marcel Dekker.
- Barefoot, S.F., Chen, Y.R., Hughes, T.A., Bodine, A.B., Shearer, M.Y. and Hughes, M.D. (1994). Identification and purification of a protein that induces production of the *Lactobacillus acidophilus* bacteriocin lactacin B, *Appl. Environ. Microbiol.*, 60:3522-3528.
- Benzet, R., Jung, G., Sahl, H.-G., (1991). Mechanism of channel-forming lantibiotics in black lipid membranes. In: Jung, G., Sahl, H.-G. (Eds.), Nisin and Novel Lantibiotics. ESCOM, Leiden, The Netherlands, pp. 359– 372.

- Betzl, D., Ludwig, W., and Schleifer, K.H. (1990) Identification of lactococci and enterococci by colony hybridization with 23S rRNA-targeted oligonucleotide probes, *Appl. Environ. Microbiol.*, 56:2927-2929.
- Bloom, H. and Mortvedt, C. (1991). Anti-microbial substances produced by food associated micro-organisms, *Biochem. Soc. Trans.*, 19:694-698.
- Brasil SP., (2004). Isolation of bacteriocin-producing LAB from meat and meat products and its spectrum of inhibitory activity, *J. Microbiol.*, 35:137-144
- Breukink, E., Weidemann, I., van Kraaij, C., Kuipers, O. P., Sahl, H.-G., and de Kruijff, B. (1999). Use of cell wall precursor lipid II by a poreforming peptide antibiotic. *Science*, 286:2361-2364.
- Brooks, J.L., Moore, A. S., Patchett, R. A., Collins, M.D., and Kroll, R.G. (1992). Use of the polymerase chain reaction and oligonucleotide probes of the rapid detection and identification of *Carnobacterium* species from meat, *J. Appl. Bacteriol.*, 72:249-301.
- Brosius J, Palmel JL, Kennedy JP, and Noller HF (1978) Complete nucleotide sequence of a 16S ribosomal RNA gene from *Escherichia coli*. Proc.NaOH Acad.Sci.USA,75:4801-4805
- Bruno, M.E.C. and Montville, T.J. (1993). Common mechanistic action of bacteriocin from lactic acid bacteria, *Appl. Environ. Microbiol.*, 59:3003-3010.
- Bryan-Jones, D.G., and Whittenbury, R. (1969). Haematin-dependent oxidative phosphorylation in *Streptococcus faecalis*, *J. Gen. Microbiol.*, 58:247-260.

Caplicee and Fitzgerald G.F. (1999). Food fermentations: role of microorganisms in food production and preservation. *Int. J. Food Microbiol.*, 50:131-149.

Carlsson, J., Iwami, Y. and Yamada, T. (1983). Hydrogen peroxide excretion by oral streptococci and effect of lactoperoxidase thiocyanate-hydrogen peroxide, *Infect. Immun.*, 40:70-80.

Chung, T.C., Alexsson, L., Lindgren, S.E., Dobrogosz, W.J., (1989). In vitro studies on reuterin synthesis in *Lactobacillus reuteri*. *Microbial Ecology of Health and Disease* 2: 137–141.

Chen, G. and Ressell, J.B. (1989). Transport of glutamine by *Streptococcus bovis* and conversion of glutamine to pyroglutamic acid and ammonia, *J. Bacteriol.*, 171:2981-2985.

Christensen, D.P. and Hutkins, R.W. (1992). Collaps of the proton motive force in *Listeria monocytogenes* caused by a bacteriocin produced by *Pediococcus acidilactici*, *Appl. Environ. Microbiol.*, 58:3312-3315.

Collins, M.D., Aquirre, M., Facklam, R.R., Shallcross, J., and Williams, A.M. (1992). *Gobiacatella sanguis* gen. nov., sp. nov., a new Gram-positive catalase-negative bacterium from human sources, *J. Appl. Bacteriol.*, 73:433-437.

Collins, M.D., Ash, C., Farrow, J.A.E., Wallbanks, S., and Williams, A.M. (1989). 16S ribosomal ribonucleic acid sequence analysis of lactococci and related taxa. Description of *Vagococcus fluvialis* gen. nov., sp. Nov., *J. Appl. Bacteriol.*, 67:453-460.

- Collins, M.D., Farrow, J.A.E., Philips, B.A., Ferusu, S., and Jones, D. (1987). Classification of *Lactobacillus divergens*, *Lactobacillus piscicola* and some catalase-negative, asporogenous, rod-shaped bacteria from poultry in a new genus, *Carnobacterium*, *Int. J. Syst. Bacteriol.*, 37:310-316.
- Collins, M.D., Williams, A.M., and Wallbanks, S. (1990). The phylogeny of *Aerococcus* and *pediococcus* as determined by 16S rRNA sequence analysis: description of *Tetragenococcus* gen. nov., *FEMS Microbiol. Lett.*, 70:255-262.
- Collins, M.D., Rodrigues, U., Ash, C., Aguirre, M., Farrow, J.A.E., Martinezmurcia, A., Phillips, B.A., Williams, A.M., and Wallbanks, S. (1991). Phylogenetic analysis of the genus *Lactobacillus* and related lactic acid bacteria as determined by reverse transcriptase sequencing of 16S rRNA, *FEMS Microbiol. Lett.*, 77:5-12.
- Collins, M.D., Samelis, J., Metaxopoulos, J., and Wallbanks, S. (1993). Taxonomic studies on some leuconostoc-like organisms from fermented sausages: description of a new genus *Weissella* for the *Leuconostoc parmesenteroides* group of species, *J. Appl. Bacteriol.*, 75:595-603.
- Condon, S. (1987). Responses of lactic acid bacteria to oxygen, *FEMS Microbiol. Rev.*, 46:269-280.
- Daeschel, M.A. (1989). Antimicrobial substances from lactic acid bacteria for use as food preservative. *Food Technology*, 43, 164-167.
- Delly, M., Mollet, B., and Hottinger, H. (1990). DNA probe for *Lactobacillus delbrueckii*, *Appl. Environ. Microbiol.*, 56:1967-1970.

De Vos, W.M., Kuipers, O.P., van der Meer, J.R. and Siezen, R.J. (1995). Maturation pathway of nisin and other lantibiotics: post-translationally modified antimicrobial peptides exported by Gram-positive bacteria, *Mol. Microbiol.*, 17:427-437.

De Vos, W.M., (1999). Gene expression systems for lactic acid bacteria. Current Opinion in *Microbiology* 2:289– 295.

Devriese, L.A., Pot, B., and Collins, M.D. (1993). Phenotypic identification of the genus *Enterococcus* and differentiation of phylogenetically distinct enterococcal species and species groups, *J. Appl. Bacteriol.* 75:399-408.

De Vuyst, L., Vandamme, E.J., (1994b). In: De Vuyst, L., Vandamme, E.J. (Eds.), Nisin, a lantibiotic produced by *Lactococcus lactis* subsp. *lactis*:properties, biosynthesis fermentation and application. Blackie Academic Press, Glasgow, pp. 151–222.

Dicks, L.M.T., Dellaglio, F., and Collins, M.D. (1995). Proposal to reclassify *Leuconostoc oenos* as *Oenococcus oeni* (corrig.). gen. nov., comb. Nov., *Int. J. Syst. Bacteriol.*, 45:395-397.

Dodd, H.M., Horn, N., Gasson, M.J., (1990). Analysis of the genetic determinants for the production of the peptide antibiotic nisin. *Journal of General Microbiology* 136:555– 566.

Donnelly, C. W. (2001). Listeria monocytogenes: A continuing challenge. *Nutrition Reviews*, 59:183–194.

Doores, S., (1993). In: Davidson, P.M, Branen, A.L. (Eds.), *Organic Acids*. Marcel Dekker, New York, pp. 95–136.

Dutton, C.J., Haxell, M.A., McAuthur, H.A.I, Wax, R.G., (2002). *Peptide Antibiotics, Discovery , Modes of Action and Applications*. Marcel Dekker, New York.

Ehrmann, M., Ludwig, W., and Schleifer, K.H. (1992). Species specific oligonucleotide probe for the identification of *Streptococcus thermophilus*, *Syst. Appl. Microbiol.*, 15:453-455.

Eklund, T. (1984). The effect of carbon dioxide on bacterial growth and uptake processes in the bacterial membrane vesicles. *International Journal of Food Microbiology*, 1:179–185.

Engesser, D.M., and Hammes, W.P. (1994). Non-heme catalase activity of lactic acid bacteria, *Syst. Appl. Microbiol.*, 17:11-19.

Erkkila S. and Petaja E. (2000). Screening of commercial meat starter cultures at low pH and in the presence of bile salts for potential probiotic use. *J. Meat Sci.* 55:297-300.

Evans, J.B. (1986). Genus *Aerococcus*, in *Bergey 's Manual of Systematic Bacteriology*, Vol. 2 (P.H.A. Sneath, N.S. Mair, M.E. Sharpe, and J.G. Holt,eds.), Williams and Wilkins, Baltimore, p. 1080.

Farber, J.M. (1991). Microbiological aspects of modified-atmosphere packaging technology—a review. *Journal of Food Protection*, 54:58–70.

Faruk A. (2002). Antimicrobial activity and characteristic of bacteriocins produced by Vaginal *Lactobacilli*. *J Med Sci.*,33:7-13.

Fontain, E.A., Claydon, E. and Tayler-Robinson, D. (1996). Lactobacilli from women with or without bacterial vaginosis and observations on the significance of hydrogen peroxide, *Microb. Ecol. Health Dis.*, 9:135-141.

Ganzle, M.G, Holtzel, A., Walter, J., Jung, G., Hammes, W.P. (2000). Characterization of reutericyclin produced by *Lactobacillus reuteri* LTH2584. *Applied and Environmental Microbiology* 66: 4233-4333.

Garvie, E.I. (1986b). Genus Pediococcus, in *Bergey's Manual of Systematic Bacteriology*, Vol. 2 (P.H.A. Sneath, N.S. Mair, M.E. Sharpe, and J.G. Holt, eds.), Williams and Wilkins, Baltimore, pp. 1075-1079.

Giovannoni, S.J., DeLong, E.F., Olsen, G.J., and Pace, N.R. (1988). Phylogenetic group-specific oligodeoxynucleotide probes for identification of single microbial cells, *J. Bacteriol.*, 170:720-726.

Gonzales, B., Glassker, R., Kunji, E.R.S., Driessens, A.J.M., Suarez, J.E. and Konings, W.N. (1996). Bactericidal mode of action of plantaricin C, *Appl. Environ. Microbiol.*, 62:2701-2709.

Grimont, F., and Grimont, P.A.D. (1992). Identification and typing by rRNA gene restriction patterns, in *Proceedings of the Conference on Taxonomy and Automated identification of Bacteria* (J. Schindler, ed.), Czechoslovak Society for Microbiology, Prague, pp.15-18.

Gross, E. and Morell, J.L. (1971). The structure of nisin, *J. Am. Chem. Soc.*, 93:4634-4635.

Hammes, W.P., and Vogel, R.F. (1995). The genus *Lactobacillus*, in *The Genera of Lactic Acid Bacteria* (B.J.B. Wood and W.H. Holzapfel, eds.), Chapman and Hall, London, pp.19-54.

Hardie, J.M. (1986a). Genus *Streptococcus* Rosenbach 1884, in *Bergey's Manual of Systematic Bacteriology*, Vol. 2 (P.H.A. Sneath, N.S. Mair, M.E. Sharpe, and J.G. Holt, eds.), Williams and Wilkins, Baltimore, pp. 1043-1047.

Harris, L.J., Fleming, H.P. and Klaenhammer, T.R. (1992). Developments in nisin research, *Fd. Res. Int.*, 25:57-66.

Hensiek, R., Krupp, G., and Stackebrandt, E. (1992). Development of diagnostic oligonucleotide probes for four *Lactobacillus* species occurring in the intestinal tract, *Syst. Appl. Microbiol.*, 15:123-128.

Hertel, C., Ludwig, W., Obst, M., Vogel, R.F., Hammes, W.P., and Schleifer, K.H. (1991). 23S rRNA-targeted oligonucleotide probes for the rapid identification of meat lactobacilli, *Syst. Appl. Microbiol.*, 14:173-177.

Holtzel, A., Ganzle, M.G., Nicholson, G.J., Hammes, W.P., Jung, G. (2000). The first low-molecular-weight antibiotic from lactic acid bacteria: reutericyclin, a new tetramic acid. *Angewandte Chemie. International Edition* 39:2766–2768.

Horn, N., Swindell, S., Dodd, H., Gasson, M.J., (1991). Nisin biosynthesis genes are encoded by a novel conjugative transposon. *Molecular and General Genetics* 228, 129– 135.

Huttunen, E., Noro, K. and Yang, Z. (1995). Purification and identification of antimicrobial substances produced by two *Lactobacillus casei* strains, *Int. Dairy J.*, 5:503-513.

Ingram, M. (1975). The lactic acid bacteria: a broad view, in *Lactic Acid Bacteria in Beverages and Food* (J.G. Carr, C.V. Cutting, and G.C. Whiting, eds.). Academic Press, London, pp. 1-13.

Immonen, Y., Ye, S., Ra, R., Qiao, M., Paulin, L., Saris, P.E.J., (1995). The codon usage of the nisin Z operon in *Lactococcus lactis* N8 suggests a non-lactococcal origin of the conjugative nisin-sucrose transposon. Sequence 5, 203– 218.

Jack, R.W., Tagg, J.R. and Ray, B. (1994). Bacteriocins of Gram-positive bacteria. *Microbiol. Rev.*, 59:171-200.

Jacop. F.L. woff, A., Siminovitch, A. and Wollman, E. (1953). Definition de quelques themes relatifs a la lysogenie, *Ann. Inst. Pasteur*, 84:222-224.

Janssen, P.H., Evers, S., Rainey, F.A., Weiss, N., Ludwig, W., Harfoot, C.G., and Schink, B. (1995). *Lactosphaera* gen. nov., a new genus of lactic acid bacteria, and transfer of *Ruminococcus pasteurii* Schink 1984 to *Lactosphaera pasteurii* comb. nov., *Int. J. Syst. Bacteriol.*, 45:565-571.

Javis, B. and Farr, J. (1971). Partial purification, specificity and mechanism of action of the nisin-inactivating enzyme from *Bacillus cereus*, *Biochim. Biophys. Acta*, 227:232-240.

Jay, J.M. (1982). Antimicrobial propertier of diacetyl. *J. Appl. Environ. Microbiol.* 44: 525.

- Jay, J.M. (1986). Modern food microbiology. New York: Van Nostrand Reinhold.
- Jimenez-Diaz, R., Rios-Sanchez, R.M., Desmazeaud, M., Ruiz-Barba, J.L. and Piard, J.C. (1993). Plantaricins S and T, two new bactriocins produced by *Lactobacillus plantarum* LPCO10 isolated from a green olive fermentation, *Appl. Environ. Microbiol.*, 59:1416-1424.
- Jimenez-Diaz, R., Ruiz-Barba, J.L., Cathcart, D.P., Holo, H., Nes, I.F., Sletten, K.H. and Warner, P.J. (1995). Purification and partial amino acid sequence of plantaricin S, a bacteriocin produced by *Lactobacillus plantarum* LPO10, the activity of which depends on the complementary action of two peptides, *Appl. Environ. Microbiol.*, 61:4459-4463.
- Johansson, M.L., Molin, G., Pettersson, B., Uhlen, M., and Ahrne, S. (1995a). Characterization and species recognition of *Lactobacillus plantarum* strains by restriction fragment length polymorphism (RFLP) of the 16S rRNA gene, *J. Appl. Bacteriol.*, 79:536-541.
- Kacem M. (1989). Lactic acid bacteria isolated from olives fermented in Western Algeria for possible use as probiotic. *Appl. Environ. Microbiol.*, 64:5216-5225.
- Kamau, D.N., Doores, S. and Pruitt, K.M. (1990). Enhanced thermal destruction of *Listeria monocytogenes* and *Staphylococcus aureus* by the lactoperoxidase system, *Appl. Environ. Microbiol.*, 56:2711-2716.
- Kandler, O. (1983). Carbohydrate metabolism in lactic acid bacteria, *Aatonie van Leeuwen-hoek*, 49:209-224.
- Kandler, O. (!984). Current taxonomy of lactobacilli, *Dev. Ind. Microbiol.*, 25:109-123.

Kandler, O., and Weiss, N. (1986). Regular, non-sporing Gram-positive rods, in *Bergey 's Manual of Systematic Bacteriology*, Vol. 2 (P.H.A. Sneath, N.S. Mair, M.E. Sharpe, and J.G. Holt,eds.), Williams and Wilkins, Baltimore, pp. 1208-1234.

Kelly, W.J., Asmundson, R.V., Huang, C.M., (1996b). Characterization of plantaricin KW30, a bacteriocin produced by *Lactobacillus plantarum*. *J. Appl.* 81:657-662.

Kim, S.H. and Worobo R.W., (2000). Characterization and Purification of a Bacteriocin Produced by a potential Probiotic Culture, *Lactobacillus acidophilus* 30SC. *J. Dairy Sci*, 83:2747-2752.

Klaenhammer, T. R. (1988). Bacteriocin of lactic acid bacteria. *Biochimic.* 70:337-349.

Klaenhammer, T.R. (1993). Genetics of bacteriocins produce by lactic acid bacteria. *FEMS Microbiology Review*, 12:39–86.

Klijn, N., Weerkamp, A. H., and de Vos, W.M. (1991). Identification of mesophilic lactic acid bacteria by using polymerase chain reaction-amplified variable regions of 16S rRNA and specific DNA probes, *Appl. Environ. Microbiol.*, 57:3390-3393.

Koneman, E.W., H.M. Sommers., S.D. Allen., and V.R. Dowell. (1983). Color Atlas and Textbook of Diagnostic Microbiology. 2th edition. *JB. Lippincott Company. Philadelphia.* P.425-458.

Kok, J., Holo, H., van Belkum, M.J., Haandrikman, A.J. and Nes, I.F. (1993). Non-nisin bacteriocins in lactococci: biochemistry, genetics and mode of action, in *Bacteriocins of Lactic Acid Bacteria* (D. Hoover and L. Steenson, eds.), Academic Press, New York, pp.121-150.

Kuipers, O.P., Yap, W.M.G.J., Rollemijn, H.S., Beerthuyzen, M.M., Siezen, R.J., De Vos, W.M., (1991). Expression of wild type and mutant nisin genes in *Lactococcus lactis*. In: Jung, G., Sahl, H.-G. (Eds.), Nisin and Novel Lantibiotics. *ESCOM Science Publishers, Leiden*, pp. 250–259.

Kuipers, O.P., Beerthuyzen, M.M., Siezen, R.J. and de Vos, W.M. (1993). Characterization of the nisin gene cluster *nisABTCIPR* of *Lactococcus lactis*, requirement of expression of the *nisA* and *nisI* genes for development of immunity, *Eur. J. Biochem.*, 216:281-291.

Kuipers, O.P., Beerthuyzen, M.M., de Ruyter, P.G.G.A., Luesink, E.J., de Vos, W.M., (1995). Autoregulation of nisin biosynthesis in *Lactococcus lactis* by signal transduction. *Journal of Biological Chemistry* 270:27299–27304.

Lancefield, R.C. (1993). A serological differentiation of human and other groups of hemolytic streptococci, *J. Exp. Med.*, 59:571-591.

Lane, D.J., Pace, B., Olsen, G.J., Stahl, D.A., Sogin, M.L., and Pace, N.R. (1985). Rapid determination of 16S ribosomal RNA sequences for phylogenetic analysis, *Proc. Natl. Acad. Sci. USA*, 82:6955:6959.

Lane, D.L., Field, K.G., Olsen, G.J., and Pace, N.R. (1988). Reverse transcriptase sequencing of ribosomal RNA for phylogenetic analysis, *Meth. Enzymol.*, 167:138-144.

Lindgren, S.E. and Dobrogosz, W.J. (1990). Antagonistic activities of lactic acid bacteria in food and feed fermentations, *FEMS Microbiol. Rev.*, 87:149-163.

Linnett, P.E. and Strominger, J.L. (1973). Additional antibiotic inhibitors of peptidoglycan synthesis. *Antimicrob. Agents Chemother.*, 4:231-236.

Martinez-Murcia, A.J., and Collins, M.D. (1990). A phylogenetic analysis of the genus Leucunostoc based on reverse transcriptase sequencing of 16S rRNA, *FEMS Microbiol. Lett.*, 70:73-84.

Meisel, J., Wolf, G., and Hammes, W.P. (1994). Heme-dependent cytochrome formation in *Lactobacillus maltaromicus*, *Syst. Appl. Microbiol.*, 17:20-23.

Moberg, L. (1989). Good manufacturing practices for refrigerated foods. *Journal of Food Protection*, 52:363-367.

Morris, J.G. (1976). Oxygen and the obligate anaerob, *J. Appl. Bact.*, 40:229-244.

Mulders, J.W.M., Boerrigter, I.J., Rollema, H.S., Siezen, R.J., De Vos, W.M., (1991). Identification and characterization of the lantibiotic nisin Z, a natural nisin variant. European Journal of Biochemistry 201:581– 584.

Mundt, J.O. (1986). Enterococci, in *Bergey's Manual of Systematic Bacteriology*, Vol. 2 (P. H. A. Sneath, N. S. Mair, M.E. Sharpe, and J.G. Holt, eds.), Williams and Wilkins, Baltimore, pp. 1063-1065.

Nagy, E., Petterson, M. and Mardh, P.A. (1991). Antibiosis between bacteria isolated from the vagina of women with and without signs of bacterial vaginosis, *Acta Pathol. Microbiol. Immunol. Scand.*, 99:739-744.

Nes, I.F., Diep, D.B., Havarstein, L.S., Brurberg, M.B., Eijsink, V. and Holo, H. (1996). Biosynthesis of bacteriocins in lactic acid bacteria, A.V. *Leeuwenhoek*, 70:113-128.

Nissen, H., Holck, A., and Dainty, R.H. (1994). Identification of *Carnobacterium spp.* and *Leuconostoc spp.* in meat by genus-specific 16S rRNA probes, *Lett. Appl. Microbiol.*, 19:165-168.

Nissen-Meyer, J., Havarstein, L.S., Holo, H., Sletten, K. and Nes, I.F. (1993). Association of the lactococcin A immunity factor with the cell membrane: purification and characterization of the immunity factor with, *J. Gen. Micobiol.*, 139:1503-1509.

Orla-Jensen, S. (1919). *The Lactic Acid Bacteria*, Host and son, Copenhagen. Piard, J.C., and Desmazeaud, M. (1991). Inhibiting factors produced by lactic acid bacteria: 1. Oxygen metabolites and catabolism endproduct. *Lait*, 71:525- 541.

Piard, J.C., and Desmazeaud, M. (1992). Inhibiting factors produced by lactic acid bacteria : 2. Bacteriocins and Other antibacterial substances. *Lait*, 72:113- 142.

Pot, B., Ludwig, W., Kersters, K., and Schleifer, K.H. (1994b). Taxonomy for lactic acid bacteria, in *Bacteriocins of Lactic Acid Bacteria* (L. De Vuyst and E.J. Vandamme, eds.), Chapman and Hall, London, pp.13-90.

Petrick, H.A.R., Ambrosio, R.E., and Holzapfel, W.H. (1988). Isolation of a DNA probe for *Lactobacillus curvatus*, *Appl. Environ. Microbiol.*, 54:405-408.

Pilloud, N., and Mollet, B. (1990). DNA probes for the detection of *Lactobacillus helveticus*, *Syst. Appl. Microbiol.*, 13:345-349.

Quadri, L.E.N., Sailer, M., Roy, K.L., Vederas, J.C. and Stiles, M.E. (1994). Chemical and genetic characterization of bacteriocins produced by *Carnobacterium piscicola* LV17B, *J. Biol. Chem.*, 269:12204-12211.

Quadri, L.E.N., Sailer, M., Terebiznik, M.R., Roy, K.L., Vederas, J.C. and Stiles, M.E. (1995). Characterization of the protein conferring immunity to the antimicrobial peptide carnobacteriocin B2 and expression of carnobacteriocins B2 and BMI, *J. Bacteriol.*, 177:1144-1151.

Rauch, P.J.G., and De Vos, W.M., (1992). Characterization of the novel nisin-sucrose conjugative transposon Tn5276 and its integration into *Lactococcus lactis*. *Journal of Bacteriology* 174:1280-1287.

Ray, B. (1993). Sublethal injury, bacteriocins, and food microbiology, *ASM News*, 59:285-291.

Ray, B., and Daeschel, M. (1992). Food Biopreservatives of Microbial Origin. CRC Press, Florida, USA.

Reddy, G.V. and Shahani, K.M. (1971). Isolation of antibiotic from *Lactobacillus bulgaricus*, *J. Dairy Sci.*, 54:748.

Reddy, G.V., Shahani, K.M., Friend, B.A. and Chandan, R.C. (1984). Natural antibiotic activity of *Lactobacillus acidophilus* and *bulgaricus* III : Production and partial purification of bulgarican from *Lactobacillus bulgaricus*. *Cult. Dairy Prod. J.*, 19:7-11.

Renata Bromberg. (2004). Isolation of bacteriocin-producing Lactic acid bacteria from meat and meat products and its spectrum of inhibitory activity. *Journal of Microbiology* 35: 137-144.

Resinger, P., Seidel, H., Tschesche, H. and Hammers, W.P. (1980). The effect of nisin on murein synthesis, *Arch. Microbiol.*, 127:187-193.

Ritchey, T.W., and Seeley, H.W.J. (1976). Distribution of cytochrome-like respiration in streptococci, *J. Gen. Microbiol.*, 93:195-203.

Rodrigues, U.M., Aguirre, M., Facklam, R.R., and Collins, M.D. (1991). Specific and intraspecific molecular typing of lactococci based on polymorphism of DNA encoding rRNA, *J. Appl. Bacteriol.*, 71:509-516.

Rodriguez, J.M.,and Dodd, H.M., (1996). Genetic determinants for the biosynthesis of nisin, a bacteriocin produced by *Lactococcus lactis*. *Microbiologia* 12:61–74.

Ruhr, E., Sahl, H.-G., (1985). Mode of action of the peptide antibiotic nisin and influence on the membrane potential of whole cells and on artificial membrane vesicles. *Antimicrobial Agents and Chemotherapy* 27:841–845.

Ryan. M.p., Rea, M.C., Hill, C. and Ross, R.P. (1996). An application in cheddar cheese manufacture for a strain of *Lactococcus lactis* producing a novel broad-spectrum bacteriocin, lacticin 3147, *Appl. Environ. Microbiol.*, 62:612-619.

Sahl. H.G., Jack, R.W. and Bierbaum, G. (1995). Biosynthesis and biological activities of lantibiotics with unique post-translational modifications, *Eur. J. Biochem.*, 230:827-853.

- Sahl, H.G., Kordel, M. and Benz, R. (1987). Voltage-dependent depolarization of bacterial membranes and artificial lipid bilayers by peptide antibiotic nisin, *Arch. Microbiol.*, 149:120-124.
- Sanders, M.E. (1993). Effects of consumption of lactic cultures on human health. *Adv. Fd. Nutr. Res.*, 37:67-130.
- Sambrook, J., Fritsch, E.F., and Maniatis, T. (1989). *Molecular Cloning: A Laboratory Manual*, 2nd ed. Ccold Spring Harbor Laboratory, Cold Spring Harbor, NY.
- Salama, M., Sandine, W., and Giovannoni, S. (1991). Development and application of oligonucleotide probes for identification of *Lactococcus lactis* subsp. *cremoris*, *Appl. Environ. Microbiol.*, 57:1313-1318.
- Schlegel, H.G. (1985). *Allgemeine Mikrobiologie*, Thieme, Stuttgart.
- Schleifer, K.H. (1986). Gram-positive cocci, in *Bergey's Manual of Systematic Bacteriology*, Vol. 2 (P.H.A. Sneath, N.S. Mair, M.E. Sharpe, and J.G. Holt, eds.), Williams and Wilkins, Baltimore, pp.999-1103.
- Schleifer, K.H., and Kilpper-Balz, R. (1984). Transfer of *Streptococcus faecalis* and *Streptococcus faecium* to the genus *Enterococcus* nom. rev. as *Enterococcus faecalis* comb. nov. and *Enterococcus faecium* comb. nov., *Int. J. Syst. Bacteriol.*, 34:31-34.
- Schleifer, K.H., and Kilpper-Balz, R. (1987). Molecular and hemotaxonomic approaches to the classification of streptococci, enterococci and lactococci: a review, *Syst. Appl. Microbiol.*, 10:1-19.

Schleifer, K.H., Kraus, J., Dvorak, C., Kilpper-Balz, R., Collins, M. D., and Fischer, W. (1985). Transfer of *Streptococcus lactis* and related streptococci to the genus *Lactococcus* gen. nov., *Syst. Appl. Microbiol.*, 6:183-195.

Schleifer, K.H., and Stackebrandt, E. (1983). Molecular systematics of prokaryotes, *Annu. Rev. Microbiol.*, 37:143-187.

Sharpe, M.E. (1979). Identification of the lactic acid bacteria, in *Identification Methods for Microbiologists*, 2nd ed., *Soc. Appl. Bacteriol. Technical Series*, No. 14 (F.A. Skinner and D.W. Lovelock, eds.), Academic Press, London, pp. 246-255.

Sharpe, M.E. (1981). The genus *Lactobacillus*, in *The Prokaryotes: A Handbook on Habitats, Isolation and Identification of Bacteria* (M.P. Starr, H. Stolp, H.G. Truper, A. Balows, and H.G. Schlegel, eds.), Springer-Verlag, Berlin, pp.1653-1674.

Shiba, T., Wakamiya, T., Fukase, K., Ueki, Y., Teshima, T., Nishikawa, M., (1991). Structure of the lanthionine peptide nisin, an ancovenin and lanthiopeptin. In: Jung, G., Sahl, H.-G. (Eds.), *Nisin and Novel Lantibiotics*. ESCOM Science Publishers, Leiden, pp. 113– 122.

Siegers, K. and Entain, K.D. (1995). Genes involved in immunity to the lantibiotic nisin produced by *Lactococcus lactis* 6F3, *Appl. Environ. Microbiol.*, 61:1082-1089.

All rights reserved

Silva. M., Jacobus, N.V., Deneke, C., and Gorbach, S.L. (1987). Antimicrobial substance from a human and Blackman G.E. (1994). The significance of hydrogen-ion concentration in the study of toxicity, *Symp. Soc. Exp. Biol.* 3:253–265.

Simon. E.W. and Blackman, G.E. (1949). The significance of hydrogen-ion concentration in the study of toxicity, *Symp. Soc. Exp. Biol.* 3:253 – 265.

Simpson, W.J. (1994). Comments on the mode of division of *Pediococcus spp.*, *Lett. Appl. Microbiol.*, 18:69-70.

Simpson, W.J., and Taguchi, H. (1995). The genus *Pediococcus* with notes on the genera *Tetragenococcus* and *Aerococcus*, in *The Genera of Lactic Acid Bacteria* (B.J.B. Wood and W.H. Holzapfel, eds.), Chapman and Hall, London, p. 125- 172.

Siragusa, G.R. (1992). Production of bacteriocin inhibitory to *Listeria* species by *Enterococcus hirae*, *Appl. Environ. Microbiol.*, 58:3508-3513.

Spillmam, H., Puhan, Z., and Banhegyi. M. (1987). Antimicrobielle aktivital thermophiler lactobacillen *Milchwissenschaft*, 33:148–153.

Stackebrandt, E., and Teuber, M. (1988). Molecular taxonomy and phylogenetic position of lactic acid bacteria, *Biochimie*, 70:317-324.

A Stahl, M., Pettersson, B., Molin, G., Uhlen, M., and Ahrne, S. (1994). Restriction fragment length polymorphism of *Lactobacillus reuteri* and *Lactobacillus fermentum*, originating from intestinal mucosa based on 16S rRNA genes, *Syst. Appl. Microbiol.*, 17:108-115.

Stevens, K.A., Sheldon, B.W., Klapes, N.A., Klaenhammer, T.R. (1992). Effective treatment conditions on nisin inactivation of Gram negative bacteria. *Journal of Food Protection* 55:763– 766.

Talarico, T.L., Dobrogosz, W.J. (1989). Chemical characterization of an antimicrobial substance produced by *Lactobacillus reuteri*. *Antimicrobial Agents and Chemotherapy* 33:674– 679.

Tamminen M. (2004). Screening of lactic acid bacteria from fermented vegetables by carbohydrate profiling and PCR-ELISA., *Applied Microbiology* 39:439-444.

Toba, T., Samant, S.K., Yoshioka, E. and Itoh, T. (1991). Reutericin 6, a new bacteriocin produced by *Lactococcus reuteri* LA 6, *Lett. Appl. Microbiol.*, 13:281-286.

Todorov, S.D., and Dicks, L.M.T. (2006). Screening for bacteriocin-producing lactic acid bacteria from boza, a traditional cereal beverage from Bulgaria Comparison of the bacteriocins. *Process Biochemistry* 41: 11-19.

Toit, M. (1998). Characterization and selection of probiotic Lactobacilli for a preliminary minipig feeding trial and their effect on serum cholesterol levels, faeces pH and faeces moistire content. *J. Food Microbiol.* 40:93-104.

Tortora, G.J., B.R. Funke. and C.L. Case. (1992). Microbiology. 4th edition., *Journal the Benjamin Cummings Publishing Company*. Inc. California. P. 672-676.

All rights reserved

Untereiner, S., (1989). Etude theorique da la fermentation du chou en choucroute.

Diplome de fin detudes, ENSIA-Massy, France.

van Belkum, M.J., Kok, J., Venema, G., Holo, H., Nes, I.F., Konings, W.N. and Abbe, T. (1991). The bacteriocin lactococcin A specifically increases the permeability of lactococcal cytoplasmic membranes in a voltage-independent, protein-mediated manner. *J. Bacteriol.*, 173:7934-7941.

van der Wal, F.J., Luijink, J. and Oudega, B. (1995). Bacteriocin release proteins: mode of action structure, and biotechnical application, *FEMS Microbiol. Rev.*, 17:381- 399.

van Niel, C.B., Kluyver, A.J. and Derx, H.G. (1929). Uber das Butteraroma, *Biochem. Z.*, 210:234-251.

Venema, K., Venema, G. and Kok, J. (1995). Lactococcal bacteriocins:mode of action and immunity, *Trends Microbiol.*, 3:299-303.

Vogel, R.F., Bocker, G., Stoltz, P., Ehrmann, M., Fanta, D., Ludwing, W., Pot, B., Kersters, K., Schleifer, K.H., and Hammes, W.P. (1994). Identification of lactobacilli from sourdough and description of *Lactobacillus pontis* sp. nov., *Int. J. Syst. Bacteriol.*, 44:223-229.

Wagner, M.K., and Moberg, L.J. (1989). Present and future use of traditional antimicrobials. *Food Technology*, 43:143–147.

Weisburge WG, Burns SM, Pelletier DA and Lane DJ (1991): 16S ribosomal DNA amplification for phylogenetic study. *J. Bacterial.* 143, 697-763

Whittenbury, R. (1964). Hydrogen peroxide formation and catalase activity in the lactic acid bacteria. *J. Gen. Microbiol.*, 35:13-26.

Whittenbury, R. (1978). Biochemical characteristics of *Streptococcus* species, in *Streptococci. Soc. Appl. Bacteriol. Symp. Series No. 7* (F.A. Skinner and L.B. Quesnel, eds.), Academic Press, London. Pp. 51-69.

Wiedemann, I., Breukink, E., van Kraaij, C., Kuipers, O., Bierbaum, G., de Kruijff, B., Sahl, H.-G., (2001). Specific binding of nisin to the peptidoglycan precursor lipid II combines pore formation and the inhibition of cell wall biosynthesis for potent antibiotic activity. *Journal of Biological Chemistry* 276:1772-1779.

Weiss, N. (1991). The genera *Pediococcus* and *Aerococcus*, in *The Prokaryotes*, Vol. 2, 2nd ed. (E. Balows, H.G. Truper, M. Davorkin, W. Harder, and K.-H. Schleifer, eds.), Springer-Verlag, New York, pp. 1502-1507.

Williams. A.M., and Collins, M.D. (1992). Genus-and species-specific oligonucleotide probes derived from 16S r RNA for the identification of vagococi., *Lett. Appl. Microbiol.*, 14:1-12.

Winkowski, k., Crandall, A.D. and Montville, T.J. (1993). Inhibition of *Listeria monocytogenes* by *Lactobacillus bavaricus* MN in beef systems at refrigeration temperatures, *Appl. Environ. Microbiol.*, 59:2552-2557.


Wolf, G., Strahl, A., Meisel, J., and Hammes, W.P. (1991). Heme-dependent catalase activity of lactobacilli, *Int. J. Food Microbiol.*, 12:133-140.

Wood, B.J.B., and Holzapfel, W.H. (eds.) (1995). *The Genera of Lactic Acid Bacteria*, Chapman and Hall. London.

Woese, C.R. (1987). Bacterial evolution, *Microbiol. Rev.*, 51:221-271.

Yang, D., and Woese, C.R. (1989). Phylogenetic structure of the “leuconostocs”: and interesting case of a rapidly evolving organism, *Syst. Appl. Microbiol.*, 12:145-149.

Yang, R., Johnson, M.C. and Ray, B. (1992). Novel method to extract large amounts of bacteriocins from lactic acid bacteria, *Appl. Microbiol.*, 58:3355-3359.



ลิขสิทธิ์มหาวิทยาลัยเชียงใหม่
Copyright[©] by Chiang Mai University
All rights reserved