

REFERENCES

- Alexopoulos, G. S. (2005). *Depression in the elderly*. Retrieved June 15, 2006 from
http://www.sciencedirect.com/science?_ob=JournalURL&_cdi=4886&_aut_h=y&_acct=C000010978&_version=1&_urlVersion=0&_userid=131617&md5=dac9b86a647bd8a89c28d8ff976d03c6.html
- American Psychiatric Association. (2000). Diagnostic and statistical manual of mental disorders (4th ed., text revision). Washington, DC: Author.
- Anderson, D. N. (2001). Treating depression in old age: The reasons to be positive. *Age and ageing*, 30(1), 13-17.
- Arean, P. A., & Cook, B. L. (2002). Psychotherapy and combined psychotherapy/pharmacotherapy for late life depression. *Biological Psychiatry*, 52(3), 293-303.
- Balletto, J., Hart, J. A., Lee, L., & Littman, A. (2003). *Depression*. Retrieved March 23, 2003, from <http://www.healthandage.com>
- Barker, P. (1999). Therapeutic nursing of the person in depression. In M. Clinton & S. Nelson (Eds.), *Advanced practice in mental health nursing*. Oxford: Blackwell Science.
- Barry, L. C., Allore, H. G., Guo, Z., Bruce, M. L., & Gill, T. M. (2008). Higher burden of depression among older women: The effect of onset, persistence, and mortality over time. *Archives of General Psychiatry*, 65(2), 172-178.
- Beck, A. (1967). *Depression clinical, experimental, and theoretical aspects*. New York: Harper & Row.

- Beck, A. T., & Steer, R. A. (1993). *Manual for the Beck depression inventory*. San Antonio, TX: The Psychological Corporation.
- Beck, A. T., Rush, A. J., Shaw, B. F., & Emery, G. (1979). *Cognitive therapy of depression*. New York: The Guilford Press.
- Beck, J. S. (1995). *Cognitive therapy: Basics and beyond*. New York: The Guilford Press.
- Bhikkhu Bodhi. (1999). *The noble eightfold path: The way to the end of suffering*. Retrieved August 9, 2007, from <http://www.accesstoinsight.org/lib/authors/bodhi.html>
- Blazer, D. (1998). *Emotional problems in later life: Intervention strategies for professional caregivers*. New York: Springer.
- Bradvik, L., & Berglund, M. (2001). Late mortality in severe depression. *Acta Psychiatrica Scandinavica*, 103, 111-116.
- Brage, D. G. (1995). Adolescent depression: A review of the literature. *Archives of Psychiatric Nursing*, 9(1), 45-55.
- Buddhadasa Bhikkhu. (1991). *Mindfulness with the breathing: A manual for serious beginners*. Bangkok: O.S Printing House.
- Buschmann, M. T., Dixon, M. A., & Tichy, A. M. (1995). Geriatric depression. *Home Healthcare Nurse*, 13(3), 47-56.
- Bush, J. H. (2004). Orientation to CBT. Retrieved June 27, 2004, from <http://www.Orientationcbt.html>
- Butler, A. C., & Beck, J. S. (2000). Cognitive therapy outcomes: A review of meta-analysis. *Journal of the Norwegian Psychological Association*, 37, 1-9.

- Callahan, C. M. (2001). Quality improvement research on late life depression in primary care. *Medical care*, 39(8), 772-784.
- Casacalenda, N., Perry, J. C., & Looper, K. (2002). Remission in major depressive disorder: A comparison of pharmacotherapy, psychotherapy, and control conditions. *American Journal of Psychiatry*, 159(8), 1354-1360.
- Casey D. (1994). Depression in the elderly. *Southern Medicine Journal*, 87, 559-563.
- Cassano, P., & Fava, M. (2002). Depression and public health: An overview. *Journal of Psychometric Research*, 53, 849-857.
- Castonguay, L. G., Goldfried, M. R., Wiser, S., Raue, P. J., & Hayes, A. M. (1996). Predicting the effect of cognitive therapy for depression: a study of unique and common factors. *Journal of Consulting and Clinical Psychology*, 64(3), 497-504
- Chantamoon, E. (1996). *The prevalence of dementia and major depression in elderly in Roi-Et Province*. Unpublished master's thesis, Chulalongkorn University, Bangkok, Thailand.
- Chatkaew, A. (2003). *Effect of cognitive therapy on depression of female youths in a welfare institution*. Unpublished master's thesis, Chiang Mai University, Chiang Mai, Thailand.
- Churujiporn, V. (1999). Coping styles of Thai depressive population. *Journal of Mental Health of Thailand*, 7(3), 176-179.
- Cole, M. D., & Dendukuri, N. (2003). Risk factors for depression among elderly community subjects: A systematic review and meta-analysis. *American Journal of Psychiatry*, 160(6), 1147-1156

- Day, P. H., & Deutsch, S. H. (2004). *Using mindfulness-based therapeutic interventions in psychiatric nursing practice-part I: Description and empirical support for mindfulness-based interventions*. Indianapolis: Indiana University School of Nursing.
- Department of Health and Human Services, Administration on Aging. (2001). *Older adults and mental health: Issues and opportunities*. Retrieved from <http://www.aoa.dhhs.gov/mh/report2001/default.html>
- Department of Mental Health. (2006). *Cumulative reports*. Retrieved December 20, 2007, from <http://www.dmh.moph.go.th/english.html>
- Disayavanish, C. (2005). *The four foundations of mindfulness*. Retrieved April 27, 2005, from <http://www.chamlongclinic-psych.com/mindfulness.html>
- Disayavanish, P. (1994). The effect of buddhist insight meditation on stress and anxiety. Unpublished doctoral dissertation, Illinois State University, Normal.
- Dwivedi, K. N. (2000). *Mindfulness in mental health*. Retrieved August 7, 2007, from <http://www.priory.com/psych/mindfulness.html>
- Evans, M., & Mottram, P. (2000). Diagnosis of depression in elderly patients. *Advances in Psychiatric Treatment*, 6, 49-56.
- Foongpaisarn, K. (1996). *Factors related to depression among the elderly people in municipal area of Kamphaeng Phet Province*. Unpublished master's thesis, Mahidol University, Bangkok, Thailand.
- Fram, D. H. (2006). *Depression in the elderly*. Retrieved June 5, 2008, from <http://www.webmd.com/depression/depression-elderly?page=2.html>

Franklin, D. J. (2002). *Depression-information and treatment*. Retrieved May 2, 2006, from <http://www.psychologyinfo.com/depression/index.html>

Frisch, N. C., & Frisch, L. E. (Eds.). (2002). *The client experiencing depression*. In *Psychiatric mental health nursing* (2nd ed., p. 245-273). New York: Delma.

Gareri, P., Fazio, P. D., & Sarro, G. D. (2002). Neuropharmacology of depression in aging and age-related diseases. *Ageing Research Reviews*, 1, 113-134.

Greenberger, D., & Padesky, C. A. (1995). *Mind over mood*. New York: The Guilford Press.

Gunaratana Mahathera, H. (2007). *Mindfulness in plain english*. Retrieved August 7, 2007, from http://www.vipassana.com/meditation/mindfulness_in_plain_english_19.php

Harvard Mental Health Letter. (2003). *Depression in old age*. Retrieved December 20, 2007, from http://findarticles.com/p/articles/mi_go2292/is_200309/ai_n9240068.html

Hillert, L., Hedman, K. B., Dolling, B. F., & Arnetz, B. B. (1998). Cognitive behavioral therapy for patients with electric sensitivity: A multidisciplinary approach in a controlled study. *Psychotherapy and Psychosomatic*, 67, 302-310.

Hollon, S. D., Thase, M. E., & Markowitz, J. C. (2002). Treatment and prevention of depression. *Psychological Science*. 3(2), 39-77.

Inskip, H. M., Harris, E. C., & Barraclough, B. (1998). Life time risk of suicide for affective disorder, alcoholism, and schizophrenia. *British Journal of Psychiatry*, 172, 35-37.

Jeffrey, S. H., & Charles, S. R. (2000). Removing the barriers to effective depression treatment in old age. *Journal of the American Geriatrics Society*. 48(8), 1012-1013.

Jeste, D. V. (2003). *Depression in older person*. Retrieved May 3, 2007, from http://www.nami.org/Template.cfm?Section=By_Illness&template=/ContentManagement/ContentDisplay.cfm&ContentID=7515.html

Kabat-Zinn, J. (1994). *Wherever you go, there you are: mindfulness in everyday life*. New York: Hyperion.

Katon, W, Lin, E., Russo, J., & Unutzer, J. (2003). Increase medical costs of a population-based sample of depressed elderly patients. *Archives General Psychiatry*, 60, 897-903.

Katz, I. R. (2004). Depression and frailty: The need for multidisciplinary research. *The American Journal of Geriatric Psychiatry*, 12, 1-6.

Keller, M. B. (2003). Past, present, and future direction for defining optimal treatment outcome in depression. *Journal of the American Medical Associate*, 289(23), 3152-3160.

Kelner, N. L., & Warren, B. J. (2003). Depression. In N. L. Kelner, L. H. Schwecke, & C. E. Bostrom (Eds.), *Psychiatric nursing*. (P. 340 - 375). St. Louis: Mosby.

Kessler, R. C. (2001). Epidemiology of women and depression. *Journal of Affective Disorders*, 74(1), 5-13.

Kitching, D., & Raphael, B. (2001). *Consensus guidelines for assessment and management of depression in the elderly*. Retrieved January 2, 2007, from <http://www.health.nsw.gov.au.html>

- Koder, D. A., Brodaty, H., & Anstey, K. J. (1996). Review cognitive therapy for depression in the elderly. *International Journal of Geriatric Psychiatry, 11*, 97-107.
- Koenig, H. G., & Blazer, D. G. (1996). Minor depression in late life. *American Journal of Geriatric Psychiatry, 4*(1), 14-21.
- Kraus, C. A., Kunik, M. E., & Stanley, M. A. (2007). *Use of cognitive behavioral therapy in late-life psychiatric disorders*. Retrieved January 2, 2007, from <http://www.geri.com/geriatrics/article.html>
- Kuehner, C. (2003). Gender differences in unipolar depression: An update of epidemiological findings and possible explanation. *Acta Psychiatrica Scandinavica, 108*, 163-174.
- Kunsongkeit, W. (2004). *Spiritual health of Thai people*. Unpublished doctoral dissertation, Chang Mai University, Thailand.
- Kwon, S. M., & Oei, T. P. S. (2003). Cognitive change process in a group cognitive behavior therapy of depression. *Journal of Behavior Therapy and Experimental Psychiatry, 34* (1), 73-85.
- Landreville, P., & Gervais, P. W. (1997). Psychotherapy for depression in older adults with a disability: Where do we go from here? *Aging & Mental Health, 1*(3), 197-209.
- Linsuwanont, P. (2001). *A study of elderly depression in Rongmuang Sub-district, Pratumwan, Bangkok Metropolis*. Unpublished master's thesis Mahidol University, Bangkok, Thailand.

- Ma, S. H., & Teasdale, J. D. (2004). Mindfulness-based cognitive therapy for depression: replication and exploration of differential relapse prevention effects. *Journal of Consulting and Clinical Psychology*, 72(1), 31-40.
- Maynard, C. K. (1993). Comparison of effectiveness of group interventions for depression in women. *Archives of Psychiatric Nursing*, 7(5), 277-283.
- Meurman, J. H., & Syvalahti, E. (2003). *Depressive symptoms in relation to oral health and related factors in a middle-aged population*. Retrieved December 12, 2005, from <http://herkules.oulu.fi/isbn9514270215.html>
- Miller, J., Fletcher, K., & Kabat-Zinn, J. (1995). Three - year follow - up and clinical implications of a mindfulness-based stress reduction intervention in the treatment anxiety disorders. *General Hospital Psychiatry*. 17. 192-200.
- Moutier, C., Wetherell, J. L., & Zisook, S. (2003). *Combined psychotherapy and pharmacotherapy for late-life depression*. Retrieved July 6, 2007, from <http://www.cmellc.com/geriatrictimes/q031014.html>
- Namsui, K. (2005). Effect of cognitive therapy program on depression of persons with HIV/AIDS. Unpublished master's thesis, Chiang Mai University, Chiang Mai, Thailand.
- Namto, S. S. (1989). *Moment to moment mindfulness*. FawnSkin: Vipassana dhura Meditation Society.
- National Alliance on Mental Illness. (2007). *Depression in older person*. Retrieved January 2, 2008, from http://www.nami.org/Template.cfm?Section =By_Illness&template=/ContentManagement/ContentDisplay.cfm&ContentID=7515.html

- National Institute of Mental Health. (2007). *Older adults: Depression and suicide facts*. Retrieved January 2, 2008, from <http://www.nimh.nih.gov/health/publications/older-adults-depression-and-suicide-facts.shtml>
- National Institutes of Health. (1998). *Late-life depression: An NIH consensus update*. Retrieved January 2, 2008, from <http://www.consensus.nih.gov/1991/1991DepressionLateLife086.html>
- Nunnally, J. C., & Bernstein, I. H. (1994). Psychometric Theory. In McGraw-Hill (Ed.), *Psychology* (3rd ed., p. 264-265). New York: McGraw-Hill.
- Nyanatiloka Thera. (2000). *The Buddha's path to deliverance* (5th ed.). Kandy: Buddhist Publication Society.
- O'Donohue, W. J., Fisher, J. E. & Hayes, S. C. (2003). *Cognitive behavior therapy: a guide for clinicians*. New York: Wiley.
- Palsson, S. P., Ostling, S., & Skoog, I. (2001). The incidence of first-onset depression in a population followed from the age of 70 to 85. *Psychological Medicine*, 31(7), 1159-1168.
- Phra Ajahn Plien Panyapatipo. (2006). *A Guide book to sati-sampajanna (mindfulness and self awareness*. (5th ed.). Chiang Mai: Silpa Siam Packging & Printing.
- Phra Bhasakorn Bhavilai. (2008). *Karma for today's traveler* (4th ed.). Chiang Mai: Nuntapun Printing.
- Phra Dhammadisuddhikavi. (2002). *A Buddhist way of mental training* (2nd ed.). Bangkok: Chuan Printing Press.
- Phra Rajaprommajarn. (2004). *Path to Nibbana* (complete edition). Chiang Mai: Chang Puerk.

Polit, D. F., & Beck, C. T. (2004). *Nursing research: Principal and methods* (7th ed.). Philadelphia: J.B. Lippincott.

Pritchard, C., & Baldwin, D. S. (2002). Elderly suicide rates in Asian and English-speaking countries. *Acta Psychiatrica Scandinavica*. 105 (4), 271-275.

Raj, A. (2004). *Depression in the elderly*. Retrieved December 18, 2007, from
<http://www.postgraduated.com>

Rupke, S. J., Blecke, D., & Renfrow, M. (2006). Cognitive therapy for depression. *Journal of the American Academy of Family Physicians*, 73(1), 30-35.

Rush, A. J., & Beck, A. T. (1995). Cognitive therapy. In H. I. Kaplan & B. J. Sadock (Eds.), *Comprehensive textbook of psychiatry*. (8 th ed., p. 2167- 2173) Baltimore: Williams and Wilkins.

Sacco, W. P., & Beck, A. T. (1995). Cognitive therapy and therapy. In E. E. Beckham & W. R. Leber (Eds.). *Handbook of depression* (2 nd ed., p.329-351). New York: Guilford Press.

Sadek, N., & Nemeroff, C. B. (2000). Update on the neurobiology of depression. Retrieved December 1, 2005, from <http://www.Medscape.com/Medscape/psychiatry/Treatment Update/2000/tu03/public/toc-tu03.html>.

Sanderson, W. C. & McGinn, L. K. (2001). Cognitive - behavioral therapy of depression. In M. M. Weissman & M. Weissman (Eds.). *Treatment of depression : Bridging the 21 st century*. New York: American Psychiatric Publishing Inc.

Sanseeha, L. (1993). *Depression and negative automatic thoughts of late adolescents*. Unpublished master's thesis, Chiang Mai University, Chiang Mai, Thailand.

- Scogin, F. (2007). *Depression and suicide in older adults resource guide*. Retrieved December 18, 2007, from <http://www.apa.org/pi/aging/depression.html>
- Segal, Z. V., Williams, J. M. G., & Teasdale, J. D. (2002). *Mindfulness-based cognitive therapy for depression*. New York: The Guilford Press.
- Serby, M., & Yu, M. (2003). Overview: Depression in the elderly. *The Mount Sinai Journal of Medicine*, 70(1), 38-44.
- Sidik, S. M., Zulkefli, A. M., & Shah, S. A. (2003). Factors associated with depression among elderly patients in a primary health care clinic in Malaysia. *Asia Pacific Family Medicine*, 2, 148-152.
- Siriwanarungsun, P., Kongsuk, T., Sririwan, Y., Charachsing, A., Bauolek, N., Rujirashakhorn, S., et al. (2003). *Quality of life and depression of Thai senior citizen in 2003*. Nonthaburi, Thailand: Department of Mental Health.
- Stauart, G. W. (2001). Emotional Responses and mood disorders. In G. W. Stauart, & M. T. Laraia, (Eds.). *Principles and practice of psychiatric nursing* (7th ed., p. 330-363). St. Louis. Mosby.
- Suan Prung Psychiatric Hospital. (2007). *Rate of psychiatric patients: Annual report*. Chiang Mai: Author.
- Teasdale, J. D., Segal, Z. V., Williams, J., Mark, G., Ridgeway, V. A., Soulsby, J. M., et al. (2000). Prevention of relapse/recurrence in major depression by mindfulness-base cognitive therapy. *Journal of Consulting & Clinical Psychology*, 68(4), 615-623.
- Thase, M. E., & Beck, A. T. (1993). *An overview of cognitive therapy*. USA.
- Thompson, L. W. (1996). Cognitive-behavioral therapy and treatment for late-life depression. *Journal of Clinical Psychiatry*, 57(5), 29-37.

- Thompson, L. W., Coon, D. W., Gallagher-Thomson, D., Sommer, B. R., & Koin, D. (2001). Comparison of desipramine and cognitive-behavioral therapy in the treatment of elderly outpatients with mild-to-moderate depression. *American Journal of Geriatric Psychiatry*, 9, 225-240.
- Varcarolis, E. M. (2002). *Foundations of psychiatric mental health nursing: A clinical approach* (4th ed.). Philadelphia: W.B. Saunders.
- Ven Punyananda Bhikkhu (n.d.). Why meditation?. In Mahachulalongkornrajavidyalaya Buddhist University. *The handbook experiencing thai Buddhist culture and meditation retreat*. Chiang Mai: Mahachulalongkornrajavidyalaya Press.
- Venerable Ajahn Sumedho. (1992). *The four noble truths*. Retrieved August 8, 2007 from <http://www.amaravati.org/abmnew/documents/4noble2/data/02pre.html>
- Walker, D. A., & Clarke, M. (2001). Cognitive behavioral psychotherapy: A comparison between younger and older adults in two inner city mental health teams. *Aging Mental Health*, 5(2), 197-199.
- Westbrook, D., Kennerley, H., & Kirk, J. (2007). *An introduction to cognitive behavioral therapy*. London: Sage.
- Wilson, K. C., Scott, M., Abou-Salch, M., Burns, R., & Copeland, J. R. (1995). Long-term effects of cognitive-behavioral therapy and lithium therapy on depression in the elderly. *The British Journal of Psychiatry*, 167, 653-658.
- World Health Organization. (2002). *Mental health: Responding to the call for action*. Retrieved December 19, 2003, from www.allaboutdepression.com

World Health Organization. (2003). *Depression*. Retrieved December 19, 2003, from
www.who.int

Yupho, D. (1988). *Vipassanā-Bhāvanā advance self-study practical insight meditation methods for self-testing and realization of consequences*. Bangkok: Mahachulalongkornrajavidyalaya Press.

Zerhuzen, J. D., Boyle, K., & Wilson, W. (1991). Out of darkness :Group cognitive therapy for depressed elderly. *Journal of Psychosocial of Mental Health Service*, 29(9), 16-21.

Zimney, E. (2008). *Definition of Depression*. Retrieved June 5, 2008, from
<http://www2.healthtalk.com/go/mental-health/depression/disease-basics.html>

จัดทำโดย ภาควิชาสังคมวิทยาและมานุษยวิทยา

Copyright © by Chiang Mai University
All rights reserved