

REFERENCES

- Akrishnan VR, Menon VP. Potential role of antioxidants during ethanol-induced changes in the fatty acid composition and arachidonic acid metabolites in male Wistar rats. *Cell Biol Toxicol.* 2001; 17(1): 11-22.
- Ali AT., Crowther NJ. Body fat distribution and insulin resistance. *S. Afr. Med. J.* 2005; 95: 878-880.
- Ammon HP., Wahl MA. Pharmacology of *Curcuma longa*. *Planta Med.* 1991; 57: 1-7.
- Araujo CC., Leon LL. Biological activities of *Curcuma longa* L. *Mem. Inst. Oswaldo Cruz.* 2001; 96: 723-728.
- Arun N., Nalini N. Efficacy of turmeric on blood sugar and polyol pathway in diabetic albino rats. *Plant Foods Hum. Nutr.* 2002; 57: 41-52.
- Asai A., Miyazawa T. Dietary curcuminoids prevent high-fat diet-induced lipid accumulation in rat liver and epididymal adipose tissue. *J Nutr.* 2001; 131(11): 2932-5.
- Atlanta GA.: Centers for Disease control. 2005. "Prevalence of overweight and obesity among adults: United States 1985–2003". [Online]. Available: <http://www.cdc.gov/needphp/dnpa/obesity/trend/maps>

- Babu PS, Srinivasan K. Hypolipidemic action of curcumin, the active principle of turmeric (*Curcuma longa*) in streptozotocin induced diabetic rats. *Mol Cell Biochem.* 1997; 166(1-2): 169-75.
- Bergman RN., Ader M. Free fatty acids and pathogenesis of type 2 diabetes mellitus. *Trends Endocrinol. Metab.* 2000; 11: 351-356.
- Bevilacqua S., Buzzigoli G., Bonadonna R., Brandi LS., Oleggini M., Boni C., Geloni M., Ferrannini E. Operation of Randle's cycle in patients with NIDDM. *Diabetes.* 1990; 39: 383-389.
- Bierhous A. The dietary pigment curcumin reduces endothelial tissue factor gene expression by inhibiting binding of AP-1 to the DNA and activation of NF κ B . *Thromb Haematostasis.* 1997; 77: 772-82.
- Bigger JT., Jr., Fleiss JL., Rolnitzky LM., Steinman RC. Frequency domain measures of heart period variability to assess risk late after myocardial infarction. *J. Am. Coll. Cardiol.* 1993; 21: 729-736.
- Bjorntorp P. Obesity. *Lancet.* 1997; 350: 423-426.
- Bjorntorp P., Bergman H., Varnauskas E. Plasma free fatty acid turnover rate in obesity. *Acta Med. Scand.* 1969; 185: 351-356.
- Bloomfield DM., Kaufman ES., Bigger JT., Jr., Fleiss J., Rolnitzky L., Steinman R. Passive head-up tilt and actively standing up produce similar overall changes in autonomic balance. *Am. Heart J.* 1997; 134: 316-320.
- Blunberg VS. "Obesity and the heart". In: Bjorntrop P, Brodoff BN, eds. *Obesity.* Philadelphia: Lippincott; pp. 517-531, 1992.

Boden G. Role of fatty acids in the pathogenesis of insulin resistance and NIDDM. *Diabetes*. 1997; 46: 3-10.

Boden G. Effects of free fatty acids on gluconeogenesis and glycogenolysis. *Life Sci*. 2003; 72: 977-988.

Boden G., Chen X. Effects of fat on glucose uptake and utilization in patients with non-insulin-dependent diabetes. *J Clin Invest*. 1995; 96: 1261-1268.

Boden G., Chen X., Ruiz J., White JV., Rossetti L. Mechanisms of fatty acid-induced inhibition of glucose uptake. *J Clin Invest*. 1994; 93: 2438-2446.

Bonnemeier H., Richardt G., Potratz J., Wiegand UK., Brandes A., Kluge N., Katus HA. Circadian profile of cardiac autonomic nervous modulation in healthy subjects: differing effects of aging and gender on heart rate variability. *J Cardiovasc Electrophysiol*. 2003; 14: 791-799.

Bonora E., Kiechl S., Willeit J., Oberhollenzer F., Egger G., Targher G., Alberiche M., Bonadonna RC., Muggeo M. Prevalence of insulin resistance in metabolic disorders: the Bruneck Study. *Diabetes*. 1998; 47: 1643-1649.

Borst SE., Conover CF. High-fat diet induces increased tissue expression of TNF-alpha. *Life Sci*. 2005; 77: 2156-2165.

Bray GA., York DA., Fisler JS. Experimental obesity: a homeostatic failure due to defective nutrient stimulation of the sympathetic nervous system. *Vitam Horm*. 1989; 45: 1-125.

- Brouet I., Ohshima H. Curcumin, an anti-tumour promoter and anti-inflammatory agent, inhibits induction of nitric oxide synthase in activated macrophages. *Biochem Biophys Res Commun.* 1995; 206(2): 533-40.
- Centers for Disease Control and Prevention. "Obesity Trends 1984– 2002". U.S.
- Chalkley SM., Hettiarachchi M., Chisholm DJ., Kraegen EW. Long-term high-fat feeding leads to severe insulin resistance but not diabetes in Wistar rats. *Am J Physiol Endocrinol Metab.* 2002; 282: E1231-E1238.
- Chattopadhyay I., Biswas K., Bandyopadhyay U., Banerjee R.K. Turmeric and curcumin : biological actions and medicinal applications. *Current Science.* 2004; 48(1): 44-53.
- Chearwae W., Wu C.P., Chu H.Y., T. Lee R., Ambudkar SV., Limtrakul P. Curcuminoids purified from turmeric powder modulate the function of human multidrug resistance protein 1 (ABCC1). *Cancer Chemotherapy and Pharmacology.* 2006; 57: 376-388.
- Chen HW., Huang HC. Effect of curcumin on cell cycle progression and apoptosis in vascular smooth muscle cells. *Br J Pharmacol.* 1998; 124(6): 1029-40.
- Chen X., Iqbal N., Boden G. The effects of free fatty acids on gluconeogenesis and glycogenolysis in normal subjects. *J Clin Invest.* 1999; 103: 365-372.
- Clore JN., Glickman PS., Nestler JE., Blackard WG. In vivo evidence for hepatic autoregulation during FFA-stimulated gluconeogenesis in normal humans. *Am J Physiol.* 1991; 261: E425-E429.

- Corbi GM., Carbone S., Ziccardi P., Giugliano G., Marfella R., Nappo F., Paolisso G., Esposito K., Giugliano D. FFAs and QT intervals in obese women with visceral adiposity: effects of sustained weight loss over 1 year. *J Clin Endocrinol Metab* 2002; 87: 2080-2083.
- Crespin SR., Greenough WB., III, Steinberg D. Stimulation of insulin secretion by long-chain free fatty acids. A direct pancreatic effect. *J Clin Invest.* 1973; 52: 1979-1984.
- Cripps TR., Malik M., Farrell TG., Camm AJ. Prognostic value of reduced heart rate variability after myocardial infarction: clinical evaluation of a new analysis method. *Br Heart J.* 1991; 65(1): 14-9.
- DeFronzo RA., Ferrannini E. Insulin resistance: A multifaceted syndrome responsible for NIDDM, obesity, hypertension, dyslipidemia, and atherosclerotic cardiovascular disease. *Diabetes Care.* 1991; 14: 173-194.
- Dekker JM., Schouten EG., Klootwijk P., Pool J., Swenne CA., Kromhout D. Heart rate variability from short electrocardiographic recordings predicts mortality from all causes in middle-aged and elderly men. The Zutphen Study. *Am J Epidemiol.* 1997; 145: 899-908.
- Deshpanda SS., Maru G.B. Effects of curcumin on the formation of benzo[a] pyrene derived DNA adducts in vitro. *Cancer Lett.* 1995; 96: 71-80.
- Dhalla NS., Liu X., Panagia V., Takeda N. Subcellular remodeling and heart dysfunction in chronic diabetes. *Cardiovasc Res.* 1998; 40: 239-247.
- Eckel RH., Grundy SM., Zimmet PZ. The metabolic syndrome. *Lancet.* 2005; 365: 1415-1428.

El-Gamal A., Gallagher D., Nawras A., Gandhi P., Gomez J., Allison DB., Steinberg JS., Shumacher D., Blank R., Heymsfield SB. Effects of obesity on QT, RR, and QTc intervals. *Am J Cardiol.* 1995; 75: 956-959.

Fallen EL., Kamath MV., Ghista DN. Power spectrum of heart rate variability: a non-invasive test of integrated neurocardiac function. *Clin Invest Med.* 1988; 11: 331-340.

Fanelli C., Calderone S., Epifano L., De VA., Modarelli F., Pampanelli S., Perriello G., De FP., Brunetti P., Gerich JE. Demonstration of a critical role for free fatty acids in mediating counterregulatory stimulation of gluconeogenesis and suppression of glucose utilization in humans. *J Clin Invest.* 1993; 92: 1617-1622.

Frayn KN., Maycock PF. Skeletal muscle triacylglycerol in the rat: methods for sampling and measurement, and studies of biological variability. *J Lipid Res.* 1980; 21: 139-144.

Gao YY., Lovejoy JC., Sparti A., Bray GA., Keys LK., Partington C. Autonomic activity assessed by heart rate spectral analysis varies with fat distribution in obese women. *Obes Res.* 1996; 4: 55-63.

Goberna R., Tamarit J., Jr., Osorio J., Fussganger R., Tamarit J., Pfeiffer EF. Action of B-hydroxy butyrate, acetoacetate and palmitate on the insulin release in the perfused isolated rat pancreas. *Horm Metab Res.* 1974; 6: 256-260.

Goldstein BJ. Insulin resistance as the core defect in type 2 diabetes mellitus. *Am J Cardiol.* 2002; 90: 3G-10G.

Gonzalez-Manchon C., Ayuso MS., Parrilla R. Control of hepatic gluconeogenesis: role of fatty acid oxidation. *Arch Biochem Biophys.* 1989; 271: 1-9.

Grassi G., Seravalle G., Cattaneo BM., Bolla GB., Lanfranchi A., Colombo M., Giannattasio C., Brunani A., Cavagnini F., Mancia G. Sympathetic activation in obese normotensive subjects. *Hypertension.* 1995; 25: 560-563.

Han DH., Hansen PA., Host HH., Holloszy JO. Insulin resistance of muscle glucose transport in rats fed a high-fat diet: a reevaluation. *Diabetes.* 1997; 46: 1761-1767.

Huang HC., Jan TR., Yeh SF. Inhibitory effect of curcumin, an anti-inflammatory agent, on vascular smooth muscle cell proliferation. *Eur J Pharmacol.* 1992; 221(2-3): 381-4.

Huang MT., Ma W., Lu YP., Chang RL., Fisher C., Manchand PS., Newmark HL., Conney AH. Effects of curcumin, demethoxycurcumin, bisdemethoxycurcumin and tetrahydrocurcumin on 12-O-tetradecanoylphorbol-13-acetate-induced tumor promotion. *Carcinogenesis.* 1995; 16(10): 2493-7.

Hubert HB., Feinleib M., McNamara PM., Castelli WP. Obesity as an independent risk factor for cardiovascular disease: a 26-year follow-up of participants in the Framingham Heart Study. *Circulation.* 1983; 67: 968-977.

Jacob S., Machann J., Rett K., Brechtel K., Volk A., Renn W., Maerker E., Matthaei S., Schick F., Claussen CD., Haring HU. Association of increased intramyocellular lipid content with insulin resistance in lean nondiabetic offspring of type 2 diabetic subjects. *Diabetes*. 1999; 48: 1113-1119.

Jensen MD., Haymond MW., Rizza RA., Cryer PE., Miles JM. Influence of body fat distribution on free fatty acid metabolism in obesity. *J Clin Invest*. 1989; 83: 1168-1173.

Jones PP., Snitker S., Skinner JS., Ravussin E. Gender differences in muscle sympathetic nerve activity: effect of body fat distribution. *Am J Physiol*. 1996; 270: E363-E366.

Jouven X., Charles MA., Desnos M., Ducimetiere P. Circulating nonesterified fatty acid level as a predictive risk factor for sudden death in the population. *Circulation*. 2001; 104: 756-761.

Kahn BB., Flier JS. Obesity and insulin resistance. *J Clin Invest*. 2000; 106: 473-481.

Kahn SE., Prigeon RL., Schwartz RS., Fujimoto WY., Knopp RH., Brunzell JD., Porte D., Jr. Obesity, body fat distribution, insulin sensitivity and Islet beta-cell function as explanations for metabolic diversity. *J Nutr*. 2001; 131: 354S-360S.

Karason K., Molgaard H., Wikstrand J., Sjostrom L. Heart rate variability in obesity and the effect of weight loss. *Am J Cardiol*. 1999; 83: 1242-1247.

- Kassab S., Kato T., Wilkins FC., Chen R., Hall JE., Granger JP. Renal denervation attenuates the sodium retention and hypertension associated with obesity. *Hypertension*. 1995; 25: 893-897.
- Kim JA., Park YG., Cho KH., Hong MH., Han HC., Choi YS., Yoon D. Heart rate variability and obesity indices: emphasis on the response to noise and standing. *J Am Board Fam Pract*. 2005; 18: 97-103.
- Kim JK., Fillmore JJ., Chen Y., Yu C., Moore IK., Pypaert M., Lutz EP., Kako Y., Velez-Carrasco W., Goldberg IJ., Breslow JL., Shulman GI. Tissue-specific overexpression of lipoprotein lipase causes tissue-specific insulin resistance. *Proc Natl Acad Sci U. S. A.* 2001; 98: 7522-7527.
- Kim JY. High-fat diet-induced muscle insulin resistance: relationship to visceral fat mass. *Am.J.Physiol Regul.Integr.Comp Physio*. 2000; 279.6: R2057-R2065.
- Kim SP., Ellmerer M., Van Citters GW., Bergman RN. Primacy of hepatic insulin resistance in the development of the metabolic syndrome induced by an isocaloric moderate-fat diet in the dog. *Diabetes*. 2003; 52: 2453-2460.
- Kleiger RE., Miller JP., Bigger JT., Jr., Moss AJ. Decreased heart rate variability and its association with increased mortality after acute myocardial infarction. *Am J Cardiol*. 1987; 59: 256-262.
- Kleiger RE., Stein PK., Bigger JT., Jr. Heart rate variability: measurement and clinical utility. *Ann. Noninvasive. Electrocardiol*. 2005; 10: 88-101.
- Kleiger RE., Stein PK., Bosner MS., Rottman JN. Time domain measurements of heart rate variability. *Cardiol Clin*. 1992; 10: 487-498.

Kopelman PG. Obesity as a medical problem. *Nature*. 2000; 404: 635-643.

Koyama K., Chen G., Lee Y., Unger RH. Tissue triglycerides, insulin resistance, and insulin production: implications for hyperinsulinemia of obesity. *Am J Physiol*. 1997; 273: E708-E713.

Kraegen EW., Clark PW., Jenkins AB., Daley EA., Chisholm DJ., Storlien LH. Development of muscle insulin resistance after liver insulin resistance in high-fat-fed rats. *Diabetes*. 1991; 40: 1397-1403.

Kramer D., Shapiro R., Adler A., Bush E., Rondinone CM. Insulin-sensitizing effect of rosiglitazone (BRL-49653) by regulation of glucose transporters in muscle and fat of Zucker rats. *Metabolism*. 2001; 50: 1294-1300.

Laederach-Hofmann K., Mussgay L., Ruddel H. Autonomic cardiovascular regulation in obesity. *J Endocrinol*. 2000; 164: 59-66.

Lamarche B. Abdominal obesity and its metabolic complications: implications for the risk of ischaemic heart disease. *Coron. Artery Dis*. 1998; 9: 473-481.

Langager AM., Hammerberg BE., Rotella DL., Staus HM. Very low-frequency blood pressure variability depends on voltage-gated L-type Ca^{2+} channels in conscious rats. *Am J Physiol Heart Circ Physiol*. 2007; 292: H1321-H1327.

Lemieux S., Prud'homme D., Tremblay A., Bouchard C., Despres JP. Anthropometric correlates to changes in visceral adipose tissue over 7 years in women. *Int J Obes Relat Metab Disord*. 1996; 20: 618-624.

Maheshwari RK., Singh AK., Gaddipati J., Srimal RC. Multiple biological activities of curcumin: a short review. *Life Sci.* 2006; 78: 2081-2087.

Malaisse WJ., Malaisse-Lagae F. Stimulation of insulin secretion by noncarbohydrate metabolites. *J Lab Clin Med.* 1968; 72: 438-448.

Malik M. Heart rate variability. *Curr Opin Cardiol.* 1998; 13: 36-44.

Malik M. "Measurement of heart rate variability". In: Malik M, Camm J, eds. *Heart Rate Variability*. Armonk, NY: Futura, pp. 33-132, 1995.

Malliani A., Pagani M., Lombardi F., Cerutti S. Cardiovascular neural regulation explored in the frequency domain. *Circulation.* 1991; 84: 482-492.

Manosontorn S. Attributable Fraction of Obesity in Thailand. *Journal of Health Science.* 2005; 14,(2): 337-344.

Manzella D., Barbieri M., Ragno E., Paolisso G. Chronic administration of pharmacologic doses of vitamin E improves the cardiac autonomic nervous system in patients with type 2 diabetes. *Am J Clin Nutr.* 2001; 73: 1052-1057.

Manzella D., Barbieri M., Rizzo MR., Ragno E., Passariello N., Gambardella A., Marfella R., Giugliano D., Paolisso G. Role of free fatty acids on cardiac autonomic nervous system in noninsulin-dependent diabetic patients: effects of metabolic control. *J Clin Endocrinol Metab.* 2001; 86: 2769-2774.

Manzella D., Grella R., Marfella R., Giugliano D., Paolisso G. Elevated post-prandial free fatty acids are associated with cardiac sympathetic overactivity in Type II diabetic patients. *Diabetologia.* 2002; 45: 1737-1738.

- Martini G., Riva P., Rabbia F., Molini V., Ferrero GB., Cerutti F. Heart rate variability in childhood obesity. *Clin Auton Res.* 2001; 11: 87-91.
- Massillon D., Barzilai N., Hawkins M., Prus-Wertheimer D., Rossetti L. Induction of hepatic glucose-6-phosphatase gene expression by lipid infusion. *Diabetes.* 1997; 46: 153-157.
- Matsumoto T., Miyawaki C., Ue H., Kanda T., Yoshitake Y., Moritani T. Comparison of thermogenic sympathetic response to food intake between obese and non-obese young women. *Obes Res.* 2001; 9: 78-85.
- Matsumoto T., Miyawaki T., Ue H., Kanda T., Zenji C., Moritani T. Autonomic responsiveness to acute cold exposure in obese and non-obese young women. *Int J Obes Relat Metab Disord.* 1999; 23: 793-800.
- Matthews DR., Hosker JP., Rudenski AS., Naylor BA., Treacher DF., Turner RC. Homeostasis model assessment: insulin resistance and beta-cell function from fasting plasma glucose and insulin concentrations in man. *Diabetologia.* 1985; 28(7): 412-9.
- McGuinness OP, Ejiofor J, Audoly LP, Schrom N. Regulation of glucose production by NEFA and gluconeogenic precursors during chronic glucagon infusion. *Am J Physiol.* 1998; 275(3 Pt 1): E432-9.
- Messerli FH., Nunez BD., Ventura HO., Snyder DW. Overweight and sudden death. Increased ventricular ectopy in cardiopathy of obesity. *Arch Intern Med.* 1987; 147: 1725-1728.

Morgan DA., Anderson EA., Mark AL. Renal sympathetic nerve activity is increased in obese Zucker rats. *Hypertension*. 1995; 25: 834-838.

Moss AJ. Measurement of the QT interval and the risk associated with QTc interval prolongation: a review. *Am J Cardiol*. 1993; 72: 23B-25B.

Muscelli E., Emdin M., Natali A., Pratali L., Camastra S., Gastaldelli A., Baldi S., Carpeggiani C., Ferrannini E. Autonomic and hemodynamic responses to insulin in lean and obese humans. *J Clin Endocrinol Metab*. 1998; 83: 2084-2090.

Must A., Spadano J., Coakley EH., Field AE., Colditz G., Dietz WH. The disease burden associated with overweight and obesity. *JAMA*. 1999; 282: 1523-1529.

Nieves DJ., Cnop M., Retzlaff B., Walden CE., Brunzell JD., Knopp RH., Kahn SE. The atherogenic lipoprotein profile associated with obesity and insulin resistance is largely attributable to intra-abdominal fat. *Diabetes*. 2003; 52: 172-179.

Nirmala C., Puvanakrishnan R. Protective role of curcumin against isoproterenol induced myocardial infarction in rats. *Mol Cell Biochem*. 1996; 159(2): 85-93.

Nishiyama T., Mae T., Kishida H., Tsukagawa M., Mimaki Y., Kuroda M., Sashida Y., Takahashi K., Kawada T., Nakagawa K., Kitahara M. Curcuminoids and sesquiterpenoids in turmeric (*Curcuma longa* L.) suppress an increase in blood glucose level in type 2 diabetic KK-Ay mice. *J Agric Food Chem*. 2005; 53(4): 959-63.

- Ogden CL., Carroll MD., Curtin LR., McDowell MA., Tabak CJ., Flegal KM. Prevalence of overweight and obesity in the United States, 1999-2004. *JAMA* 2006; 295: 1549-1555.
- Oliver MF., Opie LH. Effects of glucose and fatty acids on myocardial ischaemia and arrhythmias. *Lancet*. 1994; 343: 155-158.
- Olszanecki R., Jawien J., Gajda M., Mateuszuk L., Gebaska A., Korabiowska M., Chlopicki S., Korbut R. Effect of curcumin on atherosclerosis in apoE/LDLR-double knockout mice. *J Physiol and Pharmacol*. 2005; 56(4): 627-35.
- Oparil S., Oberman A. Nontraditional cardiovascular risk factors. *Am J Med Sci*. 1999; 317: 193-207.
- Pagani M., Lombardi F., Guzzetti S., Rimoldi O., Furlan R., Pizzinelli P., Sandrone G., Malfatto G., Dell'Orto S., Piccaluga E. Power spectral analysis of heart rate and arterial pressure variabilities as a marker of sympatho-vagal interaction in man and conscious dog. *Circ Res*. 1986; 59: 178-193.
- Pan DA., Lillioja S., Kriketos AD., Milner MR., Baur LA., Bogardus C., Jenkins AB., Storlien LH. Skeletal muscle triglyceride levels are inversely related to insulin action. *Diabetes*. 1997; 46: 983-988.
- Paolisso G., Gambardella A., Tagliamonte MR., Saccomanno F., Salvatore T., Gualdiero P., D'Onofrio MV., Howard BV. Does free fatty acid infusion impair insulin action also through an increase in oxidative stress? *J Clin Endocrinol. Metab*. 1996b; 81: 4244-4248.

Paolisso G., Gualdiero P., Manzella D., Gambardella A., Verza M., Gentile S., Varricchio M. Hyperinsulinemia is associated with ventricular premature complexes. *Metabolism*. 1996a; 45: 1248-1253.

Paolisso G., Gualdiero P., Manzella D., Rizzo MR., Tagliamonte MR., Gambardella A., Verza M., Gentile S., Varricchio M., D'Onofrio F. Association of fasting plasma free fatty acid concentration and frequency of ventricular premature complexes in nonischemic non-insulin-dependent diabetic patients. *Am J Cardiol*. 1997; 80: 932-937.

Paolisso G., Manzella D., Rizzo MR., Ragno E., Barbieri M., Varricchio G., Varricchio M. Elevated plasma fatty acid concentrations stimulate the cardiac autonomic nervous system in healthy subjects. *Am J Clin Nutr*. 2000; 72: 723-730.

Park JJ., Swan PD. Effect of obesity and regional adiposity on the QTc interval in women. *Int J Obes Relat Metab Disord*. 1997; 21: 1104-1110.

Park SY., Kim YW., Kim JE., Kim JY. Age-associated changes in fat metabolism in the rat and its relation to sympathetic activity. *Life Sci*. 2006; 79: 2228-2233.

Patil TN, Srinivasan M. Hypocholesteremic effect of curcumin in induced hypercholesteremic rats. *Indian J Exp Biol*. 1971; 9(2): 167-9.

Peiris AN., Thakur RK., Sothmann MS., Gustafson AB., Hennes MI., Wilson CR., Kissebah AH. Relationship of regional fat distribution and obesity to electrocardiographic parameters in healthy premenopausal women. *South Med J.* 1991; 84: 961-965.

Pelkonen R., Miettinen TA., Taskinen MR., Nikkila EA. Effect of acute elevation of plasma glycerol, triglyceride and FFA levels on glucose utilization and plasma insulin. *Diabetes.* 1968; 17: 76-82.

Peterson HR., Rothschild M., Weinberg CR., Fell RD., McLeish KR., Pfeifer MA. Body fat and the activity of the autonomic nervous system. *N Engl J Med.* 1988; 318: 1077-1083.

Phan TT., See P., Lee ST., Chan SY. Protective effects of curcumin against oxidative damage on skin cells in vitro: its implication for wound healing. *J Trauma.* 2001; 51: 927-31.

Poehlman ET., Gardner AW., Goran MI., Arciero PJ., Toth MJ., Ades PA., Calles-Escandon J. Sympathetic nervous system activity, body fatness, and body fat distribution in younger and older males. *J Appl Physiol.* 1995; 78: 802-806.

Pomeranz B., Macaulay RJ., Caudill MA., Kutz I., Adam D., Gordon D., Kilborn KM., Barger AC., Shannon DC., Cohen RJ. Assessment of autonomic function in humans by heart rate spectral analysis. *Am J Physiol.* 1985; 248: H151-H153.

Quiles JL., Mesa MD., Ramirez-Tortosa CL., Aguilera CM., Battino M., Gil A., Ramirez-Tortosa M.C. Curcuma longa extract supplementation reduces oxidative stress and attenuates aortic fatty streak development in rabbits. *Arterioscler Thromb Vasc Biol.* 2002; 22(7): 1225-31.

Quintana M., Storck N., Lindblad LE., Lindvall K., Ericson M. Heart rate variability as a means of assessing prognosis after acute myocardial infarction. A 3-year follow-up study. *Eur Heart J.* 1997; 18: 789-797.

"Race, education and prevalence of hypertension". *Am. J. Epidemiol.* 1977; 106: 351-361.

Rao DS, Sekhara NC, Satyanarayana MN, Srinivasan M. Effect of curcumin on serum and liver cholesterol levels in the rat. *J Nutr.* 1970; 100(11): 1307-15.

Reddy AC., Lokesh BR. Effect of dietary turmeric (*Curcuma longa*) on iron-induced lipid peroxidation in the rat liver. *Food Chem Toxicol.* 1994; 32(3): 279-83.

Reddy AC., Lokesh BR. Studies on spice principles as antioxidants in the inhibition of lipid peroxidation of rat liver microsomes. *Mol Cell Biochem.* 1992; 111(1-2): 117-24.

Rigalleau V., Binnert C., Minehira K., Stefanoni N., Schneiter P., Henchoz E., Matzinger O., Cayeux C., Jequier E., Tappy L. In normal men, free fatty acids reduce peripheral but not splanchnic glucose uptake. *Diabetes.* 2001; 50: 727-732.

Robinson M.K. and Thomas A. Obesity and cardiovascular disease. New York : Taylor and Francis group, 2006.

Rocchini AP., Mao HZ., Babu K., Marker P., Rocchini AJ. Clonidine prevents insulin resistance and hypertension in obese dogs. Hypertension. 1999; 33: 548-553.

Roden M., Ludwig C., Nowotny P., Schneider B., Clodi M., Vierhapper H., Roden A., Waldhausl W. Relative hypoleptinemia in patients with type 1 and type 2 diabetes mellitus. Int J Obes Relat Metab Disord. 2000; 24: 976-981.

Roden M., Price TB., Perseghin G., Petersen KF., Rothman DL., Cline GW., Shulman GI. Mechanism of free fatty acid-induced insulin resistance in humans. J Clin Invest. 1996; 97: 2859-2865.

Roden M., Stingl H., Chandramouli V., Schumann WC., Hofer A., Landau BR., Nowotny P., Waldhausl W., Shulman GI. Effects of free fatty acid elevation on postabsorptive endogenous glucose production and gluconeogenesis in humans. Diabetes. 2000; 49: 701-707.

Rukkumani R., Aruna K., Varma PS., Rajasekaran KN., Menon VP. Comparative effects of curcumin and its analog on alcohol- and polyunsaturated fatty acid-induced alterations in circulatory lipid profiles. J Med Food. 2005; 8: 256-260.

Rukkumani R., Sri BM., Vishwanathan P., Menon VP. Comparative effects of curcumin and photo-irradiated curcumin on alcohol- and polyunsaturated fatty acid-induced hyperlipidemia. Pharmacol Res. 2002; 46: 257-264.

Ryan C., Hollenberg M., Harvey DB., Gwynn R. Impaired parasympathetic responses in patients after myocardial infarction. *Am J Cardiol.* 1976; 37: 1013-1018.

Ryysy L., Hakkinen AM., Goto T., Vehkavaara S., Westerbacka J., Halavaara J., Yki-Jarvinen H. Hepatic fat content and insulin action on free fatty acids and glucose metabolism rather than insulin absorption are associated with insulin requirements during insulin therapy in type 2 diabetic patients. *Diabetes.* 2000; 49: 749-758.

Saloranta C., Koivisto V., Widen E., Falholt K., DeFronzo RA., Harkonen M., Groop L. Contribution of muscle and liver to glucose-fatty acid cycle in humans. *Am J Physiol.* 1993; 264: E599-E605.

Samuel VT., Liu ZX., Qu X., Elder BD., Bilz S., Befroy D., Romanelli AJ., Shulman GI. Mechanism of hepatic insulin resistance in non-alcoholic fatty liver disease. *J Biol Chem.* 2004; 279: 32345-32353.

Scherrer U., Randin D., Tappy L., Vollenweider P., Jequier E., Nicod P. Body fat and sympathetic nerve activity in healthy subjects. *Circulation.* 1994; 89: 2634-2640.

Schmitz-Peiffer C., Browne CL., Oakes ND., Watkinson A., Chisholm DJ., Kraegen EW., Biden TJ. Alterations in the expression and cellular localization of protein kinase C isozymes epsilon and theta are associated with insulin resistance in skeletal muscle of the high-fat-fed rat. *Diabetes.* 1997; 46: 169-178.

Schoonjans K., Staels B., Auwerx J. Role of the peroxisome proliferator-activated receptor (PPAR) in mediating the effects of fibrates and fatty acids on gene expression. *J Lipid Res.* 1996; 37: 907-925.

Sgoifo A., Koolhaas JM., Musso E., De Boer SF. Different sympathovagal modulation of heart rate during social and nonsocial stress episodes in wild-type rats. *Physiol Behav.* 1999; 67: 733-738.

Shaltout HA., bdel-Rahman AA. Mechanism of fatty acids induced suppression of cardiovascular reflexes in rats. *J Pharmacol Exp Ther.* 2005; 314: 1328-1337.

Shao ZM., Shen ZZ., Liu CH., Sartippour MR., Go VL., Heber D., Nguyen M. Curcumin exerts multiple suppressive effects on human breast carcinoma cells. *Int J Cancer.* 2002; 98: 234-40.

Shimoguchi K. [Analysis of immunoprotection against fatal experimental *Klebsiella pneumoniae pneumonia*]. *Kansenshogaku Zasshi.* 1990; 64: 1482-1492.

Shishodia S., Sethi G., Aggarwal BB. Curcumin: getting back to the roots. *Ann N Y Acad Sci.* 2005; 1056: 206-217.

Shulman GI. Cellular mechanisms of insulin resistance. *J Clin. Invest.* 2000; 106: 171-176.

Sowers JR., Whitfield LA., Catania RA., Stern N., Tuck ML., Dornfeld L., Maxwell M. Role of the sympathetic nervous system in blood pressure maintenance in obesity. *J Clin Endocrinol Metab.* 1982; 54: 1181-1186.

Sreejayan N., Rao MN. Curcuminoids as potent inhibitors of lipid peroxidation. *J Pharm Pharmacol.* 1994; 46(12): 1013-6.

Srimal RC., Dhawan BN. Pharmacology of diferuloyl methane(curcumin), a non-steroidal anti-inflammatory agent. *J Pharm Pharmacol.* 1973; 25(6): 447-52.

Srinivasan K., Viswanad B., Asrat L., Kaul CL., Ramarao P. Combination of high-fat diet-fed and low-dose streptozotocin-treated rat: a model for type 2 diabetes and pharmacological screening. *Pharmacol Res.* 2005; 52: 313-320.

Srihari RT., Basu N., Siddiqui HH. Anti-inflammatory activity of curcumin analogues. *Indian J Med Res.* 1982; 75: 574-8.

Srivastava R., Srimal RC. Modification of certain inflammation-induced biochemical changes by curcumin. *Indian J Med Res.* 1985; 81: 215-23.

Staels B., Fruchart JC. Therapeutic roles of peroxisome proliferator-activated receptor agonists. *Diabetes.* 2005; 54: 2460-2470.

Stamler R., Stamler J., Riedlinger WF., Algera G., Roberts RH. Weight and blood pressure: Findings in hypertension screening of 1 million Americans. *JAMA* 1978; 240: 1607-1610.

Stein PK., Bosner MS., Kleiger RE., Conger BM. Heart rate variability: a measure of cardiac autonomic tone. *Am Heart J.* 1994; 127: 1376-1381.

Stein PK., Kleiger RE. Insights from the study of heart rate variability. *Annu Rev Med.* 1999; 50: 249-261.

Storlien LH., Jenkins AB., Chisholm DJ., Pascoe WS., Khouri S., Kraegen EW.

Influence of dietary fat composition on development of insulin resistance in rats.

Relationship to muscle triglyceride and omega-3 fatty acids in muscle phospholipid. *Diabetes*. 1991; 40: 280-289.

Sturm R., Wells KB. Does obesity contribute as much to morbidity as poverty or smoking?. *Public Health*. 2001; 115: 229-235.

Sztajzel J. Heart rate variability: a noninvasive electrocardiographic method to measure the autonomic nervous system. *Swiss Med Wkly*. 2004; 134: 514-522.

Task Force of the European Society of Cardiology and the North American Society of Pacing and Electrophysiology. "Heart rate variability: standards of measurement, physiological interpretation and clinical use". *Circulation*. 1996a; 93: 1043-1065.

Tentolouris N., Tsigos C., Perea D., Koukou E., Kyriaki D., Kitsou E., Daskas S., Daifotis Z., Makrilakis K., Raptis SA., Katsilambros N. Differential effects of high-fat and high-carbohydrate isoenergetic meals on cardiac autonomic nervous system activity in lean and obese women. *Metabolism*. 2003; 52: 1426-1432.

Terathongkum S., Pickler RH. Relationships among heart rate variability, hypertension, and relaxation techniques. *J Vasc Nurs*. 2004; 22: 78-82.

The National Heart, Lung and Blood Institute. Clinical Guidelines on the Identification, Evaluation, and Treatment of Overweight and Obesity in Adults-- The Evidence Report. *Obes Res.* 1998; 6 Suppl 2: 51S-209S.

Tiikkainen M., Tamminen M., Hakkinen AM., Bergholm R., Vehkavaara S., Halavaara J., Teramo K., Rissanen A., Yki-Jarvinen H. Liver-fat accumulation and insulin resistance in obese women with previous gestational diabetes. *Obes Res.* 2002; 10: 859-867.

Tsuji H., Venditti FJ., Jr., Manders ES., Evans JC., Larson MG., Feldman CL., Levy D. Reduced heart rate variability and mortality risk in an elderly cohort. The Framingham Heart Study. *Circulation.* 1994; 90: 878-883.

Tsuji H., Venditti FJ., Jr., Manders ES., Evans JC., Larson MG., Feldman CL., Levy D. Determinants of heart rate variability. *J Am Coll Cardiol.* 1996; 28: 1539-1546.

The National Institutes of Health, NHLBI document. "Clinical Guidelines on the Identification, Evaluation, and Treatment of Overweight and Obesity in Adults: The Evidence Report". Dept. of Health and Human Services, U.S. 2000: 98-408.

Unnikrishnan MK., Rao M.N. Curcumin inhibits nitrogen dioxide induced oxidation of hemoglobin. *Mol Cell Biochem.* 1995; 146(1): 35-7.

Valensi P., Thi BN., Lormeau B., Paries J., Attali JR. Cardiac autonomic function in obese patients. *Int J Obes Relat Metab Disord.* 1995; 19: 113-118.

Van Ravenswaaij-Arts CM., Kollee LA., Hopman JC., Stoelinga GB., van Geijn HP.

Heart rate variability. *Ann Intern Med.* 1993; 118: 436-447.

Verwaerde P., Senard JM., Galinier M., Rouge P., Massabuau P., Galitzky J., Berlan M., Lafontan M., Montastruc JL. Changes in short-term variability of blood pressure and heart rate during the development of obesity-associated hypertension in high-fat fed dogs. *J Hypertens.* 1999; 17: 1135-1143.

Villareal RP., Liu BC., Massumi A. Heart rate variability and cardiovascular mortality. *Curr Atheroscler.* 2002; Rep. 4: 120-127.

Vlay SC., Mallis GI, Brown EJ., Jr., Cohn PF. Documented sudden cardiac death in prolonged QT syndrome. *Arch Intern Med.* 1984; 144: 833-835.

Wan R., Camandola S., Mattson MP. Intermittent fasting and dietary supplementation with 2-deoxy-D-glucose improve functional and metabolic cardiovascular risk factors in rats. *FASEB J.* 2003; 17: 1133-1134.

Wan R., Camandola S., Mattson MP. Intermittent food deprivation improves cardiovascular and neuroendocrine responses to stress in rats. *J Nutr.* 2003; 133: 1921-1929.

Williamson JR., Kreisberg RA., Felts PW. Mechanism for the stimulation of gluconeogenesis by fatty acids in perfused rat liver. *Proc Natl Acad Sci USA.* 1966; 56: 247-254.

Wyne KL. Free fatty acids and type 2 diabetes mellitus. *Am J Med.* 2003; 115 Suppl 8A: 29S-36S.

- Xu J., Fu Y., Chen A. Activation of peroxisome proliferator-activated receptor- γ contributes to the inhibitory effects of curcumin on rat hepatic stellate cell growth. *Am. J. Physiol. Gastrointest. Liver Physiol.* 2003; 285: G20-G30.
- Yao QH., Wang DQ., Cui CC., Yan ZY., Chen IB., Yao XW., Wang JK., Lian JF. Curcumin ameliorates Left ventricular function in rabbits with pressure overload inhibition of the remodeling of the Left ventricular collagen network associated with suppression of myocardial Tumor Necrosis Factor- α and Matrix Metalloproteinase – 2 expression. *Biol Pharm Bull.* 2004; 27(2): 198-202.
- Yegnanarayan R., Saraf AP., Balwani JH. Comparison of anti-inflammatory activity of various extracts of *Curcuma longa* (Linn). *Indian J Med Res.* 1976; 64(4): 601-8.
- Young JB., Landsberg L. Stimulation of the sympathetic nervous system during sucrose feeding. *Nature.* 1977; 269: 615-617.
- Young JB., Saville E., Rothwell NJ., Stock MJ., Landsberg L. Effect of diet and cold exposure on norepinephrine turnover in brown adipose tissue of the rat. *J Clin Invest.* 1982; 69: 1061-1071.
- Zahorska-Markiewicz B., Kuagowska E., Kucio C., Klin M. Heart rate variability in obesity. *Int J Obes Relat Metab Disord.* 1993; 17: 21-23.
- Zipes DP. Sudden cardiac death. Future approaches. *Circulation.* 1992; 85: 1160-1166.
- Zipes DP., Wellens HJ. Sudden cardiac death. *Circulation.* 1998; 98: 2334-2351.