

## REFERENCES

1. สำนักงานคณะกรรมการป้องกันและปราบปรามยาเสพติด. “สถิติการจับกุมยาเสพติดทั่วประเทศ แสดงจำนวนคดีจำแนกตามชนิดยาเสพติด ปี ๒๕๔๓-๒๕๔๗” [ระบบออนไลน์] แหล่งที่มา: <http://www.oncb.go.th/c2-statistic.html> (21 ตุลาคม 2548).
2. สำนักงานคณะกรรมการป้องกันและปราบปรามยาเสพติด. “สรุปผลการประมาณการผู้เกี่ยวข้องกับยาเสพติดในประเทศไทย” [ระบบออนไลน์]. แหล่งที่มา: <http://www.oncb.go.th/document/situation5.html> (10 ตุลาคม 2548).
3. Adam P. “Drug characterization of methamphetamine pills as a scientific tool to help identify drug production and trafficking network”. Ph.D. Thesis, Chiang Mai University, 2005.
4. Kintz P. “Amphetamine determination in hair.” in Analytical and practical aspects drug testing in hair, Taylor & Francis group, CRC Press Boca Raton., New York, 2006:143-144.
5. Yudko E, Hall VH, McPherson SB. Methamphetamine use: clinical and forensic aspects, Boca Raton, FL: CRC Press, 2003.
6. Moffat CA, Osselton MD, Widdop B. “Monographs.” in Clarke’s Analysis of Drugs and Poison. 3rd ed. Chicago: Pharmaceutical Press, 2004;612-4: 1226-8.
7. Ellison J, Dobies D. Methamphetamine abuse presenting as dysuria. *J. Annals of Emergency. Medicine.* 1984; 13: 198-200.

8. Cook CE, Jeffcoat AR, Sadler BM, et al. Pharmacokinetics of oral methamphetamine and effects of repeated daily dosing in humans. *J. Drug Metab. Dispos.* 1992; 20: 856–862.
9. McGregor C, Srisurapanont M, Jittiwutikarn J, Laobhripatr S, Wongtan T, White J. "The nature, time course and severity of methamphetamine withdrawal". *J. Addiction* . 2005; 100 (9): 1320-9.
10. Karch SB. "Pharmacokinetics of specific drugs" in Drug Abuse Handbook. Washington D.C., CRC Press, 1998: 175.
11. Brust JCM. Neurological aspects of substance abuse. Butterworth-Heinemann, Stoneham, MA. 1993.
12. Sribanditmongkol P, Chokjamsai M, Tampitak S. Methamphetamine overdose and fatality: 2 cases report. *J. Med Assoc Thai*: 2000; 83(9): 1120-3.
13. Bennett B, Hollingsworth C, Martin RH. Methamphetamine-induced alterations in dopamine transporter function. *J. Brain Res* . 1998; 82 (1-2): 219-27.
14. Wu AHB. "Mechanism of Interferences for Gas Chromatography/Mass Spectrometry Analysis of Urine for Drugs of Abuse," *J. Annals of Clinical Laboratory Science* 1995; 25: 319–329.
15. Moeller MR, Steinmeyer S, Kraemer T. Determination of drugs of abuse in blood. *J. Chromatogr. B.* 1998; 713(1): 91-109.
16. Wenning R. Potential problems with the interpretation of hair analysis results. *J. Forensic Sci Int.* 2000; 107: 5-12.
17. Moeller MR, Fey P, Wening R, Sachs H. Hair analysis as evidence in forensic cases. *J. Forensic Sci Int.* 1993; 63:185-206.

18. Zhoa H, Brenneisen R, Scholer A, et al. *J. Anal. Toxicol.* 2001; 25L: 258.
19. Baumgartner AM, Jones PF, Baumgartner WA, Black CT. Radioimmuno assay of hair for determining opiates abuse histories. *J Nuclear Med.* 1979; 20: 748-752.
20. Kramer P, Maurer HH. Liquid chromatography-mass spectrometry in forensic and clinical toxicology. *J. Chromatogr. B.* 1998; 713 :163.
21. Gentili S, Cornetta M, Macchia T. Rapid screening procedure based on headspace solid-phase microextraction and gas chromatography-mass spectrometry for the detection of many recreational drugs in hair. *J. Chromatogr. B.* 2004; 801: 289-96.
22. Saitoh M, Uzuka M, Sakamoto M. "Rate of hair growth", in Montagna W, Dobson RL (eds): *Advances in Biology of the Skin, Vol IX, Hair Growth.* Oxford, Pergamon. 1969: 83.
23. "Hair physiology". Hair restoration, The Blackrock clinic, Blackrock, co. Dublin, Ireland. [Online].  
Available : <http://hairtransplantsurgery.ie> (2008, February 20)
24. Kintz P. "Hair analysis." in Clarke's *Analysis of Drugs and Poison.* 3rd ed. Chicago: Pharmaceutical Press, 2004: 124-133.
25. Society of Hair Testing. Recommendation for hair testing in forensic cases, *J. Forensic Sci Int.* 2004; 145: 83-4.
26. Garside DGB. *Determination of Cocaine and Opioids in Hair,* CRC Press, Boca Raton, FL, 1996.
27. Sribanditmongkol P, Thampitak S, Sathawornwiwat P, Jittiwutikarn J. Hair analysis for morphine in heroin addicts in Thailand. *Thai J Health Research.* 2001; 15:13-27.

28. Lucas AC, Barnejo AM, Taberero MJ, Fernandes P, Strano-Rossi S. Use of solid-phase microextraction (SPME) for determination of methadone EDDP in human hair by GC-MS, *J. Forensic Sci Int.* 2000; 107: 225-232.
29. Peterson S, MaLachlan-Troup N, Cordero R, Dohnal M, Carman S. Quantitative screening for drugs of abuse in hair using GC-MS. *J. Anal. Toxicol.* 2001; 25: 203-208.
30. Goldberger BA, Caplan YH, Maguire T, Cone EJ. Testing human hair for drugs of abuse, III: identification of heroin and 6-acetylmorphine as indicators of heroin use. *J. Anal. Toxicol.* 1991; 15: 226-231.
31. Goldberger BA, Darraj AG, Caplan YH, Cone EJ. Detection of methadone, methadone metabolites, and other illicit drugs of abuse in hair of methadone treatment subjects. *J. Anal. Toxicol.* 1998; 22: 526-530.
32. Girod C, Staub C. Analysis of drugs of abuse in hair by automated solid-phase extraction, GC/EI/MS and GC ion trap/CI/MS *J. Forensic Sci Int.* 2000; 107: 261-271.
33. Uhl M. Tandem mass spectrometry: a helpful tool in hair analysis for the forensic expert. *J. Forensic Sci Int.* 2000; 107:169-179.
34. Sporkert F, Pragst F. Determination of methadone and its metabolites EDDP and EMDP in human hair by headspace solid-phase microextraction and gas chromatography-mass spectrometry. *J. Forensic Sci Int.* 2000; 107 (1-3):225-232.
35. [Lachenmeier](#) DW, [Kroener](#) L, [Musshoff](#) F, [Madea](#), B. Application of tandem mass spectrometry combined with gas chromatography and headspace solid-

phase dynamic extraction for the determination of drugs of abuse in hair samples by Rapid Commun Mass Spectrom. 2003; 17: 472-478.

36. Berens G, Yegles M, Wenning R. Study of methadone and EPPD enantiomer ratio in hair, urine and serum by capillary electrophoresis, Proceedings TIAFT, Paris, 2003: 26-30.
37. Balíkova MA, Habrdova V. Hair analysis for opiates: evaluation of washing and incubation procedures. *J. Chromatography. B.* 2003; 789(11): 93-100.
38. Nakahara Y, Kikura R, Takahashi K. Hair analysis for drugs of abuse VIII. Effective extraction and determination of 6-acetylmorphine and morphine in hair with trifluoroacetic acid and methanol for the confirmation of retrospective heroin use by GCIMS. *J Chromatogr B:Biomedical Appl.* 1994; 657: 93-101.
39. Gentili S, Torresi A, Marsili R, Chirotti M, Macchia T. Simultaneous detection of amphetamine-like drug with headspace solid-phase microextraction and gas chromatography–mass spectrometry. *J. Chromatogr. B.* 2002; 780: 183-192.
40. Arthur CL, Pawliszyn J. Solid Phase Microextraction with Thermal Desorption Using Fused Silica Optical Fibers. *Anal. Chem.* 1990; 62: 2145.
41. Arthur CL, Killam LM, Motlagh S, Lim M, Potter DW, Pawliszyn J. Analysis of substituted benzene compounds in groundwater using solid-phase microextraction. *J. Environ Sci. Technol.* 1992; 26(5): 979-983.
42. Almeida CMM, Boas LV. Analysis of BTEX and other substituted benzenes in water using headspace SPME-GC-FID: method validation. *J. Environ. Monit.*, 2004; 6: 80 – 88.

43. Randolph CG, Alfredo JC, Michael DW, Stephen LM. Analysis of Volatile Fragrance and Flavor Compounds by Headspace Solid Phase Microextraction Combined with Gas Chromatography/Mass Spectrometry. An instrumental analysis experiment; Department of Chemistry and Biochemistry, The University of South Carolina, Columbia, SC. [online]  
Available: <http://www.chem.sc.edu/.../lab/images/RGFig1.JPG> (2007, May 10)
44. Wong RC, Tse HY. "Drug-testing technologies and applications", in Drug of abuse: Body fluid testing. Humana press 2005: 29-63.
45. "Chromatography". Analytical techniques. [online]  
Available: <http://www.fao.org> (2008, January 10)
46. "Schematic view of the ion source based on electron impact ionization and the quadrupole mass filter typically found in a GC-MS instrument". Microbial cell factories. [online].  
Available: <http://www.microbialcellfactories.com/> (2008, January 14)
47. Agilent technologies. "An introduction to mass spectrometry for GC-the single quadrupole" in Mass spectrometry solution seminar on 22-26 August 2005. p. 2.
48. Guidance for Industry. Analytical Procedures and Methods Validation Chemistry, Manufacturing, and Controls Documentation Food and Drug Administration (FDA), 2005. [online]  
Available: <http://www.fda.gov/cder/guidance/2396dft.htm> (2006, January 10)
49. Junting L, Peng C. Solid-phase microextraction (SPME) of drugs and poisons from biological samples. *J. Forensic Sci Int.* 1998; 97:93-100.

50. Chia KJ, Huang SD. Simultaneous derivatization and extraction of amphetamine-like drugs in urine with headspace solid-phase microextraction followed by gas chromatography-mass spectrometry. *J. Anal. Chim. Acta.* 2005; 539: 49-54.
51. Nantarat S. "Determination of amphetamine, methamphetamine and ephedrine in urine by gas chromatography-mass spectrometry". M.S. Thesis, Chiang Mai University, 1998.
52. Bournissen FG, Rokach B, Karaskov T, Koren G. Methamphetamine detection in maternal and neonatal hair: implications for fetal safety. *J. Archives of Disease in Childhood - Fetal and Neonatal Edition.* 2007; 92:351-355.
53. Nishida M, Yashiki M, Namera A, Kimura K. Single hair analysis of methamphetamine and amphetamine by solid phase microextraction coupled with in matrix derivatization. *J. Chromatogr. B.* 2006; 842: 106-110.
54. Lord HL, Pawliszyn J. Method optimization for analysis of amphetamines in urine by solid phase microextraction. *Anal. Chem.* 1997; 69: 3899-3906.
55. Nik JN, Bska AJ, Zygmunt B. Determination of volatile aliphatic amines in air by solid-phase microextraction coupled with gas chromatography with flame ionization detection *J. Chromatogr. A.* 2003; 1016(1): 1-9.
56. Musshoff F, Lachenmeier DW, Kroener L, Madea K. Automated headspace solid-phase dynamic extraction for the determination of amphetamines and synthetic designer drugs in hair samples. *J. Chromatogr. A.* 2002; 958: 231-238.

57. Raikos N, Christopoulou K, Theodoridis G, Tsoukali H, Psaroulis D. Determination of amphetamines in human urine by headspace solid-phase microextraction and gas chromatography. *J. Chromatogr. B* 2003; 789: 59-63.
58. Okajima K, Narera A, Yashiki M, Tsukue I, Kojima T. Highly sensitive methamphetamine and amphetamine in human whole blood using headspace solid-phase microextraction and gas chromatography-mass spectrometry. *J. Forensic Sci Int.* 2001; 116:15-22.
59. Pragst F, Balikova M.A. State of art in hair analysis for detection of drug and alcohol abuse. *J. Clin. Chim. Acta.* 2006; 370: 1749.
60. Miki A, Katagi M, Zaitso K, Nishioka H, Tsuchihashi H. Development of a two-step injector for GC-MS with on-column derivatization, and its application to the determination of amphetamine-type stimulants (ATS) in biological specimens. *J. Chromatogr. B.* 2002; 865: 25-32.
61. Xiang P, Shen M, Zhuo X. Hair analysis for ketamine and its metabolites. *J. Forensic Sci Int.* 2006; 162: 131-134.