TABLE OF CONTENT



| 1.2.10 Oncogene-induced senescence | 52 |
|---|-----------------|
| 1.2.11 Senescence and tumorigenesis | 57 |
| 1.2.12 CD44 structure and interactions | 58 |
| 1.2.13 CD44 functions and relating signals | 62 |
| 1.3 Objective | 68 |
| CHAPTER II MATERIALS AND METHODS | |
| 2.1 Materials | 69 |
| 2.2 Methods | |
| 2.2.1 Preparation methods | |
| 2.2.1.1 Versican Isolation | 69 |
| 2.2.1.2 WT and A subdomain less versican mouse | 70 |
| embryonic fibroblasts culture preparation | |
| 2.2.2 Analytical methods | |
| 2.2.2.1 Cell proliferation assay | 72 |
| 2.2.2.2Colony formation in soft agarose gel | 73 |
| 2.2.2.3Tumor growth <i>in vivo</i> | 73 |
| 2.2.2.4 β-galactosidase staining | 73 |
| 2.2.2.5Immunocytochemistry and immunohistochemistry | S ⁷⁴ |
| 2.2.2.6 Sodium dodecyl sulfate-polyacrylamide gel | 77 |
| Electrophoresis (SDS-PAGE) and immunoblot analysis | |
| 2.2.2.7 Chondroitin sulfate chain analysis | 79 |
| 2.2.2.8 Real-time reverse transcription-polymerase | 81 |
| chain reaction (real-time RT-PCR) | |

| 2.2.2.9Sandwich ELISA for hyaluronan content determination | 82 |
|--|------------------|
| 2.2.2.10 Cell adhesion assay | 83 |
| 2.2.2.11 Attachment assay | 83 |
| 2.2.2.12 Protein assay | 84 |
| 2.2.2.13 Collagen assay | 84 |
| 2.2.2.14 Treatment with an anti-CD44 antibody | 85 |
| and hyaluronidase | |
| 2.2.2.15 Treatment with a hyaluronan fragments | 86 |
| CHAPTER III RESULTS | |
| 3.1 Cell behavior and morphology | |
| 3.1.1 Fibroblasts culture established from $Cspg2^{\Delta 3/\Delta 3}$ embryo | 87 |
| 3.1.2 $Cspg2^{\Delta 3/\Delta 3}$ embryo fibroblasts expressed low proliferation | 94 |
| rate and premature senescence characteristic at early passages | |
| 3.1.3 $Cspg2^{\Delta 3/\Delta 3}$ fibroblasts escaped from senescence, and | 99 |
| expressed high proliferation rate after growing more than ten pas | sages |
| 3.1.4 The transformed $Cspg2^{\Delta 3/\Delta 3}$ fibroblasts were malignant cells | 104 |
| 3.1.5 Adhesion ability of the transformed $Cspg2^{\Delta 3/\Delta 3}$ fibroblasts | 111 |
| 3.1.6Attaching ability of the transformed $Cspg2^{\Delta 3/\Delta 3}$ fibroblasts | S ₁₁₃ |
| 3.1.7 Exogenous versican adding to the $Cspg2^{\Delta 3/\Delta 3}$ fibroblasts | 115 |
| did not recover cell proliferation rate to the WT fibroblasts level | |
| 3.2 Extracellular matrix proteins expression | |
| 3.2.1 Versican expression decreased in the $Cspg2^{\Delta 3/\Delta 3}$ fibroblasts | 118 |

х

the $Cspg2^{\Delta 3/\Delta 3}$ fibroblasts

3.2.3 Chondroitin sulfate content in versican from the 122 $Cspg2^{A3/A3}$ fibroblasts culture was decreased in the ECM

but increased in cultured medium

3.2.4 HA deposition is altered in levels and structure in the 126 $Cspg2^{\Delta 3/\Delta 3}$ fibroblasts

3.2.5 Expression of fibronectin, laminin in the $Cspg2^{\Delta 3/\Delta 3}$ 129

and the transformed $Cspg2^{\Delta 3/\Delta 3}$ cell cultures are comparable to the WT

3.2.6 Collagen expression decreased in the $Cspg2^{\Delta 3/\Delta 3}$, but 131

recovered in the transformed $Cspg2^{\Delta 3/\Delta 3}$ cells culture

3.3 Signaling pathway elucidation

- 3.3.1p53 protein was upregulated in the Cspg2^{A3/A3} fibroblasts, 134 but abrogated in the transformed Cspg2^{A3/A3} fibroblasts
 3.3.2Phosphorylated ERK1/2 was upregulated in the 135 Cspg2^{A3/A3} fibroblasts, but downregulated in the transformed Cspg2^{A3/A3} fibroblasts
 3.3.3Phosphorylated ERK1/2 was upregulated by upstream 137 molecules of the MAPK signal transduction cascade
 - 3.3.4 Anti-CD44 suppressed ERK1/2 phosphorylation in the 138 $Cspg2^{\Delta3/\Delta3}$ fibroblasts culture

| 3.3.5HA-CD44 interaction mediated ERK1/2 phosphorylation | on 141 |
|--|--------|
| in the hyaluronidase treated WT fibroblasts culture | |
| 3.3.6 β 1-integrin and EGF receptor did not involve ERK1/2 | 143 |
| phosphorylation in the $Cspg2^{\Delta 3/\Delta 3}$ fibroblasts culture | |
| 3.3.7 Exogenous HA treatment enhanced phosphorylation of | f 144 |
| ERK1/2 | |
| CHAPTER IV DISCUSSION | 146 |
| REFERENCES | 5 154 |
| APPENDIX | 177 |
| CURRICULUM VITAE | 187 |
| | |
| | |
| ALUNTERST | |

ลิขสิทธิ์มหาวิทยาลัยเชียงใหม่ Copyright[©] by Chiang Mai University All rights reserved

xii

LIST OF TABLES

| Table | | | |
|-------|--|-----|--|
| 1 | Concentration of hyaluronan in tissues and tissue fluids | 41 | |
| 2 | Chondroitin sulfate derived disaccharide composition in proteoglycan | 124 | |
| | extracted from cell layers and cultured media of fibroblasts. | | |
| 3 - 5 | Chondroitin sulfate disaccharide composition of versican | 187 | |
| | and aggrecan. | | |

ลิขสิทธิ์มหาวิทยาลัยเชียงใหม่ Copyright[©] by Chiang Mai University All rights reserved

| LIST OF FIGURES | | |
|-----------------|---|------|
| Figu | re | Page |
| 1 | Structure of extracellular matrix composes of several extracellular molecules. | 7 |
| 2 | The modular organization of Hyaluronan Binding Proteins. | 10 |
| 3 | Schematic representation of the human versican gene and | 16 |
| | intron/exon organization. | |
| 4 | A schematic of versican structure, composing of core protein | 17 |
| | (G1, CS chain binding region, and G3 domain). | |
| 5 | The interaction of versican with other molecules. | 26 |
| 6 | Schematic diagram of interaction of the G1 domain of versican | 27 |
| | (Ver), link protein (LP), and hyaluronan (HA). | |
| 7 | The chemical structure of hyaluronan. | 39 |
| 80 | Schematic of CD44-HA interactions. | 44 |
| 9 V | Multiple pathways mediate Oncogene-induced senescence. | 56 |
| 10 | Schematic diagram of the structure of the CD44 gene. | 66 |
| 11 | CD44 protein structure. | 67 |
| 12 | The genomic organization of $Cspg2^{\Delta 3/\Delta 3}$, the targeting vector, | 88 |
| | and the mutated allele by homologous recombinant. | |
| 13 | Phenotype of Cspg2 gene modified mouse and embryo. | 89 |

| 14 | Agarose Gel electrophoresis of PCR products from mice embryo tails. | 91 |
|----|---|-----|
| 15 | Morphology of the WT and $Cspg2^{\Delta 3/\Delta 3}$ fibroblasts at passages 5. | 93 |
| 16 | Proliferation rates of MEFs at different phenotype, | 95 |
| | measured by BrdU incorporation ELISA assay. | |
| 17 | Proliferation rates of MEFs at different phenotype, | 96 |
| | measured cell counting assay. | |
| 18 | Staining of β -galactosidase activity in fibroblasts culture. | 98 |
| 19 | Proliferation rates of MEFs in each passage. | 102 |
| 20 | Morphology of MEFs in each passage. | 103 |
| 21 | Agarose Gel electrophoresis of PCR products from | 104 |
| | fibroblasts culture. | |
| 22 | Colony formation assay. Fibroblasts were grown on 7% | 106 |
| | soft agarose gel culture. | |
| 23 | The tumor formation in BALB/c nu/nu mice. | 108 |
| 24 | Hematoxylin & Eosin histological staining of tumor isolated | 110 |
| | from mice which injected with the $Cspg2^{\Delta 3/\Delta 3}$ fibroblasts | |
| 25 | Adhering ability of the $Cspg2^{\Delta 3/\Delta 3}$ fibroblasts to various | 112 |
| | extracellular matrix molecules. | |
| 26 | Attaching ability of fibroblast. | 114 |
| 27 | Dot blot analysis of versican and aggrecan which extracted | 116 |
| | from mice brain. | |
| 28 | Proliferation rate of WT, $Cspg2^{\Delta 3/\Delta 3}$, and transformed | 117 |

 $Cspg2^{\Delta 3/\Delta 3}$ fibroblasts.

| 29 | Expression level of versican mRNA in wild type, $Cspg2^{\Delta 3/\Delta 3}$ (mortal) | 119 |
|----|--|-----|
| | and transformed $Cspg2^{\Delta 3/\Delta 3}$ (IM1, IM2, IM3) fibroblasts. | |
| 30 | Localization of versican molecule in ECM of the WT, the $Cspg2^{\Delta 3/\Delta 3}$, | 121 |
| | and the transformed $Cspg2^{\Delta 3/\Delta 3}$ fibroblasts. | |
| 31 | Chondroitin sulfate derived disaccharide content of extracted | 125 |
| | proteoglycan from cell layers cultured media and total content | |
| | from cell layers and cultured media. | |
| 32 | Localization of hyaluronan molecule in ECM of the WT, | 127 |
| | the $Cspg2^{\Delta 3/\Delta 3}$, and the transformed $Cspg2^{\Delta 3/\Delta 3}$ fibroblasts. | |
| 33 | Content of hyaluronan in cell layers, and cultured media | 128 |
| | of fibroblasts cultures. | |
| 34 | Localization of fibronectin (FN) and laminin (LN) in ECM of | 130 |
| | wild type (WT), the $Cspg2^{\Delta 3/\Delta 3}$ (mortal), and the transformed | |
| | $Cspg2^{\Delta 3/\Delta 3}$ (IM) fibroblasts. | |
| 35 | Expression of fibronectin (FN) and laminin (LN) from | 131 |
| | fibroblast cultures. | |
| 36 | Localization of type I collagen in ECM of the WT, | 132 |
| | the $Cspg2^{\Delta 3/\Delta 3}$, and the transformed $Cspg2^{\Delta 3/\Delta 3}$ fibroblasts. | |
| 37 | Content of collagen in cell layers, and cultured media | 133 |
| | of fibroblasts cultures. | |
| | | |

38 Expression of p53 protein in the WT, the $Cspg2^{\Delta 3/\Delta 3}$, 135

and the transformed $Cspg2^{\Delta 3/\Delta 3}$.

| 39 | Expression of MAPK signaling protein in (ERK1/2, p38, JNK) | 136 |
|----|---|-----|
| | the WT, the $Cspg2^{\Delta 3/\Delta 3}$, and the transformed $Cspg2^{\Delta 3/\Delta 3}$. | |
| 40 | Effects of PD98059 and EGF to phosphorylation level of ERK1/2. | 138 |
| 41 | Effects of anti-CD44 treatment on phosphorylated ERK1/2. | 140 |
| 42 | Effects of hyaluronidase treatment on phosphorylation of | 142 |
| | ERK1/2 in the WT fibroblasts. | |
| 43 | Effects of combined treatment with hyaluronidase and anti-CD44. | 142 |
| 44 | Expression of integrin and phosphorylatedEGFR, the WT, | 144 |
| | the $Cspg2^{\Delta 3/\Delta 3}$, and the transformed $Cspg2^{\Delta 3/\Delta 3}$. | |
| 45 | Effects of exogenous HA on phosphorylation of ERK1/2. | 145 |
| 46 | A schematic diagram showing the mechanism underlying | 147 |
| | acquisition of the senescence and the immortality in the | |
| | $= -\sqrt{3}/\sqrt{3}$ at $= 1.1$ | |

Cspg2^{A3/A3} fibroblasts. Copyright[©] by Chiang Mai University All rights reserved

xvii



xviii



Hyaluronan Synthase

xix

HAS



Molar

XX



Phosphatidylinositol 3-kinase

PI3K

xxi



xxii



ลิขสิทธิ์มหาวิทยาลัยเชียงใหม่ Copyright[©] by Chiang Mai University All rights reserved

xxiii