

REFERENCES

- Adav, S.S., Chen, M.Y., Lee, D.J. and Ren, N.Q. 2007. Degradation of phenol by aerobic granules and isolated yeast *Candida tropicalis*. *Biotechnol. Bioeng.* 67: 1566-1572.
- Agency for Toxic Substances and Disease Registry (ATSDR). 1998. Toxicological profile for phenol. Update. Atlanta, GA.
- Ahmed A.M., G.F. Nakhla, and Farooq, S.J. 1995. Phenol degradation by *Pseudomonas aeruginosa*, *J. Environ. Sci. Health.* 30: 99-107.
- Aksu, Z. 2005. Application of biosorption for removal of organic pollutants: a review. *Process Biochem.* 40: 997-1026.
- Alexander, M. 1994. *Biodegradation and bioremediation*. San Diego, California: Academic Press pp. 1-7.
- Amore, J.E. and Hautala, E. 1983. Odor as an aid to chemical safety: odor thresholds compared with threshold limit values and volatilities for 214 industrial chemicals in air and water dilution. *J. Appl. Toxicol.* 3: 272-290.
- Annadurai G., R. S. Juang, and D.J. Lee. 2002. Microbiological degradation of phenol using mixed liquors of *Pseudomonas putida* and activated sludge. *Waste Manage.* 22: 703-710.
- Azuma, M., Ikeuchi, T., Kiritani, R., Kato, J. and Ooshima, H. 2000. Increase in xylitol production by *Candida tropicalis* upon addition of salt. *Biomass. Bioenerg.* 19: 129-135.

- Babel, W., Brinkmann, U. and Mueller, R. H. 1993. The auxiliary substrate concept. An approach for overcoming limits of microbial performances. *Acta Biotechnol.* 13: 211-242.
- Bai, J., Wen, J.P., Li, H.M. and Jiang, Y. 2007. Kinetic modeling of growth and biodegradation of phenol and m-cresol using *Alcaligenes faecalis*. *Process Biochem.* 42: 510-517.
- Bandyopadhyay, K., Das, D. and Maiti, B.R. 1998. Kinetics of phenol degradation using *Pseudomonas putida* MTCC 1194. *Bioprocess Eng.* 18(5): 373-377.
- Bandyopadhyay, K., Das, D., Bhattacharyya, P. and Maiti, B.R. 2001. Reaction engineering studies on biodegradation of phenol by *Pseudomonas putida* MTCC 1194 immobilized on calcium alginate. *J. Biochem. Eng.* 8: 179-186.
- Barker, E.L., Peter, E.B., Petrecia, H.F., and Grant, S.K. 1978. Phenol poisoning due to contaminated drinking water. *Arch. Environ. Health.* 33: 89-94.
- Bastos, A.E., Moon, D.H., Rossi, A., Trevors, J.T. and Tsai, S.M. 2000. Salt-tolerant phenol-degrading microorganisms isolated from Amazonian soil samples. *Arch. Microbiol.* 174: 346-352.
- Bechard, G., Bisailon, J.G., Beaudet, R. and Sylvestre, M. 1990. Degradation of phenol by a bacterial consortium under methanogenic conditions. *Can. Microbiol.* 36: 573-578.
- Bisping, B., Baumann, U. and Rehm, H. J. 1990. Production of glycerol by immobilized *Pichia farinose*. *Appl. Microbiol. Biotechnol.* 32: 380-386.
- Boam, A. and Zhang, S. 2001. Recovering organics from aqueous streams. *Chem. Eng.* 720: 40.

- Bosma, T.N.P., Ballemans, E.M.W., Hoekstra, N.K., te Welschar, R.A.G., Smeenk, M.M., Schran, G. and Zehnder, A.J.B. 1996. Biotransformation of organics in soil columns and an infiltration area. *Ground Water*. 34: 49-56.
- Botsford, J. L., and J. G. Harman. 1992. Cyclic AMP in procaryotes. *Microbiol. Rev.* 56:100-122.
- Brockt, D., D. W. Smith, and M. T. Madigan. 1984. *Biology of Microorganisms* (41h Edition), Prentice-Hall, Inc., Englewood Cliffs, NJ.
- Brown, V.M., Jordan, D.H.M and Tiller, B.A. 1967. The effect of temperature on the acute toxicity of phenol to rainbow trout in hard water. *Water Res.* 1: 587-597.
- Bryndová, J. 2002. Aerobic phenol degradation by a yeast *Candida tropicalis* and effect of nutrition on stress conditions. (Aerobní degradace fenolu kvasinkou *Candida tropicalis* a projevy indukované nutriním stresem). Ph.D Thesis.-K. 59553/1 State Tech. Lib.Praha, Czech. (with English summary) At: <http://www.vscht.cz/obsah/fakulty/fpbt/studium/absolventi/bryndova.pdf>. accessed on 15 september 2008.
- Budavari, S. 1996. The Merck Index: An encyclopedia of chemicals, drugs and biochemicals. Whitehouse station, Merck, N.J.
- Capasso, R., Cristinzo, G., Evidente, A. and Scognamiglio, F. 1992. Isolation, spectroscopy and selective phytotoxic effects of polyphenols from vegetable waste waters. *Phytochem.* 31: 4125-4128.
- Carmichael, R.D., Jones, A., Pickard, M.A. 1986. Semicontinuous and continuous production of chloroperoxidase by *Caldariomyces fumago* Immobilized in *k*-carrageenan. *Appl. Environ. Microbiol.* Feb. p. 276-280.

- Chablain, P. A., G. Philippe, A. Groboillot, N. Truffaut, and Guespin- Michel JF. 1997. Isolation of a soil psychrotrophic toluenedegrading *Pseudomonas* strain: influence of temperature on the growth characteristics on different substrates. *Res. Microbiol.* 148: 153-161.
- Chang, S.Y., Li, C.T., Hiang, S.Y. and Chang, M.C. 1995. Intraspecific protoplast fusion of *Candida tropicalis* for enhancing phenol degradation. *Appl. Microbiol. Biotechnol.* 43: 534-538.
- Chen, K.C., Lin, Y.H., Chen, W.H. and Liu, Y.C. 2002. Degradation of phenol by PAA-immobilized *Candida tropicalis*. *Enzym. Microb. Technol.* 31: 490-497.
- Chibata, I. and Toss, T. 1981. Use of immobilized cells. *Ann. Reo. Biophys. Bioeng.* 10: 197-216.
- Cho, Y.G., Rhee, S.K. and Lee, S.T. 2000. Influence of phenol on biodegradation of *p*-nitrophenol by freely suspended and immobilized *Nocardioides* sp. NSP41. *Biodegrade.* 11: 21-28.
- Chung, T.P., Tseng, H.Y., and Jung, R.S. 2003. Mass transfer effect and intermediate detection for phenol degradation in immobilized *Pseudomonas putida* systems. *Process Biochem.* 38: 1497-1507.
- Collier, D. N., P. W. Hager, and Phibbs P.V. Jr 1996. Catabolite repression control in the *Pseudomonads*. *Res. Microbiol.* 147: 551-561.
- Collins, L.D. and Daugulis, A.J. 1997. Characterization and optimization of a two phase partitioning bioreactor for the biodegradation of phenol. *Appl. Microbiol. Biotechnol.* 48: 18-22.
- Colwell, R.R. and Walker, J.D. 1977. Ecological aspects of microbial degradation of petroleum in the marine environment. *Int. Microbiol. Rev.* 5: 423-445.

- Cort, T. and A. Bielefeldt. 2000. Effects of surfactants and temperature on PCP biodegradation. *J. Environ. Eng.* 126: 635-643.
- Crawford, D.W., Bonnevie, N.C. and Wenning, R.J. 1995. Sources of pollution and sediment contamination in Newark Bay, New Jersey. *Exotoxicol. Environ. Safety.* 35: 85-100.
- Dean-Ross, D. 1989. Bacterial abundance and activity in hazardous waste contaminated soil. *Bull. Environ. Cont. Toxicol.* 43: 511-517.
- Dean-Ross, D. and Rahimi, M. 1995. Toxicity of phenolic compounds to sediment bacteria. *Bull. Environ. Contam. Toxicol.* 55: 245-250.
- Delaney, J.L. and Hughes, T.W. 1979. *Source assessment: Manufacture of acetone and phenol from cumene.* Prepared by Mosanto Research Corp., Dayton, OH. EPA-600/2-79-019D. NTIS PB80-150592, 500.
- Den Boeft, J., Kruiswijk, F.J., and Schulting, F.L. 1984. *Air pollution by combustion of solid fuels.* The Hague, Ministry of Housing, Physical Planning and Environment. (Publication Lucht N0.37).
- Dias, J.C.T., Rezende, R.P., Linardi, V.R., 2001. Bioconversion of nitriles by *Candida guilliermondii* CCT 7207 cells immobilized in barium alginate. *Appl. Microbiol. Biotechnol.* 56: 757-761.
- Duetz, W. A., Marques, S., de Jong, C., Ramos, J. L. and van Andel, J. G. 1994. Inducibility of the TOL catabolic pathways in *Pseudomonas putida*(pWW0) growing on succinate in continuous culture: evidence of carbon catabolite repression control. *J. Bacteriol.* 176: 2354–2361.
- Efremenko, I. and Sheintuch, M. 2006. Predicting Solute Adsorption on Activated Carbon: Phenol. *Langmuir.* 22: 3614-3621.

- Ettayebi, K., Errachidi, F., Jamaï, L., Tahri-Jouti, M.A., Sendide, K. and Ettayebi, M. 2003. Biodegradation of polyphenols with immobilized *Candida tropicalis* under metabolic induction. *FEMS Microbiol. Lett.* 223: 215-219.
- Evans, W.C. 1947. Oxidation of phenol and benzoic acid by some soil bacteria. *Biol. Chem.* 41: 373-382.
- Fabregas, J., Patino, M., Morales, E.D., Cordero, B. and Otero, A. 1996. Optimal renewal rate and nutrient concentration for the production of the marine microalga *Phaeodactylum tricoratum* in semicontinuous cultures. *Appl Environ Microbiol.* 62(1): 266-8.
- Fedorak, P.M., Semple, K. M. and Westlake, D.W.S. 1984. Oil-degrading capabilities of yeasts and fungi isolated from coastal marine environments. *Can. Microbiol.* 30: 565-71.
- Feist, C. F., and G. E. Hegeman. 1969. Phenol and benzoate metabolism by *Pseudomonas putida* : Regulation of tangential pathways. *J. Bact.* 100: 869.
- Fialova, A., Boschke, E. and Bley, T. 2004. Rapid monitoring of the biodegradation of phenol like compounds by the yeast *Candida maltosa* using BOD measurements. *Int. Biodeterior. Biodegrade.* 54: 69-76.
- Figueroa, M.C. and Weber, M.E. 2005. Aqueous micellar solvent extraction of phenol from wastewater. *Sep. Sci. Technol.* 40: 1653-1672.
- Freeman, A., Lilly, M.D., 1998. Effect of processing parameters on the feasibility and operational stability of immobilized viable microbial cells. *Enzym. Microb. Technol.* 23: 335-345.

- Galindez-Mayer, J., Ramion-Gallegos, J., Ruiz-Ordaz, N., Juarez-Ramirez, C., Salmeron-Alcocer, A. and Poggi-Varaldo, H.M. 2008. Phenol and 4-chlorophenol biodegradation by yeast *Candida tropicalis* in a fluidized bed reactor. *J. Biochem. Eng.* 38: 147-157.
- Garcia, M. T. and Pena, M. 2001. Biodegradation of phenolic industrial waste water in a fluidized bed bioreactor with immobilized cells of *Pseudomonas putida*. *Bioresour. Technol.* 80: 137-142.
- Gemeiner, P. 1992. Enzyme Engineering - Immobilized Biosystems, Ellis Horwood, UK.
- Giridhar, R. and Srivastava, A. K. 2001. Repeated fed-batch sorbose fermentation by *Gluconobacter oxydans*. *Chem. Biochem. Eng.* 15(3): 127-129.
- Girolami, V., Vianello, A., Stuparon, A., Ragazzi, E. and Veronese, I. 1981. Ovipositional deterrents in *Dacus oleae*. *Entomopathog. Exp. Appl.* 29: 178-185.
- Godjevargov, T., Alexiev, Z. and Ivanov, D. 1998. Biodegradation of phenol by covalently bound *Trichosporon cutaneum* cells to porous polyamide granules. *Process Biochem.* 8: 831– 835.
- Godjevargova, T., Ivanova, D. Aleksieva, Z and Dimova, N. 2003. Biodegradation of toxic organic components from industrial phenol production waste waters by free and immobilized *Trichosporon cutaneum* R57. *Process Biochem.* 38: 915-920.
- Goerlitz, D.F., Troutman, D.E., Gody, .E.M. 1985. Migration of wood processing chemical in contaminated ground water in a sand aquifer at Pensacola, Florida. *Environ. Sci. Technol.* 19: 955-961.

- Goldstein, R.M., Mallory, L.M. and Alexander, M. (1985). Reasons for possible failure inoculation to enhance biodegradation. *Appl. Environ. Microbiol.* 50(4): 977-983.
- Goncharuk, V.V., Kucheruk, D.D., Kochkodan, V.M. and Badekha, V.P. 2002. Removal of organic substances from aqueous solutions by reagent enhanced reverse osmosis. *Desalination.* 143: 45-51.
- González, D.M., Moreno, E., Sarmiento, J.Q. and Ramos-Cormenzana, A. 1990. Studies on antibacterial activity of waste waters from olive oil mills (Alpechin): Inhibitory activity of phenolic and fatty acids. *Chemosphere.* 20: 423-432.
- González, G., Herrera, M.G., García, M.T. and Peña, M. 2001. Biodegradation of phenolic industrial wastewater in a fluidized-bed bioreactor with immobilized cells of *Pseudomonas putida*. *Biores. Technol.* 80: 137-142.
- Graedel, T.E. 1978. *Chemical compounds in the atmosphere*. New York, Academic Press, pp. 256.
- Greenberg, A.E., Cleceri, L.S. and Eaton A.D. (eds). 1992. *Standard Methods for the Examination of Water and Wastewater*. Washing DC. Pp. 5-30-5-33.
- Groenen, P.J. 1978. *Components of tobacco smoke. Nature and quantity; potential influence on health*. Zeist, The Netherlands, CIVO-TNO Institute (Rep. No. R/5787).
- Guieyesse, B., P. Wilkstrom, M. Forsman, and B. Mattiasson. 2001. Biomonitoring of continuous microbial community adaptation towards more efficient phenol degradation in a fed-batch bioreactor. *Appl. Microbiol. Biotechnol.* 56: 780-787.

- Gurujeyalekshmi, G., Oreil, P. 1988. Isolation of phenol degrading *Bacillus stearothermophilus* and partial characterization of the phenol hydroxylase. *Appl. Environ. Microbiol.* 55(2): 500-502.
- Harms, H. and Bosma, T.N.P. 1997. Mass transfer limitation of microbial growth and pollutant degradation. *Ind. Microbiol. Biotechnol.* 18: 97-105.
- Hawthorne, S.B. and Sievers, R.E. 1984. Emission of organic air pollutants from shale oil wastewaters. *Environ. Sci. Technol.* 18: 483-490.
- HazDat. 1998. *Hazardous substance database*. Agency for Toxic Substances and Disease Registry, Atlanta, G.A.
- Hickman, G.T. and Novak, J.T. 1989. Relationship between subsurface biodegradation rates and microbial density. *Environ. Sci. Technol.* 23(5): 524-532.
- Hoshino, M., and Akimoto, H. 1978. Photochemical oxidation of benzene, toluene and ethylbenzene initiated by OH radicals in the gas phase. *Bull. Chem. Soc. Jpn.* 51: 718.
- Howard, P.H. 1989. *Handbook of environmental fate and exposure data for organic chemicals*. Chelsea, Michigan, Lewis Publishers. 1: 468-476.
- HSDB 1998. *Hazardous Substances Data Bank*. National Library of Medicine, National Toxicology Information Program, Bethesda, M.D.
- Hubble, B.R., Stetter, J.R., Gebert, E., Harkness, J.B.L. and Flotard, R.D. 1981. Experimental measurements from residential wood-burning stoves. *Proc. Int. Conf. Resid. Solid Fuels: Environ. Impacts Solut.* John A A. Cooper and Dorothy Malek.

Hutchinson, D.H. and Robinson, C.W. 1988. Kinetics of the simultaneous batch degradation of p-cresol and phenol by *Pseudomonas putida*. *Appl. Microbiol. Biotechnol.* 29: 599-604.

International Agency for Research on Cancer (IARC). (1989). Phenol: In: *Some organic solvents, resin monomers and related compounds, pigments and occupational exposures in paint manufacture and painting*. Lyon, Int. Agen. Res. Cancer, pp 263-287 (IARC Monographs on the evaluation of carcinogenic risks to humans, Volume 47).

Jain, A. K., Suhas, A., Bhatnagar. 2002. Methylphenols removal from water by low-cost adsorbents. *J. Colloid Interface Sci.* 251: 39-45.

Jay, K. and Steiglitz, L. 1995. Identification and quantification of volatile organic components in emissions of waste incineration plants. *Chemosphere.* 30: 1249-1260.

Jia, X., Wen, J., Jiang, Y., Liu, X. and Feng, W. 2006. Modeling of batch phenol biodegradation in internal loop airlift bioreactor with gas recirculation by *Candida tropicalis*. *Chem. Eng. Sci.* 61: 3463-3475.

Jiang, H.L., Tay, J.H. and Tay, S.T.L. 2002. Aggregation of immobilized activated sludge cells into aerobically grown microbial granules for the aerobic biodegradation of phenol, *Lett. Appl. Microbiol.* 35: 439-445.

Jiang, H.L., Tay, J.H. and Tay, S.T.L. 2004. Changes in structure, activity and metabolism of aerobic granules as a microbial response to high phenol loading. *Appl. Microbiol. Biotechnol.* 63: 602-608.

- Jiang, Y., J. P. Wen, H. M. Li, Yang SL, and Hu ZD. 2005. The biodegradation of phenol at high initial concentration by the yeast *Candida tropicalis*. *J. Biochem Eng.* 24: 243-247.
- Jordan, W., van Barneveld, H., Gerlich, O., Kleine-Boymann, M. and Ullrich, J. 2002. Phenol, in: Ullmann's Encyclopedia of Industrial Chemistry, Wiley-VCH Verlag.
- Juarez-Ramirez, C., Riu-Ordaz, N., Cristiani-Urbina, E., and Galindez-Mayer, J. 2001. Degradation kinetics of phenol by immobilized cells of *Candida tropicalis* in a fluidized bed reactor. *World J. Microbiol. Biotechnol.* 17: 697-705.
- Kahru, A., Reiman, R. and Rätsep, A. 1998. The efficiency of different degrading bacteria and activated sludges in detoxification of phenolic leachates. *Chemosphere.* 37(2): 301-318.
- Kanekar, P.P., Sarnaik, S.S. and Kelkar, A.S. 1999. Bioremediation of phenol by alkaline bacteria isolated from alkaline lake of Lonar, India. *Appl. Microbiol.* 85: 128S-133S.
- Kao, C.M., Liu, J.K., Chen, Y.L., Chai, C.T. and Chen, S.C. 2005. Factors affecting the biodegradation of PCP by *Pseudomonas mendocina* NSYSU. *J. Hazard. Mater.* 124: 68-73.
- Kar, S., Swaminathan, T. and Baradarajan, A. 1996. Studies on biodegradation of a mixture of toxic and nontoxic pollutant using *Arthrobacter* species. *Bioprocess Eng.* 15(4): 195-199.
- Karigar, C., Mahesh, A., Nagenahalli, M. and Yun, D.J. 2006. Phenol degradation by immobilized cells of *Arthrobacter citreus*. *Biodegrade.* 17: 47-55.

- Kavitha, V. and Palanivelu, K. 2004. The role of ferrous ion in Fenton and photo-Fenton process for the degradation of phenol. *Chemosphere*. 55: 1235-1243.
- Kawachi, H., Shimizu, K., Atomi, H., Sanuki, S., Ueda, M. and Tanaka, A. 1997. Gene analysis of an NADP-linked isocitrate dehydrogenase localized in peroxisomes of the *n*-alkane-assimilating yeast *Candida tropicalis*. *Eur. Biochem.* 250: 205-211.
- Keith, L.H. • 1976. Identification of organic compounds in unbleached treated Kraft paper mill wastewaters. *Environ. Sci. Technol.* 10: 555-564.
- Kim, M.K., Singleton, I., Yin, C.R., Quan, Z.X., Lee, M. and Lee, S.T. 2005. Influence of phenol on the biodegradation of pyridine by freely suspended and immobilized *Pseudomonas putida* MK1. *Lett Appl Microb.* ISSN 0266-8254.
- Kirchner U., Westphal, A.H., Miller, M., Van Berel, W.J.H. 2003. Phenol hydroxylase from *Bacillus thermoglucosidasius* A7, a two-protein component monooxygenase with a dual role for FAD. *J. Biol Chem.* 278: 47545-47553.
- Klein, J., Hackel, U., and Wagner, F. 1979. Phenol degradation by *Candida tropicalis* whole cells entrapped in polymeric ionic networks. *ACS Symp. Ser.* 106: 101-118.
- Klibanov, A. M. 1982. Enzymatic removal of hazardous pollutants from industrial aqueous effluents. *Enzym. Eng.* 6: 319-323.
- Knoll, G. and Winter, J. 1987. Anaerobic degradation of phenol in sewage sludge: benzoate formation from phenol and carbon dioxide in the presence of hydrogen. *Appl. Microbiol. Biotechnol.* 25 (4): 384-391.

- Knupp, G., Rücker, G., Ramos-Cormenzana, A., Garrido Hoyos, S., Neugebauer, M. and Ossenkop, T. 1996. Problems of identifying phenolic compounds during the microbial degradation of olive oil mill wastewater. *Int. Biodeterior. Biodegrade.* 38: 277-282.
- Komárková, E. and Páca, J. 2000. Kinetics of phenol oxidation by *Candida tropicalis* yeast (in Czech). *Chem. Listy.* 94: 729.
- Komárková, E.; Páca, J.; Klapková, E.; Stiborová, M.; Soccol, C. R. and Sobotka, M. 2003. Physiological changes of *Candida tropicalis* population degrading phenol in fed batch reactor. *Braz. Arch. Biol. Technol.* 46: 537-542.
- Kotresha, D., and G. M. Vidyasagar. 2008. Isolation and characterization of phenol-degrading *Pseudomonas aeruginosa* MTCC 4996. *Chem. Mater Sci.* 24: 541-547.
- Krug, M., Ziegler, H. and Straube, G. 1985. Degradation of phenolic compounds by the yeast *Candida tropicalis* HP-15 : I. Physiology and growth and substrate utilization. *Basic Microbiol.* 25(2): 103-110.
- Krug, M. and Straube, G. 1986. Degradation of phenolic compounds by the yeast *Candida tropicalis* HP-15: II. Some properties of the first two enzymes of the degradation pathway. *Basic Microbiol.* 26(5): 271-281.
- Kujawski, W., Warszawski, A., Ratajczak, W., Porebski, T., Capala, W. and Ostrowska, I. 2004. Removal of phenol from wastewater by different separation techniques. *Desalination.* 163: 287-296.
- Kumar A., S. Kumar, S. Kumar 2004. Biodegradation kinetics of phenol and catechol using *Pseudomonas putida* MTCC1194. *J. Biochem. Eng.* 22: 151-159.

Kumaran, P. 1980. Microbial degradation of phenol in phenol-bearing industrial wastes. Ph.D. Thesis, Nagpur, Univ., India.

Kumaran P., and Y. L. Paruchuri. 1997. Kinetics of phenol biotransformation. *Water Res.* 31: 11-22.

Kunz, A., Reginatto, V. and Durán, N. 2001. Combined treatment of textile effluent using the sequence *Phanerochaete chrysosporium*-ozone. *Chemosphere.* 44: 281-287.

Kuwata, K., Uebori, M. and Yamazaki, Y. 1980. Determination of phenol in polluted air as p-nitrobenzene azophenol derivatives by reversed phase high performance liquid chromatography. *Anal. Chem.* 52 (6): 857-860.

Landis, W.G. and Yu, Ming-Ho. 2003. Introduction to environmental toxicology- Impacts of chemicals upon ecological systems. 3rd edition. New York, Lewis Publishers, pp. 242.

Lappin, H.M., Greaves, M.P. and Slatter, J.H. 1985. Degradation of the herbicide [2-(2-Methyl-4-Chlorophenoxy) Propionic Acid] by a synergistic microbial community. *Appl. Environ. Microbiol.* 49: 429-433.

Leahy, J.G. and Colwell, R.R. 1990. Microbial degradation of hydrocarbons in the environment. *Microbiol. Rev.* 54: 305-315.

Lee, C.M., Lu C.J. and Chuang, M.S. 1994. Effects of immobilized cells on the biodegradation of chlorinated phenols. *Water Sci. Technol.* 30(9): 87-90.

Lee, Y.G., Hwang, S.H. and Kim, S.D. 2006. Predicting the toxicity of substituted phenols to aquatic species and its changes in the stream and effluent waters. *Arch. Environ. Contam. Toxicol.* 50: 213-219.

- Lima, L.H.A., Felipe, M.G.A. and Torres, F.A.G. 2003. Reclassification of *Candida guilliermondii* FTI 20037 as *Candida tropicalis* based on molecular phylogenetic analysis. *Braz. Microbiol.* 34(1): 1-6.
- Lin, S.H., Pan, C.L. and Leu, H.G. 1999. Liquid membrane extraction of 2-chlorophenol from aqueous solution. *J. Hazard. Mater.* 65(3): 289-304.
- Lob, K. C., and P. P. Tar. 2000. Effect of additional carbon sources on biodegradation of phenol. *Bulletin Environ. Contam. Toxicol.* 64: 756-763.
- Loh, K.C., Chung, T.S. and Ang, W.F. 2000. Immobilized-cell membrane bioreactor for high-strength phenol wastewater. *J. Environ. Eng.* 126: 75-79.
- Loh K. C., and Y.G. Yu. 2000. Kinetics of carbazole degradation by *Pseudomonas putida* in presence of sodium salicylate. *Water Res.* 34: 4131-4138.
- Lyngkilde, J. and Christiansen, T.H. 1992. Redox zones of a landfill leachate pollution plume (Vejen, Denmark). *Contam. Hydrolys.* 10: 273-289.
- Manahan, S.E. 1994. *Environmental chemistry*. Boca Raton, FL: CRC Press.
- Manzano, M. A., J. A. Perales, D. Sales, and J. M. Quiroga. 1999. The effect of temperature on the biodegradation of a nonylphenol polyethoxylate in river water. *Water Res.* 33: 2593-2600.
- Margesin, R., Fonteyne, P. and Redl, B. 2005. Low-temperature biodegradation of high amounts of phenol by *Rhodococcus* spp. and basidiomycetous yeast. *Res. Microbiol.* 156: 68-75.
- Maslin, P., and R. M. Maier. 2000. Rhamnolipid enhanced mineralization of phenanthrene in organic metal co-contaminated soils. *Bioremed. J.* 4: 295-308.
- Matthews, R.W. 1992. Photocatalytic oxidation of organic contaminants in water: an aid to environmental preservation. *Pure Appl. Chem.* 64(9): 1285-1290.

- Mattiasson, B. 1983. Immobilization methods, p. 3-26. In Mattiasson B. (ed.), Immobilized cells and organelles, vol. 1. CRC Press, Inc., Boca Raton, Fla.
- McFall, S. M., B. Abraham, C. G. Narsolis, and A. M. Chakrabarty 1997. A tricarboxylic acid cycle intermediate regulating transcription of a chloroaromatic biodegradative pathway: fumarate-mediated repression of the *clcABD* operon. *J. Bacteriol.* 179: 6729-6735.
- Melcer, M. and Bridle, T.R. 1985. Discussion of metals removal and partitioning in conventional wastewater treatment plants and fate of toxic organic compounds in wastewater treatment plants. *Water Pollut. Cont. Fed.* 57: 263-264.
- Melin, E. S., J.T. Kimmo, and J. A. Puhakka. 1998. Effects of temperature on chlorophenol biodegradation kinetics in fluidized-bed reactors with different biomass carriers. *Water Res.* 32: 81-90.
- Minussi, R.C., Pastore, G.M. and Durán, N. 1998. Biotreatment of paper and pulp effluent using fungal mediators: ABTS and HBT, in: Arce, J., and Velaquez, C.I. (Eds.), *Proc. of the XIII Latinamerican Chemical Congress Rio Grande, Puerto Rico*, QAM-M-8.
- Mordocco, A., Kuek, C. and Jenkins, R. 1999. Continuous degradation of phenol at low concentration using immobilized *Pseudomonas putida*. *Enzym. Microbiol. Technol.* 25: 530-536.
- Movahedian H., R. Shokoohi, A. Parvaresh, M. Hajia, and A.J. Jafari. 2005. Evaluating the Effect of Glucose on phenol Removal Efficiency and Changing the Dominant Microorganisms in a Serial Combined Biological System. *J. Biol Sci.* 11: 1491-1494.

- Muller, C., L. Petruschka, H. Cuypers, G. Burchhardt, and H. Herrmann 1996. Carbon catabolite repression of phenol degradation in *Pseudomonas putida* is mediated by the inhibition of the activator protein PhIR. *J. Bacteriol.* 178: 2030-2036.
- Murzakov, B., Akopova, G. and Kruglova, N. 2003. The technology of bioremediation of oil polluted objects by biopreparation (Project "Biodestructor"). *Proc. 17th Forum for Applied Biotechnology.* 18-19 Sept. 2003, Belgium, Gent.
- Musto, J.D., Sane, J.N. and Warner, V.D. 1977. Quantitative determination of phenol by high-performance liquid chromatography. *Pharm. Sci.* 66: 1201-1202.
- Nakamura, Y., and T. Sawada, 2000. Biodegradation of phenol in the presence of heavy metals. *J. Chem. Technol. Biotechnol.* 75: 137-142.
- Nair, C., Jayachandran, I. K. and Shashidhar, S. 2008. Biodegradation of phenol. *Afr. J. Biotechnol.* 7(25): 4951-4958.
- Neujahr, H.Y. and Varga, J.M. 1970. Degradation of phenols by intact cells and cell-free preparations of *Trichosporon cutaneum*. *Eur. Biochem.* 13: 37-44.
- Neujahr, H.Y., Lindsjo, S. and Varga, J.M. 1974. Oxidation of phenol by cells and cell-free enzymes from *Candida tropicalis*. *Antonie van Leeuwenhoek.* 40(2): 209-216
- Oltmanns, R.H., Muller, R., Otto, M.K. and Lingens, F. 1989. Evidence for a new pathway in the bacterial degradation of 4-fluorobenzoate. *Appl. Environ. Microbiol.* 55: 2499-2504.

- Palepu, D.T., Chauhan, S.P. and Amanth, K. P. 1995. Separation technologies. In: H.M. Freeman, Editor, *Industrial Pollution Prevention Handbook* (1st ed.). McGraw-Hill, New York.
- Parales, R.E., Bruce, N.C., Schmid, A. and Wackett, L.P. 2002. Biodegradation, biotransformation, and biocatalysis (B3). *Appl. Environ. Microbiol.* 68(10): 2699-4709.
- Paraskevi N. P., and G. S. Euripides. 2005. Effect of temperature and additional carbon sources on phenol degradation by an indigenous soil *Pseudomonad*. *J. Biodegrade.* 16: 403-413.
- Parkhurst, B.R., Bradshaw, A.S., and Forte, J.L. 1979. An evaluation of the acute toxicity to aquatic biota of a coal conversion effluent and its major components. *Bull. Environ. Contam. Toxicol.* 23: 349-356.
- Patterson, J.W. 1985. *Industrial wastewater treatment technology*. Butterworths, Boston.
- Pekari, K., Vainotalo, S., Heikkila, P., Palotie, A., Luotamo, M. and Riihimaki, V. 1992. Biological monitoring of occupational exposure to low levels of benzene. *Scand. Work Environ. Health.* 18: 317-322.
- Peralta-Zamora, P., Gimenes, L.F., Cordi, L., Reyes, J., Alves, O.L. and Durán, N. 1998. Remediation of effluents from pulp and paper industry using horseradish peroxidase immobilized on ceramic materials, in: Gaylarde, C.C., Barbosa, T.C.P., Gabilan, N.H. (Eds.), *Proc. 3rd Lat. Am. Biodegradation and Biodeterioration Symposium-LABS-3*, Florianopolis, Brazil, CD-Rom Paper 6, p.1.

- Percival, L.J. and Senior, E. 1998. An assessment of the effects of the dual codisposal of phenol and waste activated sewage sludge with refuse on the refuse anaerobic fermentation and leachate quality. ISSN 0378-4738-*Water SA*24. 1: 57-70.
- Peters, M., Heinaru, E., Talpsep, H., Ward, H., Stottmeister, U., Heinaru, A. and Nurk, A. 1997. Acquisition of a deliberately introduced phenol degradation operon, *pheAB*, by different indigenous *Pseudomonas* species. *Appl. Environ. Microbiol.* 63: 4899-4906.
- Pfeffer, F.M. 1979. The 1977 screening survey for measurement of organic priority pollutants in petroleum refinery wastewaters. *ASTM Spec. Tech.Publ.* 181-190.
- Pradhan N.C., Sarkar, C.S., Niyogi, S. and Adhikari, B. 2002. Separation of phenol-water mixture by membrane pervaporation using polyimide membranes, *J. Appl. Polym. Sci.* 83: 822-829.
- Prieto, M.B., Hidalgo, A., Serra, J.L. and Llama M.J. 2002. Degradation of phenol by *Rhodococcus erythropolis* UPV-1 immobilized on Biolite® in a packed-bed reactor. *Biotechnol.* 97: 1-11.
- Radmann, E.M., Reinehr, C.O. and Costa, J.A.V. 2007. Optimization of the repeated batch cultivation of microalga *Spirulina platensis* in open raceway ponds. *Aqua.* 265(1-4): 118-26.
- Ralston, J. R., and G. R. Vela. 1974. A medium for detecting phenol-degrading bacteria. *J. Appl. Bact.* 37: 347-351.

- Rengaraj, S., Seung-hyeon, M., Sivabalan, R., Arabindoo, B., and Murugesan, V. 2002. Agricultural solid waste for the removal of organics: adsorption of phenol from water and wastewater by palm seed coat activated carbon. *Waste Manage.* 22: 543-548.
- Roane, T. M., K. L. Josephson, and I. L. Pepper. 2001. Microbial cadmium detoxification allows remediation of co-contaminated soil. *Appl. Environ. Microbiol.* 67: 3208-3215.
- Robards, K. and Ryan, D. 1998. Phenolic compounds in olives. *Analyst* 123:31R-44R.
- Rozich, A.F. and Colvin, R.J. 1986. Effect of glucose on phenol biodegradation by heterogeneous populations. *Biotechnol. Bioeng.* 28: 965-971.
- Ruiz-Ordaz, N., Juarez Ramirez, J.C., Castanon-Gonzalez, H., Lara-Rodriguez, A.R., Christiani-Urbina, E. and Galindez-Mayer, J. 2000. Aerobic bioprocesses and bioreactors used for phenol degradation by free and immobilized cells. In: Pandalai, S.G. (ed.) *Recent Research Developments in Biotechnol. and Bioeng.*, Res. Signpost. Triv. India. pp. 83-94.
- Ryan, D.R., Leukes, W.D. and Burton, S.G., 2005. Fungal bioremediation of phenolics wastewaters in an airlift reactor. *Biotechnol. Prog.* 21: 1066-1074.
- Sandrin, T.R., Chech, A.M. and Maier, R.M. 2000. A rhamnolipid biosurfactant reduces cadmium toxicity during biodegradation of naphthalene. *Appl. Environ. Microbiol.* 66(10): 4585-4588.
- Saier, M. H., Jr., S. Chauvaux, G. M. Cook, J. Deutscher, I. T. Paulsen, J. Reizer, and J.-J. Ye. 1996. Catabolite repression and inducer control in gram-positive bacteria. *Microbiol.* 142: 217-230.

- Samanta, S. K., B. Bhushan, and R. K. Jain 2001. Efficiency of naphthalene and salicylate degradation by a recombinant *Pseudomonas putida* mutant strain defective in glucose metabolism. *Appl. Microbiol. Biotechnol.* 55: 627-631.
- Santos, V. L., N. M. Heilbuth, V. R. Linardi. 2001. Degradation of phenol by *Trichosporum* sp. LE3 cells immobilized in alginate. *J. Basic Microbiol.* 41 (3-4): 171-178.
- Santos, V.L. and Linardi, V.R. 2004. Biodegradation of phenol by a filamentous fungi isolated from industrial effluents-identification and degradation potential. *Process Biochem.* 39: 1001-1006.
- Santos, V.L., Souza Monteiro, A., Telles Braga, D. and Santoro, M.M. 2009. Phenol degradation by *Aureobasidium pullulans* FE13 isolated from industrial effluents. *J. Hazard. Mater.* 161: 1413-1420.
- Saxena, S.C. and Jotshi, C.K. 1996. Management and combustion of hazardous wastes. *Prog. Energy Combust. Sci.* 22: 401-425.
- Schleissner, C., E. R. Olivera, M. Fernandez-Valverde, and J. M. Luengo. 1994. Aerobic catabolism of phenylacetic acid in *Pseudomonas putida* U: biochemical characterization of a specific phenylacetic acid transport system and formal demonstration that phenylacetyl-coenzyme A is a catabolic intermediate. *J. Bact.* 176: 7667-7676.
- Scow, K., Goyer, M., and Payne, E. 1981. *Exposure and risk assessment for phenol* (revised). Prepared for U.S. Environmental Protection Agency, Office of Water Regulations and Standards, Washington, D.C., NTIS PB85-221695, 114-116.
- Seetharam, G.B. and Saville, B.A. 2003. Degradation of phenol using tyrosinase immobilized on siliceous supports. *Water Res.* 37: 436-440.

- Shawabken, R., K. M. khleifat, I. Al-Majali, and K. tarawrawneh. 2007. Rate of biodegradation of phenol by *klebsiella oxytoca* in minimal medium and nutrient broth conditions. *J. Biorem.* 11(1): 13-19.
- Sheeja, R.Y. and Murugesan, T. 2002. Mass transfer studies on the biodegradation of phenols in up-flow packed bed reactors. *Hazard. Mater.* B89: 287-301.
- Shimizu, T., Uno, T., Dan, Y., Nei, N. and Ichikawa, K. 1973. Continuous treatment of wastewater containing phenol by *Candida tropicalis*. *Ferm. Technol.* 51: 804-812.
- Shimp, R.J. and Young, R.L. 1987. Availability of organic chemicals for biodegradation in settled bottom sediments. *Exotoxicol. Environ. Safe.* 15(1):31-45.
- Shuler, M.L. and Kargi, F. 1992. *Bioprocess Engineering*, Prentice Hall, Englewood Cliffs, NJ.
- Soda, S., M. Ike, and M. Fujita. 1998. Effects of inoculation of a genetically engineered bacterium on performance and indigenous bacteria of a sequencing batch activated sludge process treating phenol. *J. Ferment. Bioeng.* 86: 90-96.
- Spain, J.C. and van Veld, P.A. 1983. Adaptation of natural microbial communities to degradation of xenobiotic compounds: effects of concentration, exposure time, inoculum, and chemical structure. *Appl. Environ. Microbiol.* 53: 1010-1019.
- Spoelstra, S.F. 1978. Degradation of tyrosine in anaerobically stored piggery wastes and in pig faeces. *Appl. Environ. Micobiol.* 36: 631-638.
- Stanton, W.R. and DaSilva. 1978. GIAM V. Global impacts of applied microbiology. State of the Art: GIAM and its relevance to developing countries. Univ. Malaya Press, Kuala Lumpur.

Stephenson, T. 1990. Substrate inhibition of phenol oxidation by a strain of *Candida tropicalis*. *Biotechnol. Lett.* 12: 843-846.

Stephen T. L. T., Y. P. M. Benjamin, M. M. Abdul, and H. T. Joo. 2005. Comparing activated sludge and aerobic granules as microbial inocula for phenol biodegradation. *Appl. Microbiol. Biotechnol.* Online 13 January.

Suseela, G.R., . Basu, S.K., Nandy, S.C. 1991. Degradation of pentachlorophenol by *Pseudonimas aeruginosa*. *Ind. J Environ. Health.* 33(4):425-432.

Swindoll, C.M., Aelion, C.M. and Pfaender, F.K. 1988. Influence of inorganic and organic nutrients on aerobic biodegradation and on the adaptation response of subsurface microbial communities. *Appl. Environ. Microbiol.* 54: 212–217.

Symons, J.M., and McKinney, K.E. 1995. The biochemistry of nitrogen in the synthesis of activated sludge sewage. *Ind. Wastes.* 30: 875-879.

Todd, R. S. and R. H. Douglas. *Bioremediation of Organic and Metal Co-contaminated Environments: Effects of Metal Toxicity, Speciation, and Bioavailability on Biodegradation*. Department of Biology and Microbiology, University of Wisconsin Oshkosh, Oshkosh, Wisconsin, USA.

Tornai-Lehoczki, J., Péter, G. and Dlačny, D. 2003. CHROM agar *Candida* medium as a practical tool for the differentiation and presumptive identification of yeast species isolated from salads. *Int. Food Microbiol.* 86: 189-200.

TRI 1998. *Toxic Chemical Release Inventory*. National Library of Medicine, National Toxicology Information Program, Bethesda, M.D.

Tsai, S.C., Tsai, L.D. and Li, Y.K. 2005. An isolated *Candida albicans* TL3 capable of degrading phenol at large concentration. *Biosci. Biotechnol. Biochem.* 69: 2358-2367.


- US Environmental Protection Agency. 1980. Phenol. Ambient water quality criteria. Washington, D.C., USEPA (EPA 440/5-80-066).
- US Environmental Protection Agency, 2000. Guidelines for Reproductive Toxicity Risk Assessment: Notice dated October, 2000. Cas No. 108-95-2.
- Van der Meer, J.R., de Vos, W.M., Harayama, S. and Zehnder, A.J.B. 1992. Molecular mechanism of genetic adaptation to xenobiotic compounds. *Microbiol. Rev.* 56: 677-694.
- van Schie, P.M. and Young, L.Y. 2000. Biodegradation of phenol: mechanisms and applications. *Bioremed.* 4: 1-18.
- Varga, J. M., and H. Y. Neujahr, 1970. Purification and properties of catechol 1,2-oxygenase from *Trichosporon cutaneum*. *Eur. J. Biochem.* 12: 427-434.
- Verschuere, K. 1983. *Handbook of Environmental Data on Organic Chemicals*. 2nd ed. Van Nostrand Reinhold Co., New York.
- Vijaikishore, P. and Karanth, N.G. 1986. Glycerol production by immobilized cells of *Pichia farinose*. *Biothchnol Lett.* 8: 257-260.
- Vojta, V., Nahlik, J., Paca, J. and Komarkova, E. 2002. Development and verification of the control system for fed-batch phenol degradation process. *Chem. Biochem. Eng.* 16(2): 59-67.
- Wang, S. J., and K.C. Loh. 2000. Growth kinetics of *Pseudomonas putida*. *Biotechnol. Bioeng.* 68 (4): 437-447.
- Wang, Y., Tian, Y., Bin, H., Hua-bing, Z., Jian-nan, B. and Bao-li, C. 2007. Biodegradation of phenol by free and immobilized *Acinetobacter* sp. strain PD12. *J. Environ. Sci.* 19: 222-225.

- Whiteley, A.S., Wiles, S., Lilley, A.K., Philip, J. and Bailey, M.J. 2001. Ecological and physiological analyses of *Pseudomonads* species within a phenol remediation system. *Microbiol. Meter.* 44: 79-88.
- WHO. Phenol. Environmental Health Criteria-EHC 161. World Health Organization. Geneva. 1994.
- Wu, Z., Wang, Y. and Xing, Z. 2005. Study on degrading phenol by immobilized *Ralstonia metallidurans* CH34. *J. Microbiol.* 32(4): 31-36.
- Xing, B., McGill, W.B. and Dudas, M.J. 1994. Sorption of phenol by selected polymers: Isotherms, energetics, and polarity. *Environ. Sci. Technol.* 28: 466-473.
- Yamane, T., 1995. In: Bioreactor System Design (Asenjo, J.A. and Merchuk, J. C.) p. 481
- Yan, J., Jianping, W., Hongmei, L., Suliang, Y. and Zongding, H. 2005. The biodegradation of phenol at high initial concentration by yeast *Candida tropicalis*. *Biochem. Eng.* 24(3): 243-247.
- Yan, J., Jianping, W., Bai, J., Daoquan, W. and Zongding, H., 2006. Phenol biodegradation by the yeast *Candida tropicalis* in the presence of m-cresol. *J. Biochem. Eng.* 29: 223-227.
- Yang, C.F. and Lee, C.M. 2007. Enrichment, isolation, and characterization of phenol-degrading *Pseudomonas resinovorans* strain P-1 and *Brevibacillus* sp. strain P-6. *Int. Biodeterior. Biodegrade.* 59: 206-210.
- Yap, L.F., Lee, Y.K. and Poh, C.L. 1999. Mechanism for phenol tolerance in phenol-degrading *Comamonas testosteroni* strain. *Appl. Microbiol. Biotechnol.* 5(6): 833-840.

Yurii, I.M.M., and Moshe, S. 1998. Catalytic abatement of water pollutants. *Ind. Eng. Chem. Res.* 37: 309-326.

Zaitsev, G.M., Uotila, J.S., Lobanov, A.G. and Salkinoja-Salonen, M.S. 1995. Utilization of halogenated benzenes, phenols, and benzoates by *Rhodococcus opacus* GM-14. *Appl. Environ. Microbiol.* 61: 4191-4201

Zaki, S., Abd-El-Haleem, D., Abulhamd, A., Elbery, H. and AbuElreesh, G. 2008. Influence of phenolics on the sensitivity of free and immobilized bioluminescent *Acinetobacter bacterium*. *J. Hazard. Mater.* 163: 277-285.



ลิขสิทธิ์มหาวิทยาลัยเชียงใหม่
Copyright© by Chiang Mai University
All rights reserved