

TABLE OF CONTENTS

	Page
ACKNOWLEDGEMENTS	iii
ABSTRACT (ENGLISH)	iv
ABSTRACT (THAI)	v
LIST OF TABLES	ix
LIST OF FIGURES	x
ABBREVIATIONS AND SYMBOLS	xv
CHAPTER 1 INTRODUCTION	1
1.1 General aspects of zinc oxide and its applications	2
1.1.1 Physical properties	2
1.1.1.1 Crystal structure of zinc oxide	2
1.1.1.2 Mechanical properties	4
1.1.1.3 Electronic properties	4
1.1.2 Applications	5
1.2 Zinc Oxide processing route	11
1.2.1 Microwave irradiation method	11
1.2.2 Ultrasonic irradiation method	11
1.2.3 Hydrothermal/Solvothermal method	13
1.2.4 Template-based method	15
1.3 Surfactant	18
1.4 Literature Review	23

	Page
CHAPTER 2 EXPERIMENTAL PROCEDURE	31
2.1 Chemical reagents and equipments	31
2.1.1 Chemical reagents	31
2.1.2 Equipments	31
2.2 Synthesis method	32
2.2.1 Synthesis of zinc oxide via a microwave radiation method	32
2.2.1.1 Effect of liquid media	32
2.2.1.2 Effect of surfactants	33
2.2.1.3 Effect of the amount of PEG20000	34
2.2.2 Synthesis of zinc oxide via ultrasonic radiation method	35
2.2.2.1 Effect of Zn(NO ₃) ₂ to NaOH ratios	35
2.2.2.2 Effect of reaction time	37
2.2.2.3 Effect of the amount of PEG20000	38
2.3 Characterization	39
2.3.1 X-ray Diffraction (XRD)	39
2.3.2 Scanning Electron Microscopy (SEM)	40
2.3.3 Transmission Electron Microscopy (TEM)	41
2.3.4 Photoluminescence Spectroscopy	42
CHAPTER 3 RESULTS AND DISCUSSION	43
3.1 Result of the products synthesized by	43
a microwave radiation method.	
3.1.1 Effect of liquid media	43

	Page
3.1.2 Effect of surfactants	54
3.1.3 Effect of the amount of PEG20000	58
3.2 Result of products synthesized by a ultrasonic radiation method	65
3.2.1 Effect of Zn(NO ₃) ₂ to NaOH ratios	65
3.2.2 Effect of reaction time	72
3.2.3 Effect of the amount of PEG20000	75
CHAPTER 4 CONCLUSION	79
4.1 Synthesis of zinc oxide via a microwave radiation method	79
4.2 Synthesis of zinc oxide via ultrasonic radiation method	80
REFERENCES	81
APPENDICES	86
APPENDIX A ZnO JCPDS used for the present research	87
APPENDIX B ZnO Material Safety Data Sheet	91
APPENDIX C Camera constant used for the indexing of SAED pattern	97
CURRICULUM VITAE	98

LIST OF TABLES

Table	Page
1.1 Surfactant classification	21
3.1 Ring diffraction pattern values of products prepared using water as liquid media.	52
3.2 Ring diffraction pattern values of products prepared using PEG200 as liquid media.	52

ลิขสิทธิ์มหาวิทยาลัยเชียงใหม่
 Copyright© by Chiang Mai University
 All rights reserved

LIST OF FIGURES

Figure	Page
1.1 Crystal structures of ZnO	3
1.2 General purpose autoclave popularly used for hydrothermal synthesis	14
1.3 Pressure temperature map of materials processing techniques	15
1.4 Schematic illustration of the micelle monomers	19
1.5 Schematic illustration of the commonly observed geometrical shapes of surfactant micelles in aqueous solution	23
2.1 Schematic diagram used for preparation of ZnO using H ₂ O, PEG200, PEG400, PEG600 as liquid media.	33
2.2 Schematic diagram used for preparation of ZnO using 2g PEG6000, PEG10000, PEG20000 and 6g, 10g, 20g PEG20000 as surfactant.	34
2.3 Schematic diagram used for preparation of ZnO using different Zn(NO ₃) ₂ to NaOH ratios.	36
2.4 Schematic diagram used for preparation of ZnO using different prolonged time.	37
2.5 Schematic diagram used for preparation of ZnO using different amount of PEG20000 as surfactant.	38
2.6 X-ray diffractometer	39
2.7 Scanning electron microscope	40
2.8 Transmission electron microscope	41
2.9 Luminescence spectrometer	42

Figure	Page
3.1 XRD patterns of the products synthesized by a microwave radiation using H ₂ O, PEG200, PEG400, PEG600 as liquid media.	43
3.2 SEM image of the sample prepared using Zn(NO ₃) ₂ :NaOH = 1:15 as starting reagents in water solution.	45
3.3 SEM image of the sample prepared using Zn(NO ₃) ₂ :NaOH = 1:15 as starting reagents in PEG200 solution.	45
3.4 SEM image of the sample prepared using Zn(NO ₃) ₂ :NaOH = 1:15 as starting reagents in PEG400 solution.	46
3.5 SEM image of the sample prepared using Zn(NO ₃) ₂ :NaOH = 1:15 as starting reagents in PEG600 solution.	46
3.6 TEM images and SAED pattern of the product prepared using Zn(NO ₃) ₂ :NaOH = 1:15 as starting reagents in water solution.	47
3.7 TEM images and SAED pattern of the product prepared using Zn(NO ₃) ₂ :NaOH = 1:15 as starting reagents in PEG200 solution.	48
3.8 TEM images, SAED and simulated pattern of the product prepared using Zn(NO ₃) ₂ :NaOH = 1:15 as starting reagents in PEG400 solution.	49
3.9 TEM images, SAED and simulated pattern of the product prepared using Zn(NO ₃) ₂ :NaOH = 1:15 as starting reagents in PEG600 solution.	50
3.10 PL spectra of the products synthesized using a microwave radiation in H ₂ O, PEG200, PEG400 and PEG600 solution.	53
3.11 XRD patterns of the products synthesized by a microwave radiation using PEG6000, PEG10000, PEG20000 as surfactant.	54

Figure	Page
3.12 SEM image of the sample prepared using $Zn(NO_3)_2:NaOH = 1:15$ as starting reagents with PEG6000.	55
3.13 SEM image of the sample prepared using $Zn(NO_3)_2:NaOH = 1:15$ as starting reagents with PEG10000.	55
3.14 SEM image of the sample prepared using $Zn(NO_3)_2:NaOH = 1:15$ as starting reagents with PEG20000.	56
3.15 PL spectra of the products synthesized by a microwave radiation using $Zn(NO_3)_2:NaOH = 1:15$ as starting reagents with PEG6000, PEG10000 and PEG20000 as surfactant.	57
3.16 XRD patterns of the products synthesized by a microwave radiation using PEG20000 at 6, 10, 20 g as surfactant.	58
3.17 SEM image of the product prepared using $Zn(NO_3)_2:NaOH = 1:15$ as starting reagents using amount PEG 20000.	59
3.18 SEM image of the product prepared using $Zn(NO_3)_2:NaOH = 1:15$ as starting reagents using PEG 20000.	59
3.19 SEM image of the product prepared using $Zn(NO_3)_2:NaOH = 1:15$ as starting reagents using PEG 20000.	60
3.20 TEM images, SAED and simulated patterns of the product synthesized using PEG 20000 as a surfactant.	61
3.21 TEM images, SAED and simulated patterns of the product synthesized using PEG 20000 as a surfactant.	62


Figure	Page
3.22 PL spectra of the products synthesized by a microwave radiation using $\text{Zn}(\text{NO}_3)_2:\text{NaOH} = 1:15$ as starting reagents with 6 g, 10 g and 20 g PEG20000 as surfactant.	64
3.23 XRD patterns of the samples synthesized using different $\text{Zn}(\text{NO}_3)_2$ to NaOH ratios.	65
3.24 SEM image of the product prepared using 1:5 $\text{Zn}(\text{NO}_3)_2$ to NaOH ratio.	66
3.25 SEM image of the product prepared using 1:10 $\text{Zn}(\text{NO}_3)_2$ to NaOH ratio.	66
3.26 SEM image of the product prepared using 1:15 $\text{Zn}(\text{NO}_3)_2$ to NaOH ratio.	67
3.27 SEM image of the product prepared using 1:20 $\text{Zn}(\text{NO}_3)_2$ to NaOH ratio.	67
3.28 TEM images, SAED and simulated patterns of the product synthesized using $\text{Zn}(\text{NO}_3)_2$ to NaOH ratio.	69
3.29 TEM images, SAED and simulated patterns of the product synthesized using $\text{Zn}(\text{NO}_3)_2$ to NaOH ratio.	71
3.30 XRD patterns of the samples synthesized using different prolonged times.	72
3.31 SEM image of the sample synthesized using 1:10 $\text{Zn}(\text{NO}_3)_2$ to NaOH ratio.	73
3.32 SEM image of the sample synthesized using 1:10 $\text{Zn}(\text{NO}_3)_2$ to NaOH ratio.	73
3.33 SEM image of the sample synthesized using 1:10 $\text{Zn}(\text{NO}_3)_2$ to NaOH ratio.	74
3.34 XRD patterns of the samples synthesized using different amount of PEG20000.	75
3.35 SEM image of the sample synthesized using PEG20000.	76
3.36 SEM image of the sample synthesized using PEG20000.	76
3.37 SEM image of the sample synthesized using PEG20000.	77

Figure

Page

3.38 SEM image of the sample synthesized using PEG20000.

77



ลิขสิทธิ์มหาวิทยาลัยเชียงใหม่
Copyright© by Chiang Mai University
All rights reserved

ABBREVIATIONS AND SYMBOLS

Å	=	Angstrom
°C	=	Degree Celsius
eV	=	electron Volt
kV	=	Kilo Volt
g	=	Gram
h	=	Hour
mmol	=	Millimol
ml	=	Milliliter
mm	=	Millimeter
nm	=	Nanometer
µm	=	Micrometer
JCPDS	=	Joint Committee for Powder Diffraction Standards
MW	=	Molecular Weight
PL	=	Photoluminescence Spectrometry
SAED	=	Selected Area Electron Diffraction
SEM	=	Scanning Electron Microscope
TEM	=	Transmission Electron Microscope
W	=	Watt
XRD	=	X-ray Diffraction Spectrometry

ลิขสิทธิ์มหาวิทยาลัยเชียงใหม่
 Copyright © by Chiang Mai University
 All rights reserved