

REFERENCES

- [1] Haffner, S.M.; Mykkanen L.; Festa, A.; Burke, J.P. & Stern, M.P. (2000). Insulin-resistant prediabetic subjects have more atherogenic risk factors than insulin-sensitive prediabetic subjects: implications for preventing coronary heart disease during the prediabetic state. *Circulation*, 101(9), 975-980.
- [2] Ceriello, A. (2003). New insights on oxidative stress and diabetic complications may lead to a "causal" antioxidant therapy. *Diabetes Care*, 26, 1589-1596.
- [3] Khan, Z. A., & Chakrabarti, S. (2003). Growth factors in proliferative diabetic retinopathy. *Exp Diabetes Res*, 4(4), 287-301.
- [4] Baynes, J.W. (1991). Role of oxidative stress in development of complications in diabetes. *Diabetes*, 40, 405-412.
- [5] Ceriello, A. (2006). Effects of macronutrient excess and composition on oxidative stress: relevance to diabetes and cardiovascular disease. *Curr Atheroscler Rep*, 8, 472-476.
- [6] Chertow, B. (2004). Advances in diabetes for the millennium: vitamins and oxidant stress in diabetes and its complications. *MedGenMed*, 6(Suppl), 4.

- [7] Da Ros, R.; Assaloni, R. & Ceriello, A. (2004). Antioxidant therapy in diabetic complications: what is new? *Curr Vasc Pharmacol*, 2, 335-341.
- [8] Monnier, L.; Mas, E.; Ginet, C.; Michel, F.; Villon, L.; Cristol, J.P. & Colette, C. (2006). Activation of oxidative stress by acute glucose fluctuations compared with sustained chronic hyperglycemia in patients with type 2 diabetes. *JAMA*, 295, 1681-1687.
- [9] Evans, J.L.; Goldfine, I.D.; Maddux, B.A & Grodsky, G.M. (2003). Are Oxidative Stress-Activated Signaling Pathways Mediators of Insulin Resistance and Beta-Cell Dysfunction? *Diabetes*, 52(1), 1-8.
- [10] Aruoma, O.I.; Neergheen, V.S.; Bahorun, T.; & Jen, L. (2007). Free Radicals, Antioxidants and diabetes: embryopathy, retinopathy, neuropathy, nephropathy and Cardiovascular Complications. *Neuroembryol Aging*, 4, 117-137.
- [11] Chandra, A.; Mahdi, A.A.; Ahmed, S. & Singh, R.K. (2007). Indian herbs result in hypoglycemic responses in streptozotocin-induced diabetic rats. *Nutrition Research*, 27, 161-168.
- [12] Sabu, M.C. & Kuttan, R. (2004). Antidiabetic activity of Aegle marmelos and its relationship with its antioxidant properties. *Indian J Physiol Pharmacol*, 48, 81-88.
- [13] Rudnicki, M.; Oliveira, M.R.; Pereira, T.V.; Reginatto, F.H.; Dal-Pizzol, F. & Moreira, J.C.F. (2007). Antioxidant and antiglycation properties

of *Passiflora alata* and *Passiflora edulis* extracts. *Food Chemistry*, 100, 719-724.

- [14] Pan, Y.; Zhu, J.; Wang, H.; Zhang, X.; Zhang, Y.; He, C.; Ji, X. & Li, H. (2007). Antioxidant activity of ethanolic extract of *Cortex fraxini* and use in peanut oil. *Food Chemistry*, 103, 913-918.
- [15] Tang, S.Y.; Whiteman, M.; Peng, Z.F.; Jenner, A.; Yong, E.L. & Halliwell, B. (2004). Characterization of antioxidant and antiglycation properties and isolation of active ingredients from traditional Chinese medicines. *Free Radical Biology and Medicine*, 36, 1575-1587.
- [16] Su, B.N.; Pawlus, A.D.; Jung, H.A.; Keller, W.J.; McLaughlin, J.L. & Kinghorn, A.D. (2005). Chemical Constituents of the Fruits of *Morinda citrifolia* (Noni) and Their Antioxidant Activity. *J Nat Prod*, 68, 592-595.
- [17] Leong, L.P. & Shui, G. (2002) An investigation of antioxidant capacity of fruits in Singapore markets. *Food Chemistry*, 76, 69-75.
- [18] Pellegrini, N.; Serafini, M.; Colombi, B.; Del Rio, D.; Salvatore, S.; Bianchi, M. & Brighenti, F. (2003). Total antioxidant capacity of plant foods, beverages and oils consumed in Italy assessed by three different in vitro assays. *J Nutr*, 133, 2812-2819.
- [19] Saito, K.; Kohno, M.; Yoshizaki, F. & Niwano, Y. (2008). Extensive screening for edible herbal extracts with potent scavenging activity against superoxide anions. *Plant Foods Hum Nutr*, 63, 65-70.

- [20] Meng, J.; Leung, K.S.Y.; Jiang, Z.H.; Dong, X.P. & Zhao, Z.Z. (2005). Establishment of HPLC-DAD-MS fingerprint of fresh *Houttuynia cordata*. *Chem Pharm Bull*, 53, 1604-1609.
- [21] Li, W.L.; Zheng, H.C.; Bukuru, J. & De Kimpe, K. (2004). Review natural medicines used in the traditional Chinese medical system for therapy of diabetes mellitus. *J Ethnopharmacol*, 92, 1-21.
- [22] Kerdonphag, J. (2005). The medicinal plants therapy in diabetes mellitus. 1st Editor, *Seven printing group limited*, Bangkok.
- [23] Ke, B. & Higa, T. (2000). Theories and Mechanisms of Antioxidant Action of EM-X. *EM Medical Revolution*, 2, 42-56.
- [24] Schubert, S. Y.; Lansky, E. P. & Neeman, I. (1999). Antioxidant and eicosanoid enzyme inhibition properties of pomegranate seed oil and fermented juice flavonoids. *J Ethnopharmacol*, 66(1), 11-17.
- [25] Lansky, E.P. & Newman, R.A. (2007). *Punica granatum* (pomegranate) and its potential for prevention and treatment of inflammation and cancer. *J Ethnopharmacol*, 109(2), 177-206.
- [26] Cho, E.J.; Yokozawa, T.; Kim, H.Y.; Shibahara, N. & Park, J.C. (2004). *Rosa rugosa* attenuates diabetic oxidative stress in rats with streptozotocin-induced diabetes. *Am J Chin Med*, 32(4), 487- 496.
- [27] El-Alfy AT, Ahmed AAE, Fatani AJ. (2005). Protective effect of red grape seeds proanthocyanidins against induction of diabetes by alloxan in rats. *Pharmacological Research*, 52(3), 264-270.

- [28] Chanwitheesuk, A.; Teerawutgulrag, A. & Rakariyatham, N. (2005). Screening of antioxidant activity and antioxidant compounds of some edible plants of Thailand. *Food Chemistry*, 92, 491-497.
- [29] Maisuthisakul, P.; Suttajit, M. & Pongsawatmanit, R. (2007). Assessment of phenolic content and free radical-scavenging capacity of some Thai medicinal plants. *Food Chemistry*, 100, 1409-1418.
- [30] Huang, Z.; Wang, B.; Eaves, D.H.; Shikany, J.M. & Pace, R.D. (2007). Phenolic compound profile of selected vegetables frequently consumed by African Americans in the southeast United States. *Food Chemistry*, 103, 1395-1402.
- [31] Chiou, A.; Karathanos, V.T.; Mylona, A.; Salta, F.N.; Preventi, F. & Andrikopoulos, N.K. (2007). Currants (*Vitis vinifera* L.) content of simple phenolics and antioxidant activity. *Food Chemistry*, 102, 516-522.
- [32] Maisuthisakul, P.; Pongsawatmanit, R. & Gordon, M.H. (2007). Characterization of the phytochemicals and antioxidant properties of extracts from Teaw (*Cratoxylum formosum* Dyer). *Food Chemistry*, 100, 1620-1629.
- [33] Ferreira, I.C.F.R.; Baptista, P.; Vilas-Boas, M. & Barros, L. (2007). Free-radical scavenging capacity and reducing power of wild edible mushrooms from northeast Portugal: Individual cap and stipe activity. *Food Chemistry*, 100, 1511-1516.

- [34] Deiana, M.; Dessi, M.A.; Bin, K.; Liang, Y.F.; Higa, T.; Gilmour, P.S.; Jen, L.S.; Rahman, I. & Aruoma, O.I. (2002). The antioxidant cocktail effective microorganism (EMX) inhibits oxidant-induced interleukin-8 release and the peroxidation of phospholipids in vitro. *Biochem Biophys Res Commun* 296(5), 1148-1151.
- [35] Sokół-Lętowska, A.; Oszmiański, J. & Wojdyło, A. (2007). Antioxidant activity of the phenolic compounds of hawthorn, pine and skullcap. *Food Chemistry*, 103, 853-859.
- [36] Othman, A., Ismail, A., Ghani, N. A., & Adenan, I. (2007). Antioxidant capacity and phenolics content of cocoa beans. *Food Chemistry*, 100, 1523–1530.
- [37] Tepe, B.; Sokmen, M.; Akpulat, H.A. & Sokmen, A. (2006). Screening of the antioxidant potentials of six *Salvia* species from Turkey. *Food Chemistry*, 95, 200-204.
- [38] Lim, Y.Y.; Lim, T.T. & Tee, J.J. (2007). Antioxidant properties of several tropical fruits: A comparative study. *Food Chemistry*, 103: 1003-1008.
- [39] Teow, C.C.; Truong, V.; Mcfeeters, R.F.; Thompson, R.L.; Pecota, K.V. & Yencho, C.G. (2007). Antioxidant activities, polyphenol and beta-carotene contents of sweetpotato genotypes with varying flesh colors. *Food Chemistry*, 103, 829-838.

- [40] Hertog, M.G.L.; Kromhout, D.; Aravanis, C. (1995). Flavonoid intake and long-term risk of coronary heart disease and cancer in seven countries study. *Arch Intern Med*, 155, 381-386.
- [41] Yang, C.S.; Yang, G.Y.; Landau, J.M.; Kim, S.; Liao, J. (1998). Tea and tea polyphenols inhibit cell hyperproliferation, Lung tumorigenesis, and tumor progression. *Exp Lung Res*, 24, 629-639.
- [42] Bagchi, D.; Sen, C.K.; Ray, S.D.; Das, D.K.; Bagchi, M.; Preuss, H.G. & Vinson, J.A. (2003). Molecular mechanisms of cardioprotection by a novel grape seed proanthocyanidin extract. *Mutat Res*, 523–524, 87–97.
- [43] Aruoma, O.I.; Sun, B.; Fujii, H.; Vidushi, S.; Neergheen, V.S.; Bahorun, T.; Kang, K.S. & Sung, M.K. (2006). Low molecular proanthocyanidin dietary biofactor Oligonol: its modulation of oxidative stress, bioefficacy, neuroprotection, food application and chemoprevention potentials. *Biofactors*, 27, 245-265.
- [44] Yen, G.C. & Chen, H.Y. (1995). Antioxidant activity of various tea extracts in relation to their antimutagenicity. *J Agric Food Chem*, 43, 27–32.
- [45] Liviero, L.; Puglisi, P.P. & Bombardelli, E. (1994). Antimutagenic activity of procyanidins from *Vitis vinifera*. *Fitoterapia*, 65, 203–209.

- [46] Horn, R.; Vondermuhll, M.; Comte, M. & Grandroques, C. (1970). Action de quelques catechins sur l'activit  d'un enzyme (la cytochrome oxidase) de la chaine respiratoire. *Experiential*, 26, 1081–1082.
- [47] Hertog, M.G.L.; Fesken, E.; Hollman, P.; Katan, M. & Kromhout, D. (1993). Dietary antioxidant flavonoids and risk of coronary heart disease: the Zutphen Elderly study. *Lancet*, 342, 1007–1010.
- [48] Teissedre, P.; Frankel, E.N.; Waterhouse, A.L.; Peleg, H. & German, J.B. (1996). Inhibition of in vitro human LDL oxidation by phenolic oxidants from grapes and wines. *J Sci Food Agric*, 70, 55–61.
- [49] Yamakoshi, J.; Saito, M.; Kataoka, S. & Tokutake, S. (2002). Procyanidin-rich extract from grape seeds prevents cataract formation in hereditary cataractous (ICR/f) rats. *J Agric Food Chem*, 50, 4983–4988.
- [50] De Martinis, M.; Franceschi, C.; Monti, D. & Ginaldi, L. (2006). Inflammation markers predicting frailty and mortality in the elderly. *Exp Mol Pathol*, 80, 219–227.
- [51] Yang, B.; Kotani, A.; Arai, K. & Kusu, F. (2001). Estimation of the antioxidant activities of flavonoids from their oxidation potentials. *Anal Sci*, 17, 599–604.
- [52] Salah, N.; Miller, N.G.; Paganga, G.; Tigburg, L.; Bolwell, G.P. & Rice-Evans, C. (1995). Polyphenolic flavonoids as scavengers of aqueous phase radicals and as chain-breaking antioxidants. *Arch Biochem Biophys*, 322, 339–346.

- [53] Xu, R.; Yokoyama, W.H.; Irving, D.; Rein, D.; Walzem, R.L. & German, J.B. (1998). Effect of dietary catechin and vitamin E on aortic fatty streak accumulation in hypercholesterolemic hamsters. *Atherosclerosis*, 137, 29–36.
- [54] Belinky, P.A.; Aviram, M.; Fuhrman, B.; Rosenblat, M. & Vaya, J. (1998). The antioxidative effects of the isoflavan glabridin on endogenous constituents of LDL during its oxidation. *Atherosclerosis*, 137, 49–61.
- [55] Renaud, S. & Lorgeril, M. (1992). Wine, alcohol, platelets and the French paradox for coronary heart disease. *Lancet*, 339, 1523–1526.
- [56] Criqui, M.H. & Ringel, B.L. (1994). Does diet or alcohol explain the French Paradox? *Lancet*, 344, 1719–1723.
- [57] Belleville, J. (2002). The French Paradox: possible involvement of ethanol in the protective effect against cardiovascular disease. *Nutrition*, 18, 173–177.
- [58] Kaur, G.; Roberti, M.; Raul, F. & Pendurthi, U.R. (2007). Suppression of human monocyte tissue factor induction by red wine phenolics and synthetic derivatives of resveratrol. *Thromb Res*, 119, 247–256.
- [59] Zern, T.L. & Fernandez, M.L. (2005). Cardioprotective effects of dietary polyphenols. *J Nutr*, 135, 2291–2294.
- [60] Vedavanam, K.; Srijayanta, S.; O'Reilly, J.; Raman, A. & Wiseman, H. (1999). Antioxidant action and potential antidiabetic properties of an

isoflavonoid-containing soyabean phytochemical extract (SPE).
Phytother Res, 13, 601–608.

[61] Hermansen, K.; Sondergaard, M.; Hoie, L.; Carstensen, M. & Brock, B. (2001). Beneficial effects of a soy-based dietary supplement on lipid levels and cardiovascular risk markers in type 2 diabetic subjects. *Diabetes Care*, 24, 228–233.

[62] Vinson, J.A. & Zhang, J. (2005). Black and green teas equally inhibit diabetic cataracts in a streptozotocin- induced rat model of diabetes. *J Agric Food Chem*, 53, 3710–3713.

[63] Wolfram, S.; Raederstorff, P.; Preller, M.; Wang, Y.; Teixeira, S.R.; Riegger, C. & Weber, P. (2006). Epigallocatechin gallate supplementation alleviates diabetes in rodents. *J Nutr*, 136, 2512–2518.

[64] Rodriguez de Sotillo, D.V. & Hadley, M. (2002). Chlorogenic acid modifies plasma and liver concentrations of: cholesterol, triacylglycerol, and minerals in (*fa/fa*) Zucker rats. *J Nut Biochem*, 13, 717-726.

[65] Balasubashini, M.S.; Rukhumani, R.; Viswanathan, P.; Menon, V.P. (2004). Ferulic acid alleviates lipid peroxidation in diabetic rats. *Phytother Res*, 18, 310–314.

- [66] Jayaprakasam, B.; Vereed, S.K.; Olson, L.K. & Nair, M.G. (2005). Insulin secretion by bioactive anthocyanins and anthocyanidins present in fruits. *J Agric Food Chem*, 53, 28–31.
- [67] Iwai, K.; Onodera, A. & Matsue, H. (2004). Inhibitory effects of viburnum dilatatum Thunb. (gamazumi) on oxidation and hyperglycemia in rats with streptozotocin-induced diabetes. *J Agric Food Chem*, 52(4), 1002-1007.
- [68] Jin, D.Q.; Li, G.; Kim, J.S.; Yong, C.S.; Kim, J.A. & Huh, K. (2004). Preventive effects of Laminaria japonica aqueous extract on the oxidative stress and xanthine oxidase activity in streptozotocin-induced diabetic rat liver. *Biol Pharm Bull*, 27(7), 1037-1040.
- [69] Soon, Y.Y. & Tan, B.K. (2002). Evaluation of the hypoglycemic and anti-oxidant activities of Morinda officinalis in streptozotocin-induced diabetic rats. *Singapore Med J.*, 43(2), 77-85.
- [70] Pari, L. & Latha, M. (2004). Effect of scoparia dulcis (Sweet Broomweed) plant extract on plasma antioxidants in streptozotocin-induced experimental diabetes in male albino Wistar rats. *Pharmazie*, 59(7), 557-560.
- [71] Singh, N.; Kamath, V. & Rajini, P.S. (2005). Attenuation of hyperglycemia and associated biochemical parameters in STZ-induced diabetic rats by dietary supplementation of potato peel powder. *Clin Chim Acta.*, 353(1-2), 165-175.

- [72] Mahesh, T. & Menon, V.P. (2004). Quercetin allievates oxidative stress in streptozotocin-induced diabetic rats. *Phytother Res*, 18(2), 123-127.
- [73] Barthakur, N.N. & Arnold, N.P. (1991). Chemical analysis of the emblic (*Phyllanthus emblica* L.) and its potential as a food source. *Scientia Horticulturae*, 47(1-2), 99-105.
- [74] Anila, L. & Vijayalakshmi, N.R. (2003). Antioxidant action of flavonoids from *Mangifera indica* and *Emblica officinalis* in hypercholesterolemic rats. *Food chemistry*, 83,569-574.
- [75] Liu, X.; Zhao, M.; Wang, J.; Yang, B. & Jiang, Y. (2008). Antioxidant activity of methanolic extract of emblica fruit (*Phyllanthus emblica* L.) from six regions in China. *Journal of Food Composition and Analysis*, 21(3), 219-228.
- [76] Rao, T.P.; Sakaguchi, N; Juneja, Wada, L.R.E & Yokozawa, T. (2005). Amla (*Emblica officinalis* Gaertn.) extracts reduce oxidative stress in streptozotocin-induced diabetic rats. *J Med Food*, 8(3), 362-368.
- [77] Chaiyasut, C & Chansakaow, S. (2007). Inhibitory Effects of Some Thai Plant Extracts on AAPH-induced Protein Oxidation and Protein Glycation. *Naresuan University Journal*, 15(1): 35-41.
- [78] Senthilkumar, G.P. & Subramanian, S.P. (2008). Biochemical studies on the effect of *Terminalia chebula* on the levels of glycoproteins in stz-induced experimental diabetes in rats. *Journal of Applied Biomedicine*, 6(2).105-115.

- [79] Zin, Z.M.; Hamid, A.A.; Osman, A. & Saari, N. (2006). Antioxidative activities of chromatographic fractions obtained from root, fruit and leaf of Mengkudu (*Morinda citrifolia* L.). *Food Chemistry*, 94(2), 169-178.
- [80] Cho, E.J.; Yokozawa, T.; Rhyu, D.Y.; Kim, H.Y.; Shibahara, N. & Park J.C. (2003). The inhibitory effects of 12 medicinal plants and their component compounds on lipid peroxidation. *The American Journal of Chinese Medicine*, 31(6), 907-917.
- [81] Rahul, G.; Kumar, G.B.; Samta, J. & Saxena, A.M. (2008). An overview of indian novel traditional medicinal plants with anti-diabetic potentials. *African Journal of Traditional, Complimentary and Alternative Medicines*, 5(1), 1-17.
- [82] Venkateswaran, S. & Pari, L. (2003). Effect of *Coccinia indica* leaves on antioxidant status in streptozotocin-induced diabetic rats. *J. Ethnopharmacology*, 84, 163-168.
- [83] Platel, K. & Srinivasan, K. (1997). Plant foods in the management of diabetes mellitus: vegetables as potential hypoglycemic agents. *Die. Nahrung*, 41, 68-74.
- [84] Shibib, B.A.; Khan, L.A. & Rahman, R. (1993). Hypoglycemic activity of *Coccinia indica* and *Momordica charantia* in diabetic rat depression of the hepatic gluconeogenic enzymes glucose-6-phosphatase and fructose-1, 6- bisphosphatase and elevation of both liver and red-cell

shunt enzyme glucose-6-phosphate dehydrogenase. *Biochem. J.*, 292, 267-270.

[85] Kamalakkanan, N. & Prince, P.S. (2003). Hypoglycaemic effect of water extract of *Aegle marmelos* fruit in streptozotocin diabetic rats. *J. Ethnopharmacol*, 87, 207-210.

[86] Upadhyaya, S.; Shanbhag, K.K.; Suneetha, G.; Balachandra Naidu, M. & Upadhyaya, S. (2004). A study of hypoglycemic and antioxidant activity of *Aegle marmelos* in alloxan induced diabetic rats. *Ind. J. Physiol. Pharmacol*, 48, 476-480.

[87] Kamalakkanan, N. & Prince, P.S. (2005). The effect of *Aegle marmelos* fruit extract in streptozotocin diabetes: a histopathological study. *J. Herb Pharmacother*, 5, 87-96.

[88] Kesari, A.N.; Gupta, R.K.; Singh, S.K.; Diwakar, S. & Watal, G. (2006). Hypoglycemic and antihyperglycemic activity of *Aegle marmelos* seed extract in normal and diabetic rats. *J. Ethnopharmacology*, 107, 374-379.

[89] Miura, T.; Itoh, Y.; Iwamoto, N.; Kato, M. & Ishida, T. (2004).

Suppressive activity of the fruit of *Momordica charantia* with exercise on blood glucose in type 2 diabetic mice. *Biol. Pharm. Bull.*, 27, 248-250.

- [90] Krawinkel, M.B. & Keding, G.B. (2006). Bitter gourd (*Momordica charantia*): A dietary approach to hyperglycemia. *Nutr. Rev.*, 64, 331-337.
- [91] Chandra, A.; Mahdi, A.A.; Singh, R.K.; Mahdi, F.; Chander, R. (2008). Effect of Indian herbal hypoglycemic agents on antioxidant capacity and trace elements content in diabetic rats. *Journal of Medicinal Food*, 11(3), 506-512.
- [92] Mahdi, A.A.; Chandra, A.; Singh, R.K.; Shukla, L.; Mishra, S.C. & Ahmad, S. (2003). Effect of herbal hypoglycemic agents on oxidative stress and antioxidant status in diabetic rats. *Indian Journal of Clinical Biochemistry*, 18(2), 8-15
- [93] Jelodar, G.A.; Maleki, M.; Motadayen, M.H. & Sirus, S. (2005). Effect of fenugreek, onion and garlic on blood glucose and histopathology of pancreas of alloxan-induced diabetic rats. *Ind. J. Med. Sci.*, 59, 64-69.
- [94] Hattori, A.; Yamada, N.; Nishikawa, T.; Fukuda, H. & Fujino T. (2005). Anti-diabetic effects of ajoene in genetically diabetic KK-A(y) mice. *J. Nutr. Sci. Vitaminol.*, 51, 382-384.
- [95] Ahmad, M.S. & Ahmad N. (2006). Antiglycation properties of aged garlic extract: possible role in prevention of diabetic complications. *J. Nutr.*, 136, 796-799.
- [96] Zhu, Y.P. & Woerdenbag, H.J. (1995). Traditional Chinese herbal medicine. *Pharm. World Sci*, 17, 103.

- [97] Liang, X.C.; Xie, M.Z. & Guo, S.S. (1993). Clinical and experimental study on effect of jin-qi-jiang-tang-pian on qi-yin deficiency and hyperactivity of diabetes mellitus. *Zhongguo Zhong Xi Yi Jie He Za Zhi*, 13(10), 587-90, 579.
- [98] Kim, H.Y.; Yokozawa, T.; Cho, E.J. & Yamabe, N. (2004). Protective effects of the Chinese prescription Hachimi-jio-gan against diabetic oxidative stress. *J. Pharmacy and Pharmacology*, 56, 1299-1305.
- [99] Hu, C.C.; Hsiao, C.H.; Huang, S.Y.; Fu, S.H.; Lai, C.C.; Hong, T.M.; Chen, H.H. & Lu, F.J. (2004). Antioxidant activity of fermented soybean extract. *J Agric Food Chem*, 52(18), 5735-5739.
- [100] Kim, N.Y.; Song, E.J.; Kwon, D.Y.; Kim, H.P. & Heo, M.Y. (2008). Antioxidant and antigenotoxic activities of Korean fermented soybean. *Food Chem Toxicol*, 46(3), 1184-1189.
- [101] Kantachote, D.; Charernjiratrakul, W. & Aussawariangpipop, A. (2005). Characteristics of fermented plant beverages in Southern Thailand. *Songklanakar J. Sci. Technol.*, 27(3), 601-615.
- [102] Kantachote, D.; Charernjiratrakul, W. & Umsakul, K. (2008). Antibacterial activities of fermented plant beverages collected in southern Thailand. *J Biol. Sci.*, 8(8), 1280-1288.
- [103] Barclay, L. (2008). Benefits of probiotics reviewed, *Am Fam Physician*, 78, 1073-1078.

- [104] Minamiyama, Y.; Takemura, S.; Sukioka, T.; Shinkawa, H.; Kobayashi, F.; Nishikawa, Y.; Kodai S.; Mizuguchi, S.; Suehiro, S. & Okada, S. (2007). Effect of AOB, a fermented-grain food supplement, on oxidative stress in type 2 diabetic rats. *BioFactors*, 30(2), 91-104.
- [105] Datla, K.P.; Bennett, R.D.; Zbarsky V.; Ke, B.; Liang Y.F.; Higa, T.; Bahorun, T.; Aruoma, O.I. & Dexter, D.T. (2004). The antioxidant drink "effective microorganism-X (EM-X)" pre-treatment attenuates the loss of nigrostriatal dopaminergic neurons in 6-hydroxydopamine-lesion rat model of Parkinson's disease. *Journal of Pharmacy and Pharmacology*, 56(5), 649-654.
- [106] Ke, B.; Xu, Z.; Ling, Y.; Qiu, W.; Xu, Y., Higa, T. & Aruoma, O.I. (2008). Modulation of experimental osteoporosis in rats by the antioxidant beverage effective microorganism-X (EM-X). *Biomedicine & Pharmacotherapy*, In Press, doi:10.1016/j.biopha.2008.03.008.
- [107] Peerajan, S. (2006). Factors affecting antioxidant capacity and polyphenol content of lactic acid beverages fermented with *Lactobacillus* spp. and *Phyllanthus emblica* Linn. *Master of sciences in pharmaceutical sciences, Graduate school, Chaing Mai University*, p. 114-117.
- [108] Duangjitcharoen, Y.; Kantachote, D.; Ongsakul, M.; Poosaran, N. and Chaiyasut, C. (2008). Selection of probiotic lactic acid bacteria

isolated from fermented plant beverages. *Pak J Biol Sci*, 15, 11(4), 652-655.

[109] Duangjitcharoen, Y.; Kantachote, D.; Ongsakul, M.; Poosaran, N. and Chaiyasut, C. (2009). Potential use of probiotic *Lactobacillus plantarum* SS2 isolated from a fermented plant beverage: safety assessment and persistence in the murine gastrointestinal tract. *World Journal of Microbiology and Biotechnology*, 25, 315-321.

[110] Sirilun, S. (2005). Fermentation kinetics of *Morinda citrifolia* Linn and antimicrobial activity of its products. *Master of sciences in pharmaceutical sciences*, Graduate school, Chaing Mai University, p. 45, 98-111.

[111] Halliwell, B. (1999). Antioxidant defence mechanisms: From the beginning to the end (of the beginning). *Free Radic Res*, 31, 261-272.

[112] Halliwell, B. & Gutteridge J.M.C. (1999). Free radicals in biology and medicine. 3rd Edition, Oxiford: *Clarendon Press*.

[113] Halliwell, B. (1994). Free radicals, antioxidants, and human disease: curiosity, cause, or consequence?. *Lancet*, 344, 721-724.

[114] Halliwell, B. (2001). Role of free radicals in the neurodegenerative diseases: Therapeutic implications for antioxidant treatment. *Drugs and Aging*, 18(9), 685-716.

[115] Cuzzocrea, S.; Riley, D.P.; Caputi, A.P. & Salvemini, D. (2001). Antioxidant therapy: A new pharmacological approach in shock,

inflammation, and ischemia/reperfusion Injury. *Pharmacol Rev*, 53:135-159.

- [116] Ohshima, M.; Li, T.S.; Kubo, M.; Qin, S.L. & Hamano, K. (2009). Antioxidant therapy attenuates diabetes-related impairment of bone Marrow stem cells. *Circ J*, 73, 162-166.
- [117] Cutler, R.G. & Rodrigues, H. (2003). Critical Reviews of Oxidative Stress and Aging: Advances in Basic Science, Diagnostics and Intervention. *Biogerontology*, 4(4), 251.
- [118] Punchard, N.A. & Kelly, F.J. (1996). Free radicals: a practical approach. 1st edition, *Oxford University Press*.
- [119] American Diabetes Association. (2008). Standards of medical care in diabetes. *Diabetes Care*, 31(Suppl), 12-54.
- [120] Expert Committee on the Diagnosis and Classification of Diabetes Mellitus (1997). Report of the Expert Committee on the Diagnosis and Classification of Diabetes Mellitus. *Diabetes Care*, 20, 1183–1197.
- [121] Expert Committee on the Diagnosis and Classification of Diabetes Mellitus (2003). Follow-up report on the diagnosis of diabetes mellitus. *Diabetes Care*, 26, 3160–3167.
- [122] Konrad, R.J.; Mikolaeko, I.; Tolar, J.F.; LIU, K. & Kudlow, J. E. (2001). The potential mechanism of the diabetogenic action of streptozotocin : inhibition of pancreatic b-cell O-GlcNAc-selective N-acetyl-b-D-glucosaminidase. *Biochem. J.*, 356, 31-41.

- [123] Wilson, G.L. & Leiter, E.H. (1990). Streptozotocin interactions with pancreatic beta cells and the induction of insulin-dependent diabetes. In current topics in microbiology and immunology. Edited by Durberg T. Berlin, Germany, *Springerverlag*, 27-54.
- [124] Herold, K.C.; Bloch, T.N.; Vezys, V. & Sun, G. (1995). Diabetes induced with low doses of streptozotocin is mediated by V8.2+T-cells. *Diabetes*, 44, 354-359.
- [125] West, E.; Simon, O.R. & Morrison, E.Y. (1996). Streptozotocin alters pancreatic beta-cell responsiveness to glucose within six hours of injection into rats. *West Ind. Med. J.*, 45, 60-62.
- [126] Bedoya, F.J., Solano, F. & Lucas, M. (1996). Nmonomethyl-arginine and nicotinamide prevent streptozotocin-induced double strand DNA break formation in pancreatic rat islets. *Experientia*, 52, 344-347.
- [127] Schnedl, W.J.; Ferber, S.; Johnson, J.H. & Newgand, C.B. (1994). STZ transport and cytotoxicity. Specific enhancement in GLUT2-expression cells. *Diabetes*, 43, 1326-1333.
- [128] Wang, Z & Gleichmann, H. (1998). GLUT2 in pancreatic islets: crucial target molecule in diabetes induced with multiple low doses of streptozotocin in mice. *Diabetes*, 47(1), 50-56.
- [129] Yamamoto, H.; Uchigata, Y. & Okamoto, H. (1981). DNA strand breaks in pancreatic islets by *in vivo* administration of alloxan or streptozotocin. *Biochem Biophys Res Commun*, 103, 1014-1020.

- [130] Elsner, M.; Guldbakke, B.; Tiedge, M.; Munday, R. & Lenzen, S. (2000). Relative important of transport and alkylation for pancreatic beta-cell toxicity of streptozotocin. *Diabetologia*, 43, 1528-1533.
- [131] Szkudelski, T. (2001). The mechanism of alloxan and streptozotocin action in B cells of the rat pancreas. *Physiol. Res.*, 50, 536-546.
- [132] Nukatsuka, M. & Yoshimura, Y. & Nishida, M. (1990). Importance of the concentration of ATP in rat pancreatic beta cells in the mechanism of streptozotocin-induced cytotoxicity. *J Endocrinol* , 127, 161-165.
- [133] Kröncke, K.D.; Sommer, A.; Rodriguez, ML. & Kolb-Bachofen, V. (1995). Nitric oxide generation during cellular metabolization of the diabetogenic N-methyl-N-nitroso-urea streptozotocin contributes to islet cell DNA damage. *Biol Chem Hoppe-Seyler*, 376, 2357-2363.
- [134] Turk, J.; Corbett, J.A.; Ramanadham, S.; Bohrer, A. & McDaniel, M.L. (1993). Biochemical evidence for nitric oxide formation from streptozotocin in isolated pancreatic islets. *Biochem Biophys Res Commun*, 197, 1458-1464.
- [135] Heller, B.; Burkle, A.; Radons, J.; Fengler, E.; Jalowoy, A.; Muller, M.; Burkart, V. & Kolb, H. (1994). Analysis of oxygen radical toxicity in pancreatic islets at the single cell level. *Biol. Chem Hoppe-Seyler*, 375, 597-602.

- [136] Ganda, O.P.; Rossini, A.A. & Like AA. (1976). Studies on streptozotocin diabetes. *Diabetes*, 25, 595-603.
- [137] Katsumata, K.; Katsumata, K.J. & Katsumata, Y. (1992). Protective effect of diltiazem hydrochloride on the occurrence of alloxan or streptozotocin-induced diabetes in rats. *Horm Metab Res*; 24: 508-510.
- [138] Blondel, O.; Bailbe, D. & Portha, B. (1989). In vivo insulin resistance in streptozotocin-diabetic rats. Evidence for reversal following oral vanadate treatment. *Diabetologia*, 32, 185-190.
- [139] Cowell, R.C. & Russell, J.W. (2004). Neurosaative injury and antioxidant therapy in the management of diabetic neuropathy. *J Invest Med*, 52, 33-44.
- [140] Ceriello, A. (1999). Hyperglycaemia: the bridge between non-enzymatic glycation and oxidative stress in the pathogenesis of diabetic complications. *Diabetes Nutr Metab*, 12, 42-46.
- [141] Ceriello, A. & Motz, E. (2004). Is Oxidative Stress the Pathogenic Mechanism Underlying Insulin Resistance, Diabetes, and Cardiovascular Disease? The Common Soil Hypothesis Revisited. *Arteriosclerosis, Thrombosis, and Vascular Biology*, 24, 816-823.
- [142] Phillips, M.; Cataneo, R.N.; Cheema, T. & Greenberg, J. (2004). Increased breath biomarkers of oxidative stress in diabetes mellitus. *Clin Chim Acta*, 344, 189-194.

[143] Maddux, B.A.; See, W.; Lawrence, J.C.; Goldfine A,L.; Goldfine, I. & Evans, J. (2001). Protection against oxidative stress-induced insulin resistance in rat L6 muscle cells by micromolar concentrations of α -lipoic acid. *Diabetes*, 50, 404–410.

[144] Kim, K.T.; Yoo, K.M.; Lee, J.W.; Eom, S.H. & Hwang, I.K. (2007). Protective effect of steamed American ginseng (*Panax quinquefolius* L.) on V79-4 cells induced by oxidative stress. *Journal of Ethnopharmacology*, 111(3), 443- 450.

[145] Di Leo, M.A.; Ghirlanda, G.; Gentiloni S.N.; Giardina, B.; Franconi, F. & Santini, S.A. (2003). Potential therapeutic effect of antioxidants in experimental retina: a comparison between taurine and vitamin E plus selenium supplementation. *Free Radic Res*, 37, 323-330.

[146] Jasprica, I.; Mornar, A.; Debeljak, Z.; Smolčić-Bubalo, A.; Medić-Sarić, M.; Mayer, L.; Romić, Z.; Bućan K.; Balog, T.; Sobočanec S.; & Šverko, V. (2007). *In vivo* study of propolis supplementation effects on antioxidative status and red blood cells. *Journal of Ethnopharmacology*, 110, 548–554.

[147] Corbett, J.A.; Tilton, R.G.; Chang, K.; Hasan, K.S.; Ido, Y.; Wang, J.L.; Sweetland, M.A.; Lancaster, J.R.; Williamson, J.R. & Mc Daniel, M.L. (1992). Aminoguanidine, a novel inhibitor of nitric oxide formation, prevents diabetic vascular dysfunction. *Diabetes*, 41, 552–556.

- [148] Re, R.; Pellegrini, N.; Proteggente, A.; Pannala, A.; Yang, M. & Rice-Evans, C. (1999). Antioxidant activity applying an improved ABTS radical cation decolorization assay. *Free Radic Biol Med*, 26, 1231-1237.
- [149] Chirico, S. (1994). High-performance liquid chromatography-based thiobarbituric acid tests. *Methods Enzymol*, 233, 314-318.
- [150] Vinson, J.A. & Howard, T.B. (1996). Inhibition of protein glycation and advanced glycation end products by ascorbic acid and other vitamins and nutrients. *The Journal of Nutritional Biochemistry*, 7, 659-663.
- [151] McDonald, S.; Prenzler, P.D.; Antolovich, M. & Robards, K. (2001). Phenolic content and antioxidant activity of olive extracts. *Food Chemistry*, 73, 73-84.
- [152] Chang, C.C.; Yang, M.H.; Wen, H.M. & Chen, J.C. (2002). Estimation of total flavonoid content in propolis by two complementary colorimetric methods. *J Food Drug Analysis*, 10, 178-182.
- [153] Polshettiwar, S.A.; Ganjiwale, R.O.; Wadher, S.J. & Yeole, P.G. (2007). Spectrophotometric estimation of total tannins in some Ayurvedic Eye Drops. *Indian Journal of Pharmaceutical Science*, 69, 574-576
- [154] Chiou, A.; Karathanos, V.T.; Mylona, A.; Salta, F.N.; Preventi, F. and Andrikopoulos, N.K. (2007). Currants (*Vitis vinifera* L.) content of simple phenolics and antioxidant activity. *Food Chemistry*, 102(2), 516-522.

- [155] Mabinya, L.V., Mafunga, T. & Brand, J.M. (2006). Determination of ferulic acid and related compounds by thin layer chromatography. *African Journal of Biotechnology*, 5(13), 1271-1273.
- [156] Trinder, P. (1969). Determination of glucose in blood using glucose oxidase with an alternative oxygen acceptor. *Ann Clin Biochem*, 6, 24-27.
- [157] Amer, J.; Goldfarb, A. & Fibach, E. (2003) Flow cytometric measurement of reactive oxygen species production by normal and thalassaemic red blood cells. *European Journal Of Haematology*, 70, 84-90.
- [158] Ewing, J.F. & Janero, D.R. (1995). Microplate superoxide dismutase assay employing a nonenzymatic superoxide generator. *Anal. Biochem*, 232, 243-248.
- [159] Misko, T.P.; Schilling, R.J.; Salvemini, D.; Moore, W.M. & Currie, M.G. (1993). A fluorometric assay for the measurement of nitrite in biological samples. *Anal Biochem*, 214, 11-16.
- [160] Rice-Evans, C.A.; Miller, N.J. & Paganga, G. (1996). Structure-Antioxidant activity. relationships of flavonoids and phenolic acids. *Free Radical Biology and Medicine*, 20(7), 933-956.
- [161] Llaudy, M.C.; Canals, R.; Canals, J.M.; Rozes, N.; Arola, L. & Zamora, F. (2004). New method for evaluating astringency in red wine. *Journal of Agricultural and Food Chemistry*, 52, 742-746.

- [162] Scott, J.A. & King, G.L. (2006). Oxidative Stress and Antioxidant Treatment in Diabetes. *Annals of the New York Academy of Sciences*, 1031, 204-213.
- [163] Jakus, V. & Rietbrock, N. (2004). Advanced glycation end-products and the progress of diabetic vascular complications. *Physiological Research*, 53, 131-142.
- [164] Thornalley, P.J. (2002). Glycation in diabetic neuropathy: characteristics, consequences, causes and therapeutic options. *Int Rev Neurobiol*, 50, 37-57.
- [165] Kessler, M., Ubeaud, G. & Jung, L. (2003). Anti-and pro-oxidant activity of rutin and quercetin derivatives. *J. Pharm and Pharmacol*, 55, 131-142.
- [166] Singh, R.; Singh, S.; Kumar, S. & Arora, S. (2007). Studies on antioxidant potential of methanol extract/fractions of *Acacia auriculiformis* A. Cunn. *Food Chemistry*, 103, 505-511.
- [167] Ramadan, M.F.; Zayed, R. & El-Shamy, H. (2007). Screening of bioactive lipids and radical scavenging potential of some solanaceae plants. *Food Chemistry*, 103, 885-890.
- [168] Ronald, A.A. (1982). Clinical correlates of metabolic derangement of diabetes. *Clinical diabetes mellitus*. W.B. Saunders Co., U.S.A., 16-20.

- [169] Hu, C.C.; Hsiao, C.H.; Huang, S.Y.; Fu, S.H.; Lai, C.C.; Hong, T.M.; Chen, H.H. & Lu, F.J. (2004). Antioxidant activity of fermented soybean extract. *J Agric Food Chem*, 8, 52(18), 5735-5759.
- [170] Kim, N.Y.; Song, E.J.; Kwon, D.Y.; Kim, H.P. & Heo, M.Y. (2008). Antioxidant and antigenotoxic activities of Korean fermented soybean. *Food Chem Toxicol*, 46(3), 1184-1189.
- [171] Unger, R.H. (1978). Role of glucagon in the pathogenesis of diabetes: The status of controversy. *Metabolism*, 27, 1671-1709.
- [172] Felliet-Coudray, C.; Rock, E.; Coudray, C.; Grzelkowska, K.; Azais-Braesco, V. & Dardevet, D. (1999). Lipid peroxidation and antioxidant status in experimental diabetes. *Clin Chim Acta*, 284:31-43.
- [173] Segrouchni, I.; Draï, J.; Bannier, E.; Riviere, J.; Calmand, P.; Garcia, I. et al. (2002). Oxidative stress parameters in type 1, type 2 and insulin-treated type 2 diabetes mellitus: insulin treatment efficiency. *Clin Chim Acta*, 321, 89–96.
- [174] Kakkar, R.; Kalra, J.; Mantha, S.V. & Prasad, K. (1995). Lipid peroxidation and activity of antioxidant enzymes in diabetic rats. *Mol. cell Biochem*, 151, 113-119.
- [175] Yazdanparast, R.; Ardestani, A. & Jamshidi, S. (2007). Experimental diabetes treated with *Achillea santolina*: Effect on pancreatic oxidative parameters. *J Ethnopharmacol*, 112(1), 13-18.

- [176] Chui, C.H.; Cheng, G.Y.M.; Ke, B.; Lau, F.Y. & Wong, R.S.M. (2004). Growth inhibitory potential of effective microorganism fermentation extract (EM-X) on cancer cells. *International Journal of molecular medicine*, 14, 925-929.
- [177] Amer, J. & Fibach, E. (2004). Oxidative status of platelets in normal and thalassemia blood. *Thromb Haemost*, 92, 1052-1059.
- [178] Kumazawa, S.; Hamasaka, T. & Nakayama, T. (2004). Antioxidant activity of propolis of various geographic origins. *Food Chemistry*, 84, 329-339.
- [179] Banskota, A.H.; Tezuka, Y. & Kadota, Sh. (2001). Recent progress in pharmacological research of propolis. *Phytotherapy Research*, 15, 561-571.
- [180] Isla, M.I.; Moreno, M.I.N.; Sampietro, A.R. & Vattuone, M.A. (2001). Antioxidant activity of Argentine propolis extracts. *Journal of Ethnopharmacology*, 76, 165-170.
- [181] Kolankaya, D.; Selmanoğlu, G.; Sorkun, K. & Salih, B. (2002). Protective effects of Turkish propolis on alcohol-induced serum lipid changes and liver injury in men rats. *Food Chemistry*, 78, 213-217.
- [182] Russo, A., Cardile, V., Sanchez, F.; Troncoso, N.; Vanella, A. & Garabino, J.A. (2004). Chilean propolis: antioxidant activity and antiproliferative action in human tumor cell lines. *Life Sciences*, 76, 545-558.

- [183] Williams, R.J.; Spencer, J.P.E. & Rice-Evans, C.A. (2004). Flavonoids: antioxidants or signaling molecules? *Free Radical Biology and Medicine*, 36, 838–849.
- [184] Middleton, E.; Kandaswamy, Ch. & Theoharides, T.C. (2000). The effects of plant flavonoids on mammalian cells: implications for inflammation, heart disease and cancer. *Pharmacological Reviews*, 52, 673–751.
- [185] Beckman, J.S.; Beckman, T.W.; Chen, J.; Marshall, P.A. & Freeman, B.A. (1990). Apparent hydroxyl radical production by peroxynitrite: implications for endothelial injury from nitric oxide and superoxide. USA. *Proc.Natl Acad.Sci.*, 87, 1620-1624.
- [186] Tesfamariam, B. (1994). Free radicals in diabetic endothelial cell dysfunction. *Free radic. Biol. Med*, 16, 383-391.
- [187] McCune, L.M. & Johns, T. (2002). Antioxidant activity in medicinal plants associated with the symptoms of diabetes mellitus used by the indigenous peoples of the North American boreal forest. *Journal of Ethnopharmacology*, 82, 197-205.
- [188] Kim, K.T.; Yoo, K.M.; Lee, J.W.; Eom, S.H.; Hwang, I.K. & Lee, C.Y. (2007). Protective effect of steamed American ginseng (*Panax quinquefolius* L.) on V79-4 cells induced by oxidative stress. *Journal of Ethnopharmacology*, 111(3), 443- 450.

- [189] Cao, G.; Sofic, E. & Prior, R.L. (1997). Antioxidant and prooxidant behavior of flavonoids: structure-activity relationships. *Free Radical Biology and Medicine*, 22, 749-760.
- [190] Galati, G. & O'Brien, P.J. (2004). Potential toxicity of flavonoids and other dietary phenolics: significance for their chemopreventive and anticancer properties. *Free Radical Biology and Medicine*, 37, 287-303.
- [191] Landmesser, U. & Drexler, H. (2005). The clinical significance of endothelial dysfunction. *Current Opinion in Cardiology*, 20, 547-551.
- [192] Kone, B.C.; Kunczewicz, T.; Zhang, W. & Yu, Z.Y., (2003). Protein interactions with nitric oxide synthases: controlling the right time, the right place, and the right amount of nitric oxide. *American Journal of Physiology Renal Physiology*, 285, F178-F190.