

REFERENCES

- Abdalla Mäder. (2001). Preparation and characterization of a self-emulsifying pellet formulation. *European Journal of Pharmaceutics and Bio-pharmaceutics.*, 66 (2), 220-226.
- Aguiar, A.J., Krc. Jr, J., Kinkel, A. W. and Samyn, J. C. (1967). Effect of polymorphism on the absorption of chloramphenicol from chloramphenicol palmitate. *J. Pharm. Sci.*, 56, 847-853.
- Alway Beverley, Sangchantra Roongtawan and Stewart Peter J. (1996). Modelling the dissolution of diazepam in lactose interactive mixtures. *Int. J. Pharm.*, 130 (2), 213-224.
- Ametch Industry CO., Limited. 2009. "Introduce About Dicalcium Phosphate." [Online]. Available [http:// www.amtechem.com/DicalciumPhosphate.html](http://www.amtechem.com/DicalciumPhosphate.html) (10 February 2009).
- Ando Howard Y and Radebaugh Galen W, (2005). Property-Based Drug Design and Preformulation. in David Troy (Ed.), *Remington: The Science and Practice of Pharmacy* 21st Edition (pp. 738), Philadelphia : Lippincott Williams & Wilkins.
- Armstrong, Norman Anthony (2007). Tablet Manufacture. in J. Swarbrick (Ed.), *Encyclopedia of Pharmaceutical Technology* Third Edition Vol. 6 (pp. 3649, 3653, 3655, 3663, 3656, 3669), New York : Informa Healthcare USA, Inc.
- Augsburger, L.L., Zellhofer, M.J. (2007a). Tablet Formulation. in J. Swarbrick (Ed.), *Encyclopedia of Pharmaceutical Technology* Third Edition Vol. 6 (pp. 3644-49), New York : Informa Healthcare USA, Inc.
- Augsburger L.L., Brzezczko A.W., Shah, U., Hahm H.A. (2007b) Super Disintegrants: Characterization and Function. in James Swarbrick (Ed.), *Encyclopedia of PHARMACEUTICAL TECHNOLOGY* Third Edition Vol. 6 (pp. 3553), New York : Informa Healthcare USA, Inc.
- Aulton, M.E. (2002). Bioavailability-physicochemical and dosage form factors. in *Pharmaceutics The Science of Dosage Form Design* Second Edition (pp. 239), UK : Churchill Livingstone.

- Bandelin, Fred J. (1989). Compressed Tablet by Wet Granulation. in H.A. Lieberman, L. Lachman, and J.B. Schwartz (Eds.) *Pharmaceutical Dosage Forms: Tablets* 2nd edition vol 1 (pp. 158). New York : Marcel Dekker.
- Barra, J, Somma, R. (1996). Influence of the physicochemical variability of magnesium stearate on its lubricant properties: possible solutions. *Drug Dev. Ind. Pharm.*, 22(11), 1105–1120.
- Bavitz, J.F. and Schwartz, J.B. (1976). Direct compression vehicles: effect of common diluents on compression. *Drug Cosm. Ind.*, 118(4), 60-64.
- Becker, D, Rigassi T, Bauer-Brandl A. (1997). Effectiveness of binders in wet granulation: comparison using model formulations of different tableability. *Drug Dev. Ind. Pharm.*, 23(8), 791–808.
- Billany MR. and Richards JH. (1982). Batch variation of magnesium stearate and its effect on the dissolution rate of salicylic acid from solid dosage forms. *Drug Dev. Ind. Pharm.*, 8, 497–511.
- Bolhuis, G.K. and Holzer, A.W. (1996). Lubricant sensitivity. in Alderborn, G., Nystrom, C. (Eds.), *Pharmaceutical Powder Compaction Technology* (pp. 517-560), New York : Marcel Dekker, Inc.
- Bolhuis G.K., Lerk C.F., Broersma P. (1980). Mixing action and evaluation of tablet lubricants in direct compression. *Drug Dev. Ind. Pharm.*, 6, 573–589.
- Bootsma, H.P.R., Frijlink, H.W., Eissens, A., Proost, J.H., Van Doorne, H. Lerk, C.F. (1989). β -Cyclodextrin as an excipient in solid oral dosage forms: in vitro and in vivo evaluation of spray-dried diazepam- β -cyclodextrin products. *Int. J. Pharm.*, 51 (3), 213-223.
- Bossert, J., Stamm, A. (1980). Effect of mixing on the lubrication of crystalline lactose by magnesium stearate. *Drug Dev. Ind. Pharm.*, 6, 573–589.
- British Pharmacopoeia 2007 London : HMSO, : pp. 629, Appendix XII D, XII H. A (pp.108, 112 of 184). *Ph. Eur.* 5.5
- Brittain Harry G. (2007). Polymorphism: Pharmaceutical Aspects. in James Swarbrick (Ed.), *Encyclopedia of Pharmaceutical Technology* Third Edition VOLUME 6 (pp. 2940, 2942), New York : Informa Healthcare USA, Inc.

- Buehler, John D. "Medicinal simethicone containing composition and its method of production" U.S. Pat 4127650 Nov. 28, 1978.
- Cal, S., Iglesias, G., Souto, C., Concheiro, A., Gómez-Amoza, J. L. and Martínez-Pacheco, R. (1996). Effects of hydration on the properties of a roller-dried β -lactose for direct compression. *Int. J. Pharm.*, 129 (1-2), 253-261.
- Caramella, C., Columbo, P., Conte, U., Ferrari, F., Manna, A. La, Kamp, H.V. Van, and Bolhuis, G.K. (1986). Water Uptake and Disintegrating Force Measurements: Towards a General Understanding of Disintegration Mechanisms, *Drug Dev. Ind. Pharm.*, 12 (11 – 13), 1749 – 1766.
- Carstensen, J.T., Ertell, C. (1990). Physical and chemical properties of calcium phosphate for solid state pharmaceutical formulations. *Drug Dev. Ind. Pharm.*, 16 (7), 1121–1133.
- Celebi, N. (1989). *In vitro* investigation of the absorption of indomethacin from its β -cyclodextrin complex with competing agents. *Pharmazie*. 44, 133-135.
- Chan, H. K., Doelker, E. (1985). Polymorphic Transformation of Some Drugs Under Compression. *Drug Dev. Ind. Pharm.*, 11 (2-3), 315-332.
- Cheng, W.T., Lin, S.Y., & Li, M.J. (2007). Raman microspectroscopic mapping or thermal system used to investigate milling-induced solid-state conversion of famotidine polymorphs. *J. of Raman Spectroscopy.*, 38(12), 1595 - 1601.
- Choudhari, K.B., Sanghavi, N.M. (1993). Dissolution behaviour and characterization of diazepam-Pullulan coground mixtures. *Int. J. Pharm.*, 89 (3), 207-211.
- Choui, W. L., Reigelman, S. (1969). Preparation and dissolution characteristics of several fast release solid dispersions of griseofulvin. *J. Pharm. Sci.* 58, 1505-1510.
- Choui, W L, Chen, S J, Athanikar, N. (1976). Enhancement of poorly water soluble drugs by crystallisation in aqueous surfactant solutions. *J. Pharm. Sci.*, 65, 1702-1704.
- Chowhan, ZT, Chi, LH. (1986a). Drug–excipient interactions resulting from powder mixing III: solid state properties and their effect on drug dissolution. *J. Pharm. Sci.*, 75, 534–541.

- Chowhan ZT, Chi LH. (1986b). Drug–excipient interactions resulting from powder mixing IV: role of lubricants and their effect on *in vitro* dissolution. *J. Pharm. Sci.*, 75, 542–545.
- Cunha-Filho Marcílio, S.S., Martínez-Pacheco Ramón and Landín Mariana. (2008). Dissolution rate enhancement of the novel antitumoral β -lapachone by solvent change precipitation of microparticles. *European Journal of Pharmaceutics and Biopharmaceutics.*, 69 (3), 871-877.
- Cwiertnia, B. (2006). Solubility of selected derivatives of 1,4-benzodiazepin-2-one in solid dispersions in PEG 6000. *Acta Pol Pharm.*, 63(2), 135-139.
- DeCamp W.H. (1999). “THE IMPACT OF POLYMORPHISM ON DRUG DEVELOPMENT: A REGULATOR'S VIEWPOINT”, Presented to *XVIIIth Congress of the International Union of Crystallography* Glasgow, Scotland.
- Desai DS, Rubitski BA, Varia SA, Newman AW. (1993). Physical interactions of magnesium stearate with starch-derived disintegrants and their effects on capsule and tablet dissolution. *Int. J. Pharm.*, 91(2–3), 217–226.
- Doherty, C. York, P. (1989a). Microenvironmental pH control of drug dissolution. *Int. J. Pharm.*, 50, 223-232.
- Doherty, C. York, P. (1989b). The *in vitro* pH dissolution dependence and *in vivo* bioavailability of frusemide-PVP solid dispersions. *J. Pharm. Pharmacol.*, 41, 74-78.
- Drooge, D. J. van, Hinrichs, W. L. J. and Frijlink, H. W. (2004). Anomalous dissolution behaviour of tablets prepared from sugar glass-based solid dispersions. *Journal of Controlled Release.*, 97 (3), 441-452.
- Eckhart, C.G., McCorkle, T. (1978). Chlorpheniramine maleate. in Klaus Florey (Ed.), *Analytical Profiles of Drug Substances*, vol. 7 (pp 43-80), New York : Academic Press, Inc.
- Elshaboury, M.H. (1990). Physical properties and dissolution profiles of tablets directly compressed with β -cyclodextrin. *Int. J. Pharm.*, 63 (2), 95-100.
- Elzbieta, F., Krystyna (Dartford, GB2). “Water-dispersible tablets” U.S. Pat 5660860 Aug. 26, 1997
- Ertel, K.D., Carstensen, J.T. (1988). An examination of the physical properties of pure magnesium stearate. *Int. J. Pharm.*, 42, 171–180.

- Ertel, K.D., Zoglio, M.A., Ritschel, W.A., Carstensen, J.T. (1990). Physical aspects of wet granulation. IV. Effect of kneading time on dissolution rate and tablet properties. *Drug Dev. Ind. Pharm.*, 16 (6), 963–981.
- Fukunaka Tadashi, Yaegashi Yoshiko, Nunoko Taku, Ito Ryusei, Golman Boris, Shinohara Kunio. (2006). Dissolution characteristics of cylindrical particles and tablets. *Int. J. Pharm.*, 310, 146–153.
- Gander, B., Ventouras, K., Gurny, R., Doelker, E. (1985). *In vitro* dissolution medium with supra-micellar surfactant concentration and its relevance for *in vivo* absorption. *Int. J. Pharm.*, 27, 117-124.
- Ganderton D. (1969). The effect of distribution of magnesium stearate on the penetration of a tablet by water. *J. Pharm. Pharmacol.*, 21 (Suppl.), 9S–18S.
- Gibaldi M. (1984). *Biopharmaceutics and clinical pharmacokinetics*. Philadelphia : Lea and Febiger.
- Goldberg, A. H., Gibaldi, M., Kanig, J. L., Mayersohn, M. (1966). Increasing dissolution rates and gastrointestinal absorption of drugs via solid solutions and eutectics. *J. Pharm. Sci.*, 55, 581-583.
- Gole, D. J., Levinson, R., Saul Wilkinson, Paul, K., Davies J. Desmond. “Pharmaceutical and other dosage forms” U.S. Pat 5558880 Sep. 24, 1996
- Gordon, M.S., Chowhan, Z.T. (1990). The effect of aging on disintegrant efficiency in direct compression tablets with varied solubility and hygroscopicity, in terms of dissolution. *Drug Dev. Ind. Pharm.*, 16, 437–447.
- Gordon, M.S., Rudraraju, VS., Dani, K., Chowhan, ZT. (1993). Effect of the mode of super disintegrant incorporation on dissolution in wet granulated tablets. *J. Pharm. Sci.*, 82(2), 220–226.
- Gordon, R.E., Rosanske, T.W., Fonner D.E., Anderson, N.R. and Banker G.S. (1989). Granulation Technology and Tablet Characterization. in H.A. Lieberman, L. Lachman, Schwartz, J.B. (Eds.) *Pharmaceutical Dosage Forms: Tablets* 2nd edition vol 2 (pp. 325-333). New York : Marcel Dekker.
- Haleblian, J. K. (1975). Characterisation of habits and crystalline modification of solids and their pharmaceutical applications. *J. Pharm. Sci.*, 64, 1269-1288.
- Heckel, R.W. (1961). Density-pressure relationships in powder compaction. *Trans. Metall. Soc. AIME*, 221, 671–675.

- Hecq, J., Deleers, M., Fanara, D., Vranckx, H. and Amighi, K. (2005). Preparation and characterization of nanocrystals for solubility and dissolution rate enhancement of nifedipine. *Int. J. Pharm.*, 299 (1-2), 167-177.
- Heil Wolfgang, Tistam Ulf, Lipp Ralph, Tack Johannes-Wilhelm. "Oral fludara of high-purity formulation with quick release of active ingredient" U.S. Pat 7148207 Dec. 12, 2006.
- Hoener, B., Benet, L.Z. (2002). Factor influencing Drug Absorption and Availability. in G.S. Banker, C.T. Rhodes (Eds.), *Modern pharmaceuticals* 4th ed. Revised and Expanded (pp. 99, 105), New York : Marcel Dekker, Inc.
- Holm Per, Holm Jannie Egeskov, Ruhland Thomas, Nielsen Simon Dalsgaard. "Porous tablets as carriers for liquid formulations" Pub. No. WO/2006/000229 Jan. 05, 2006.
- Imai, T., Nishiyama, T., Shameem, M., Otagiri, M. (1998). Casein hydrolysate as a rapid and/or enteric dissolving additive for oral drugs. *Pharm. Dev. Technol.*, 3 (2), 225-232.
- Iwata, M., Ueda, H., (1996). Dissolution properties of glibenclamide in combinations with polyvinylpyrrolidone. *Drug Dev. Ind. Pharm.*, 22, 1161-1165.
- Jachowitz, R. (1987). Dissolution of partially soluble drugs from solid dispersion systems. *Int J Pharm.*, 35, 1-12.
- Jackson, K., Young, D., and Pant, S. (2000). Drug- excipient interactions and their affect on absorption. *Pharmaceutical Science & Technology Today*, 3 (10), 336 – 345 .
- Javadsadeh, Y., Siahi-Shadbad, M.R., Barzegar-Jalali, M., Nokhodchi, A. (2005). Enhancement of dissolution rate of piroxicam using liquisolid compacts. *IL Farmaco.*, 60, 361-365.
- Javadsadeh, Y., Jafari-Navimipour, B., Nokhodchi A. (2007). Liquisolid technique for dissolution rate enhancement of a high dose water-insoluble drug (carbamazepine). *Int. J. Pharm.*, 341, 26-34.
- Johnson, JR., Wang, L-H., Gordon, MS., Chowhan, ZT. (1991). Effect of formulation solubility and hygroscopicity on disintegrant efficiency in tablets prepared by wet granulation, in terms of dissolution. *J. Pharm. Sci.* , 80, 469-471.

- Kaneniwa, N. and Otsuka, M. (1985). Effect of grinding on the transformation of polymorphs of chloramphenicol palmitate. *Chem Pharm Bull.*, 33, 1660-1668.
- Kaneniwa, N., Imagawa, K., Otsuka, M. (1985). Effect of tableting on the degree of crystallinity and the dehydration and decomposition points of cephalexin crystalline powder. *Chem Pharm Bull.*, 33, 802-809.
- Kawakita, K. and Ludde, K.H. (1970). Some consideration on powder compression equations. *Powder Technol.*, 4, 61-68.
- Khan, K.A., Rhodes, C.T. (1975). Disintegration properties of calcium phosphate dibasic dihydrate tablets. *J. Pharm. Sci.*, 64, 166-168.
- Khatab, I., Menon, A., Sakr, A. (1993). Effect of mode of incorporation of disintegrants on the characteristics of fluid-bed wet-granulated tablets. *J. Pharm. Pharmacol.*, 45(8), 687-691.
- Kim Min-Soo, Lee Sibeum, Park Jeong-Sook, Woo Jong-Soo and Hwang Sung-Joo. (2007). Micronization of cilostazol using supercritical antisolvent (SAS) process: Effect of process parameters. *Powder Technology.*, 177 (2), 64-70.
- Kim Min-Soo, Jin Shun-Ji, Kim Jeong-Soo, Park Hee Jun, Song Ha-Seung, Neubert Reinhard H.H. and Hwang Sung-Joo. (2008). Preparation, characterization and *in vivo* evaluation of amorphous atorvastatin calcium nanoparticles using supercritical antisolvent (SAS) process. *European Journal of Pharmaceutics and Biopharmaceutics.*, 69 (2), 454-465.
- Kinget, R. (1989). Comparison of the dissolution behaviour of tablets containing the drug in a polyethyleneglycol solid dispersion with direct compressed tablets. *Pharm. Acta Helv.*, 64(3), 90-93.
- Koivisto, M., Heinänen, P., Tanninen, V.P., Lehto, V.P. (2006). Depth profiling of compression-induced disorders and polymorphic transition on tablet Surfaces with grazing incidence X-ray diffraction. *J. of Raman Spectroscopy.*, 23 (4), 813-820.
- Lee, Fang-Yu; Lee, Fang-Chen "Chargeable pharmaceutical tablets" U.S. Pat 6399591 Jun. 4, 2002.
- Lee, Fang-Yu; Chen Shan-chiung, Kuo Han-Chiang. "Orally administered pharmaceutical formulations of benzimidazole derivatives and the method of preparing the same" U.S. Pat 6228400 May 8, 2001.

- Lee Beom – Jin (2008). 6.1 Pharmaceutical Preformulation: Physicochemical Properties of excipients and Powders and Tablet Characterization. in Shayne Cox Gad (Ed.), *Pharmaceutical Manufacturing Handbook: Production and Processes*. pp. 896 – 929. New Jersey : John Wiley & Sons, Inc.
- Leinonen, U.I., Jalonen, H.U., Vihervaara, P.A, Laine, E.S.U. (1992). Physical and lubrication properties of magnesium stearate. *J. Pharm. Sci.*, 81(12), 1194–1198.
- Lesson, L.T., Carstensen, J.T. (1974). In *Dissolution Technology*, Washington, D.C : Academy of Pharmaceutical Science.
- Leuner Christian and Dressman Jennifer. (2000). Review article. Improving drug solubility for oral delivery using solid dispersions. *European Journal of Pharmaceutics and Biopharmaceutics.*, 50 (1, 3), 47-60.
- Levy, G., Gumtow, R.H. (1963). Effect of certain formulation factors on dissolution rate of the active ingredient III: tablet lubricants. *J. Pharm. Sci.*, 52, 1139–1144.
- Lewis Gareth A. (2007). Optimization Methods. in James Swarbrick (Ed.), *Encyclopedia of Pharmaceutical Technology* Third Edition Vol. 4 (pp. 2454), New York : Informa Healthcare USA, Inc.
- Lin, S.Y., Yang, J.C. (1986). Inclusion complexation of warfarins with cyclodextrins to improve some pharmaceutical characteristics. *Pharm. Weekblad Sci.*, 8, 223-228.
- Lu, W.G., Zhang, Y., Xiong, Q.M. (1995). Development of nifedipine (NE) pellets with a high bioavailability. *Chin. Pharm. J. Zhongguo Yaoxue Zazhi.*, 30 (Nov Suppl), 24–26.
- Macdona, A., Michaelis, A.F., Senkowski, B.Z. (1972). Diazepam. in Klaus Florey (Ed.), *Analytical Profiles of Drug Substances* Vol.1 (pp.79-99), New York : Academic Press, Inc.
- Melia, C.D. and Davis S.S. (1989). Review article: mechanisms of drug release from tablets and capsules. 2. Dissolution, Strategies for enhancing drug dissolution rates from tablets and capsules. *Aliment. Pharmacol. Therap.*, 3, 513-525.
- Mendell, E.J. (1972). Direct Compression Method of Producing Solid Dosage Forms. *Mfg. Chemist Aerosol News.*, 43, 43-46.

- Miyazaki, S., Artist T., Hori R., Ito K. (1974). Effect of polymorphism on the dissolution behavior and gastrointestinal absorption of chlortetracycline hydrochloride. *Chem. Pharm. Bull.*, 22 : 638-642.
- Muzikova, J. (2002). Effect of magnesium stearate on the tensile strength of tablets made with the binder *Prosolv SMCC 90*. *Ceska Slov Farm*, 51(1), 41–43.
- Muzikova, J., Horacek, J. (2003). The dry binders, *Vivapur 102*, *Vivapur 12* and the effect of magnesium stearate on the strength of tablets containing these substances. *Ceske Slov Farm*, 52(4), 176–180.
- Naggar, V.F. (1981). An *in vitro* study of the interaction between diazepam and some antacid or excipients. *Pharmazie*, 36(2), 114-7.
- Nail, S.M., Hem, S.L. (1982). Physical-chemical principles. in G.S. Banker, R.K. Chalmers, (Eds.), *Pharmaceutics and pharmacy practice* (pp.47-84), New York : Lippincott.
- Newton, D.W., Driscoll, D.F., Goudreau J.L., Ratanamaneichatara, S. (1981). Solubility characteristics of diazepam in aqueous admixture solutions: theory and practice. *American Journal of Hospital Pharmacy*, 38 (2), 179-182.
- Niazi Sarfaraz K. (2004a). Compressed Solid Formulations. in *Handbook of Pharmaceutical Manufacturing Formulations: Compressed Solid Products*. Volume 1 Part II pp. 59 New York : CRC Press LLC.
- Niazi Sarfaraz K. (2004b). Chapter 2 Guidance on Formulating Compressed Solids. in *Handbook of Pharmaceutical Manufacturing Formulations: Compressed Solid Products*. Volume 1 Part I pp.14, 16, 3656 New York : CRC Press LLC.
- Niazi Sarfaraz K. (2007). Solid-State Properties. in *Handbook of Preformulation. Chemical, Biological, and Botanical Drugs*, pp 218-231, New York: Informa Healthcare USA, Inc.
- Nozawa, Y., Mizumoto, T., Higashide, F. (1986). Increases in dissolution rate of nifedipine by roll-mixing with polyvinylpyrrolidone. *Pharm. Acta Hev.*, 61, 337-341.
- Omathanu, P. Perumal and Satheesh, K. Podaralla. (2008). 6.2 Role of Preformulation in Development of Solid Dosage Forms. in Shayne Cox Gad (Ed.), *Pharmaceutical Manufacturing Handbook: Production and Processes*. pp. 939. New Jersey : John Wiley & Sons, Inc.

- Otsuka, M. and Kaneniwa, N. (1986). Effect of seed crystals on solid-state transformation of chloramphenicol palmitate during grinding. *J. Pharm. Sci.*, 75, 506-511.
- Parrot E.L. (1989). Compression. in H.A. Lieberman, L. Lachman, J.B. Schwartz (Eds.) *Pharmaceutical Dosage Forms: Tablets 2nd edition* vol. 2 (pp. 202, 204, 219). New York : Marcel Dekker.
- Patel, H., Stalcup, A., Dansereau, R., Sakr, A. (2003). The effect of excipients on the stability of levothyroxine sodium pentahydrate tablets. *Int. J. Pharm.*, 264 , 35 – 43 .
- Peck Garnet E., Baley George J., McCurdy Vincent E., Banker Gilbert S. (1989). Tablet Formulation and Design. in H.A. Lieberman, L. Lachman, J.B. Schwartz (Eds.) *Pharmaceutical Dosage Forms: Tablets 2nd edition* vol 1 (pp. 91, 109). New York : Marcel Dekker.
- Pitha, J., Harman, S.M., Michel, M.E. (1986). Hydrophilic cyclodextrin derivatives enable effective oral administration of steroidal hormones. *J. Pharm. Sci.*, 75, 165-7.
- Proost, J.H., Bolhuis, G.K., Lerk, C.F. (1983). The effect of the swelling capacity of disintegrants on the in vitro and in vivo availability of diazepam tablets, containing magnesium stearate as a lubricant. *Int. J. Pharm.*, 13 (3), 287-296.
- Rabasco, A.M., Ginés, J.M., Fernández-Arévalo, M., Holgado, M.A. (1991). Dissolution rate of diazepam from polyethylene glycol 6000 solid dispersions. *Int. J. Pharm.*, 67 (2), 201-205.
- Ragnarsson, G., Holzer, A.W., Sjogren, J. (1979). The influence of mixing time and colloidal silica on the lubricating properties of magnesium stearate. *Int J Pharm.*, 3, 127–131.
- Rasenack, N., Hartenhauer, H., Müller, B.W. (2003). Microcrystals for dissolution rate enhancement of poorly water-soluble drugs. *Int. J. Pharm.*, 254 (2), 137-145.
- Rasool, A.A., Hussain, A.A., Dittert, L.W. (2006). Solubility enhancement of some water-insoluble drugs in the presence of nicotinamide and related compounds. *J. Pharm. Sci.*, 80 (4), 387 – 393

- Raymond, C.R., Paul J.S., Siân, C.O. (2006). *Pharmaceutical Excipients 5* CD-ROM
Published jointly by the Pharmaceutical Press and the American Pharmacists
Association.
- Rider, J.A. "Simethicone antacid tablet" U.S. Pat 4198390 Apr. 15, 1980.
- Rochville, Md. (2007). *The United States of Pharmacopoeia*, 30th, (pp. : 1726, 1728,
1912, 1915, <501> pp. 192, <711> pp. 277, <905> pp. 378, <1216> pp. 674,
<2040> pp. 728, <2091> pp. 731. The United States of Pharmacopoeia
convention Inc,
- Seo Anette, Holm Per, Kristensen Henning Gjelstrup and Schæfer Torben. (2003).
The preparation of agglomerates containing solid dispersions of diazepam by
melt agglomeration in a high shear mixer. *Int. J. Pharm.*, 259 (1-2), 161-171.
- Shah, T.J., Amin, A.F., Parikh, J.R., Parikh, R.H. (2007). Process Optimization and
Characterization of Poloxamer Solid Dispersions of a Poorly Water-soluble
Drug. *AAPS PharmSciTech.* 8(2), E1-E7, Article 29
(<http://www.aapspharmstech.org>).
- Shanbhag Anant, Rabel Shelley, Nauka Ewa, Casadevall Gemma, Shivanand Padmaja,
Eichenbaum Gary and Mansky Paul. (2008). Method for screening of solid
dispersion formulations of low-solubility compounds—Miniaturization and
automation of solvent casting and dissolution testing. *Int. J. Pharm.*, 351 (1-2),
209-218.
- Shangraw, R.F. (1989). Compressed Tablet by Direct Compression. in H.A.
Lieberman, L. Lachman, and J.B. Schwartz (Eds.) *Pharmaceutical Dosage
Forms: Tablets* 2nd edition vol 1 (pp. 153, 213). New York : Marcel Dekker.
- Shawky S, Meshia M S. (1988). Dissolution behavior of lorazepam crystallised from
aqueous surfactant solutions. *STP Pharma.* 4, 270-274.
- Sheikh-Salem, M., Fell, J.T. (1981). The influence of magnesium stearate on time
dependent strength changes in tablets. *Drug Dev. Ind. Pharm.*, 7, 669–674.
- Soliman, M.S., Khan M.A. (2005). Preparation and in vitro characterization of a semi-
solid dispersion of flurbiprofen with Gelucire 44/14 and Labrasol. *Pharmazie.*
60 (4), 288–293.
- Spireas, S., Sadu, S. (1998). Enhancement of prednisolone dissolution properties
using liquisolid compacts. *Int. J. Pharm.*, 166, 177-188.

- Supabphol, R., Stewart, P.J. (1996). Influence of the carrier on the intrinsic rate of dissolution of diazepam in interactive mixtures. *J. Pharm. Pharmacol.*, 48(12), 1249-1255.
- Swanepoel, E., Liebenberg, W., de Villiers, M.M., Dekker, T.G. (2000). Dissolution properties of piroxicam powders and capsules as a function of particle size and the agglomeration of powders. *Drug Dev. Ind. Pharm.*, 26 (10) : 1067- 1076.
- Sweetman Sean. (2007). *Martindale: The Complete Drug Reference 35* CD-ROM Prepared by the editorial staff of the Royal Pharmaceutical Society of Great Britain, the Pharmaceutical Press.
- Tokumura, T., Nanba, M., Tsushima, Y., Tatsuishi, K., Kayano, M., Machida, Y., Nagai, T. (1986). Enhancement of bioavailability of cinnarizine from its beta-cyclodextrin complex on oral administration with DL-phenylalanine as a competing agent. *J. Pharm. Sci.*, 75, 391-394.
- Uekama, K., Narisawa, S., Hirayama, F., Otagiri, M. (1983). Improvement of dissolution and absorption characteristics of benzodiazepines by cyclodextrin complexation. *Int. J. Pharm.*, 16 (3), 327-338.
- Varshosaz, J., Talari, R., Mostafavi, S.A., Nokhodchi A. (2008). Dissolution enhancement of gliclazide using *in situ* micronization by solvent change method. *Powder Technology*, 187 (3), 222-230.
- Verheyen, S., Blaton, N., Kinget, R., Van den Mooter, G. (2002). Mechanism of increased dissolution of diazepam and temazepam from polyethylene glycol 6000 solid dispersions. *Int. J. Pharm.*, 249 (1-2), 45-58.
- Verma, R.K., Garg, S. (2004). Compatibility studies between isosorbide mononitrate and selected excipients used in the development of extended release formulations. *J. Pharm. Biomed. Anal.*, 35, 449 – 458.
- Waard, H. de, Hinrichs, W.L.J., Visser, M.R., Bologna, C., Frijlink, H.W. (2008). Unexpected differences in dissolution behavior of tablets prepared from solid dispersions with a surfactant physically mixed or incorporated. *Int. J. Pharm.*, 349 (1-2), 66-73.
- Walker, S.E., Ganley, J.A., Bedford, K., Eaves, T. (1980). The filling of molten and thixotropic formulations into hard gelatin capsules. *J. Pharm. Pharmacol.*, 32, 389-393.

- Westerberg, M., Nyström, C. (1993). Physicochemical aspects of drug release XVII. The effect of drug surface area coverage to carrier materials on drug dissolution from ordered mixtures. *Int. J. Pharm.*, 90 (1), 1-17.
- White, H.S. (2000a). Histamine and Antihistaminic Drugs. in A. R Gennaro, and D. Limmer (Eds.), *Remington: The Science and Practice of Pharmacy*, 20th ed. (pp. 1466), Philadelphia: Lippincott Williams & Wilkins.
- White, H.S. (2000b). Sedative and Hypnotic Drugs. in A. R Gennaro, and D. Limmer (Eds.), *Remington: The Science and Practice of Pharmacy*, 20th ed. (pp. 1410), Philadelphia : Lippincott Williams & Wilkins.
- Wierik, G.H.P.Te, Eissens, A.C., Lerk, C.F. (1993). Preparation, characterization and pharmaceutical application of linear dextrans: IV. Drug release from capsules and tablets containing amylopectin. *Int. J. Pharm.*, 98 (1-3), 219-224.
- Wong, S.M., Kellaway, I.W., Murdan, S. (2006a). Fast-dissolving micro-particles fail to show improved oral bioavailability. *J. Pharm. Pharmacol.*, 58 (8), 1319-1326
- Wong, S.M., Kellaway, I.W., Murdan, S. (2006b). Enhancement of the dissolution rate and oral absorption of a poorly water soluble drug by formation of surfactant-containing microparticles. *Int. J. Pharm.*, 317 (1), 61-68.
- Yamamoto, K., Piyarom, S. (2000). Method used of improving dissolution and absorption characteristics of poorly-soluble drugs. *Srinakharinwirot Bhesajarn*, 5, 84-100.
- Yüksel Nilüfer, Karata Aysegül, Özkan Yalçın, Savaş er Ayhan, Özkan Sibel A., Baykara Tamer. (2003). Enhanced bioavailability of piroxicam using Gelucire 44/14 and Labrasol: *in vitro* and *in vivo* evaluation. *European Journal of Pharmaceutics and Biopharmaceutics*, 56 (3), 453-459.
- Zhao, F.Y., Stewart PJ. (2003). De-agglomeration of micronized benzodiazepines in dissolution media measured by laser diffraction particle sizing. *J. Pharm. Pharmacol.*, 55(6), 749-755.
- Zhao F.Y., Stewart PJ. (2004). Modeling the deagglomeration of micronized benzodiazepines from powder mixtures added to dissolution media. *J. Pharm. Sci.*, 93(6), 1618-1627.