

TABLE OF CONTENTS

| | Page |
|---|-------------|
| ACKNOWLEDGEMENTS | iii |
| ENGLISH ABSTRACT | iv |
| THAI ABSTRACT | vi |
| LIST OF TABLES | xiii |
| LIST OF FIGURES | xvi |
| ABBREVIATIONS AND SYMBOLS | xxv |
| CHAPTER I INTRODUCTION | 1 |
| 1.1 State of significant of the problem | 1 |
| 1.2 Objectives | 4 |
| 1.3 Scope of Study | 4 |
| 1.4 Literature Reviews | 5 |
| 1.4.1 Model Drugs | 5 |
| 1.4.1.1 Chlorpheniramine maleate | 5 |
| 1.4.1.2 Diazepam | 8 |
| 1.4.2 Tablet excipients | 10 |
| 1.4.2.1 Dicalcium phosphate dihydrate | 10 |
| 1.4.2.2 Lactose monohydrate | 14 |
| 1.4.2.3 Croscamellose sodium | 17 |
| 1.4.2.4 Polyvinyl pyrrollidone K90 | 20 |
| 1.4.2.5 Magnesium stearate | 23 |
| 1.4.2.6 Talc | 25 |
| 1.4.3 Solvents | 27 |
| 1.4.3.1 Ethyl alcohol | 27 |
| 1.4.3.2 Dichloromethane | 29 |
| 1.4.4 Tablet Manufacturing | 30 |

ลิขสิทธิ์มหาวิทยาลัยเชียงใหม่
Copyright © by Chiang Mai University
All rights reserved

| | Page |
|---|-------------|
| 1.4.4.1 Wet granulation | 30 |
| 1.4.4.2 Dry granulation | 32 |
| 1.4.4.3 Direct compression | 32 |
| 1.4.5 Tablet Evaluation | 34 |
| 1.4.5.1 Weight variation | 34 |
| 1.4.5.2 Hardness testing | 35 |
| 1.4.5.3 Tablet friability | 35 |
| 1.4.5.4 Tablet porosity | 35 |
| 1.4.5.5 Disintegration test | 36 |
| 1.4.5.6 Determination of the active ingredient content | 37 |
| 1.4.5.7 Uniformity of dosage unit | 37 |
| 1.4.6 Tablet Characterization | 37 |
| 1.4.7 Improving Drug Dissolutions | 40 |
| 1.4.8 Improvement of dissolution characteristic of diazepam | 46 |
| 1.4.9 Drug-Solution-Dropping Tablet (DSDT) | 50 |
| CHAPTER II RESEARCH DESIGNS AND METHODS | |
| EXPERIMENTAL | 57 |
| 2.1 Materials | 57 |
| 2.2 Equipments | 57 |
| 2.3 Methods | 58 |
| 2.3.1 Preparation of blank tablet | 58 |
| 2.3.2 Preparation of drug tablet by conventional method | 60 |
| 2.3.3 Evaluation of blank tablet properties | 60 |
| 2.3.3.1 Weight variation | 60 |
| 2.3.3.2 Hardness test | 60 |
| 2.3.3.3 Friability | 61 |
| 2.3.3.4 Tablet porosity | 61 |
| 2.3.3.5 Disintegration test | 61 |
| 2.3.4 Preparation of drug-solution-dropping tablet | 62 |
| 2.3.5 Study of the penetration depth of drug solution | 62 |

| | Page |
|--|-------------|
| 2.3.6 Tablet characterization studies | 63 |
| 2.3.6.1 Morphology study by digital camera | 63 |
| 2.3.6.2 Morphology study by scanning electron microscopy | 63 |
| 2.3.6.3 X-ray powder diffraction | 63 |
| 2.3.6.4 Differential scanning calorimetry (DSC) | 63 |
| 2.3.7 Assay of active ingredients | 64 |
| 2.3.8 Content Uniformity Testing | 64 |
| 2.3.9 Dissolution Profile Studies | 64 |
| 2.3.10 Stability Studies | 65 |
| CHAPTER III RESULTS | 66 |
| 3.1 Evaluation of blank tablet properties | 66 |
| 3.1.1 Weight variation | 66 |
| 3.1.2 Hardness testing | 68 |
| 3.1.3 Tablet friability | 69 |
| 3.1.4 Tablet porosity | 70 |
| 3.1.5 Disintegration test | 71 |
| 3.2 Study of the penetration of solution after dropping on blank tablet | 71 |
| 3.3 Tablet characterization studies | 72 |
| 3.3.1 Morphology studies by digital camera | 73 |
| 3.3.2 Morphology studies by scanning electron microscope | 83 |
| 3.3.3 X-ray powder diffraction of powder | 106 |
| 3.3.4 Differential scanning calorimetry | 126 |
| 3.4 Determination of the active ingredient content | 131 |
| 3.5 Content uniformity | 131 |
| 3.6 Dissolution Profiles Study | 132 |
| 3.6.1 Effect of times of solvent dropping | 132 |
| 3.6.2 Dissolution profiles of CPM-solution-dropping tablet | 135 |

| | Page |
|---|-------------|
| 3.6.3 Dissolution profiles of DZP-solution-dropping tablet | 139 |
| 3.6.4 Effect of compression force on drug dissolution of DSDT | 144 |
| 3.6.5 Comparison of DSDT dissolution profiles of tablets prepared from DC and from WG blank tablet | 147 |
| 3.7 Stability Studies | 150 |
| 3.7.1 Drug contents | 150 |
| 3.7.2 Dissolution profiles | 150 |
| CHAPTER IV DISCUSSION | 159 |
| 4.1 Evaluation of blank tablet properties | 159 |
| 4.1.1 Weight variation | 159 |
| 4.1.2 Hardness testing | 159 |
| 4.1.3 Tablet friability | 160 |
| 4.1.4 Tablet porosity | 160 |
| 4.1.5 Disintegration test | 161 |
| 4.2 Study of the depth of solution after dropping on blank tablet | 162 |
| 4.3 Tablet characterization studies | 162 |
| 4.3.1 Morphology studies by digital camera | 162 |
| 4.3.2 Morphology studies by scanning electron microscope | 163 |
| 4.3.2.1 Morphology of tablet surface | 163 |
| 4.3.2.2 Morphology of powder scratched from superficial and deeper surface of drug-solution-dropping tablet | 164 |
| 4.3.3 XRPD of powder of the tablet compositions and the powder from DSDT | 165 |
| 4.3.4 Differential scanning calorimetry analysis | 167 |
| 4.4 Determination of the active ingredient content | 168 |
| 4.5 Content uniformity testing | 168 |

| | Page |
|---|-------------|
| 4.6 Dissolution studies | 168 |
| 4.6.1 Effect of time of solvent dropping | 168 |
| 4.6.2 Dissolution profiles of CPM-solution-dropping tablet | 169 |
| 4.6.3 Dissolution profiles of DZP-solution-dropping tablet | 170 |
| 4.6.4 Effect of compression force on drug dissolution of DSDT | 172 |
| 4.6.5 Comparison of DSDT dissolution profiles of tablets prepared from DC and from WG blank tablet | 173 |
| 4.7 Stability Studies | 174 |
| 4.7.1 Drug content | 174 |
| 4.7.2 Dissolution profiles | 174 |
| CHAPTER V CONCLUSION | 176 |
| REFERENCES | 182 |
| APPENDICES | 195 |
| APPENDIX A | 196 |
| APPENDIX B | 197 |
| APPENDIX C | 198 |
| APPENDIX D | 213 |
| CURRICULUM VITAE | 214 |

LIST OF TABLES

| Table | Page |
|---|-------------|
| 1.1 Liquid -liquid partitioning data percent of chlorpheniramine maleate in organic phase | 7 |
| 1.2 Typical properties of dibasic calcium phosphate dihydrate | 12 |
| 1.3 Typical properties of lactose monohydrate | 16 |
| 1.4 Typical properties of croscamellose sodium | 18 |
| 1.5 Typical properties of povidone | 21 |
| 1.6 Typical properties of magnesium stearate | 24 |
| 1.7 Typical properties of talc | 26 |
| 1.8 Typical properties of absolute alcohol | 29 |
| 1.9 Approaches to improving the dissolution properties of poorly soluble drugs to be given orally | 41 |
| 2.1 The DC blank tablet formulation | 59 |
| 2.2 The WG blank tablet formulation | 59 |
| 3.1 Weight variation of DC and WG blank tablet of all tested CF | 67 |
| 3.2 Weight Variation Tolerances for Uncoated Tablets | 68 |
| 3.3 Hardness of DC and WG blank tablet of all tested CF | 69 |
| 3.4 Friability of DC and WG blank tablet of all tested CF | 70 |
| 3.5 The amount of CPM and DZP in 10 tablets of CPM- and DZO-solution-dropping tablet which prepared from DC blank tablet at 1000 kg of CF | 132 |
| 3.6 The amount of CPM and diazepam in DSDT after 0, 1, 2 and 3 months of the storage | 150 |
| A1 Porosity of DC and WG blank tablet of all tested CF | 196 |
| B1 Disintegration time of DC and WG blank tablet of all tested CF | 197 |
| C1 Average percent dissolved of CPM from DSDT prepared from DC blank tablet of 1000 kg CF following 0, 2, 4 and 6 times of dropping solvent | 198 |

| Table | Page |
|---|-------------|
| C2 Average percent dissolved of CPM from DSDT prepared from WG blank tablet of 1000 kg CF following 0, 2, 4 and 6 times of dropping solvent | 198 |
| C3 Average percent dissolved of DZP from DSDT prepared from DC blank tablet of 1000 kg CF following 0, 2, 4 and 6 times of dropping solvent | 199 |
| C4 Average percent dissolved of DZP from DSDT prepared from WG blank tablet of 1000 kg CF following 0, 2, 4 and 6 times of dropping solvent | 199 |
| C5 Average percent dissolved of CPM from DSDT prepared from DC blank tablet of all tested CF and the commercial CPM tablet | 200 |
| C6 Average percent dissolved of CPM DC tablet of all tested CF and the commercial CPM tablet | 200 |
| C7 Average percent dissolved of CPM from DSDT prepared from WG blank tablet of all tested CF and the commercial CPM tablet | 201 |
| C8 Average percent dissolved of CPM WG tablet of all tested CF and the commercial CPM tablet | 201 |
| C9 Average percent dissolved of DZP from DSDT prepared from DC blank tablet of all tested CF and the commercial DZP tablet | 202 |
| C10 Average percent dissolved of DZP DC tablet of all tested CF and the commercial DZP tablet | 202 |
| C11 Average percent dissolved of DZP from DSDT prepared from WG blank tablet of all tested CF and the commercial DZP tablet | 203 |
| C12 Average percent dissolved of DZP WG tablet of all tested CF and the commercial DZP tablet | 203 |
| C13 Average percent dissolved of CPM from DSDT prepared from DC blank tablet of 1000 kg CF at the beginning (0), 1, 2 and 3 months storage | 204 |
| C14 Average percent dissolved of CPM from DSDT prepared from DC blank tablet of 1400 kg CF at the beginning (0), 1, 2 and 3 months storage | 204 |
| C15 Average percent dissolved of CPM from DSDT prepared from DC blank tablet of 1800 kg CF at the beginning (0), 1, 2 and 3 months storage | 205 |
| C16 Average percent dissolved of CPM from CPM DC tablet of 1000 kg CF at the beginning (0), 1, 2 and 3 months storage | 205 |

| Table | Page |
|--|-------------|
| C17 Average percent dissolved of CPM from CPM DC tablet of 1400 kg CF at the beginning (0), 1, 2 and 3 months storage | 206 |
| C18 Average percent dissolved of CPM from CPM DC tablet of 1800 kg CF at the beginning (0), 1, 2 and 3 months storage | 206 |
| C19 Average percent dissolved of CPM from DSDT prepared from WG blank tablet of 1000 kg CF at the beginning (0), 1, 2 and 3 months storage | 207 |
| C20 Average percent dissolved of CPM from DSDT prepared from WG blank tablet of 1400 kg CF at the beginning (0), 1, 2 and 3 months storage | 207 |
| C21 Average percent dissolved of DZP from DSDT prepared from DC blank tablet of 1000 kg CF at the beginning (0), 1, 2 and 3 months storage | 208 |
| C22 Average percent dissolved of DZP from DSDT prepared from DC blank tablet of 1400 kg CF at the beginning (0), 1, 2 and 3 months storage | 208 |
| C23 Average percent dissolved of DZP from DSDT prepared from DC blank tablet of 1800 kg CF at the beginning (0), 1, 2 and 3 months storage | 209 |
| C24 Average percent dissolved of DZP from DZP DC tablet of 1000 kg CF at the beginning (0), 1, 2 and 3 months storage | 209 |
| C25 Average percent dissolved of DZP from DZP DC tablet of 1400 kg CF at the beginning (0), 1, 2 and 3 months storage | 210 |
| C26 Average percent dissolved of DZP from DZP DC tablet of 1800 kg CF at the beginning (0), 1, 2 and 3 months storage | 210 |
| C27 Average percent dissolved of DZP from DSDT prepared from WG blank tablet of 1000 kg CF at the beginning (0), 1, 2 and 3 months storage | 211 |
| C28 Average percent dissolved of DZP from DSDT prepared from WG blank tablet of 1400 kg CF at the beginning (0), 1, 2 and 3 months storage | 211 |
| C29 Average percent dissolved of DZP from DSDT prepared from WG blank tablet of 1800 kg CF at the beginning (0), 1, 2 and 3 months storage | 212 |

LIST OF FIGURES

| Figure | Page |
|---|------|
| 1.1 Chemical Structure of Chlorphenamine maleate | 6 |
| 1.2 Chemical Structure of Diazepam | 9 |
| 1.3 Structure Formula of DCP dihydrate | 11 |
| 1.4 SEM of dibasic calcium phosphate dihydrate, coarse grade | 12 |
| 1.5 Structure Formula of α lactose monohydrate | 14 |
| 1.6 SEM of Pharmatose | 15 |
| 1.7 Structure Formula of Carboymethylcellulose sodium | 17 |
| 1.8 SEM of Croscarmellose sodium (Ac-Di-Sol [®]) | 18 |
| 1.9 Structure Formula of Povidone | 20 |
| 1.10 SEM of Povidone K-30 (<i>Plasdone K-30</i>) | 21 |
| 1.11 SEM of Magnesium stearate | 23 |
| 1.12 SEM of Talc (<i>Purталc</i>) | 26 |
| 1.13 Structure Formula of ethyl alcohol | 28 |
| 1.14 Structure Formula of dichloromethane | 29 |
| 1.15 Diagram of wet granulation process | 30 |
| 1.16 Diagram of dry granulation process | 32 |
| 1.17 Diagram of direct compression | 33 |
| 1.18 A force-porosity diagram | 36 |
| 1.19 Characterization of amorphous form | 39 |
| 3.1 Effect of compression force on the porosity of DC and WG blank tablet | 70 |
| 3.2 Effect of compression force on disintegration time of DC and WG blank tablet | 71 |
| 3.3 The DC blank tablet at 1000 kg of CF after dropping 0.1% fast green FCF in alcohol | 72 |
| 3.4 The aspect of the surface of DC blank tablet of all test CF after dropping the solvent without drug | 74 |

| Figure | Page |
|--|-------------|
| 3.5 The surface of WG blank tablet at all tested CF after dropping CPM solution using a mixture of absolute alcohol and dichloromethane (1:3 by volume) as the solvent | 75 |
| 3.6 The surface of DSDT compared with DC blank tablet of 1000 kg CF | 76 |
| 3.7 The surface of DSDT compared with DC blank tablet of 1400 kg CF | 77 |
| 3.8 The surface of DSDT compared with DC blank tablet of 1800 kg CF | 78 |
| 3.9 The surface of DSDT compared with WG blank tablet of 1000 kg CF | 79 |
| 3.10 The surface of DSDT compared with WG blank tablet of 1400 kg CF | 80 |
| 3.11 The surface of DSDT compared with WG blank tablet of 1800 kg CF | 81 |
| 3.12 Layer cover of the surface of CPM-solution-dropping tablet prepared from WG blank tablet of 1800 kg CF | 82 |
| 3.13 Small crystals of diazepam on the surface of DZP-solution-dropping tablet prepared from WG blank tablet of 1800 kg CF | 82 |
| 3.14 The SEM morphology of upper surface of the tablet compressed at 1000 kg CF:- (a) DC blank tablet, (b) CPM-solution-dropping tablet and (c) DZP-solution-dropping tablet prepared from DC blank tablet | 85 |
| 3.15 The SEM morphology of the surface of the tablet compressed at 1400 kg CF:- (a) DC blank tablet, (b) CPM-solution-dropping tablet and (c) DZP-solution-dropping tablet prepared from DC blank tablet | 86 |
| 3.16 The SEM morphology of the surface of the tablet compressed at 1800 kg CF:- (a) DC blank tablet, (b) CPM-solution-dropping tablet and (c) DZP-solution-dropping tablet prepared from DC blank tablet | 87 |
| 3.17 The SEM morphology of the surface of the tablet compressed at 1000 kg CF:- (a) WG blank tablet, (b) CPM-solution-dropping tablet and (c) DZP-solution-dropping tablet prepared from DC blank tablet | 88 |
| 3.18 The SEM morphology of the surface of the tablet compressed at 1400 kg CF:- (a) WG blank tablet, (b) CPM-solution-dropping tablet and (c) DZP-solution-dropping tablet prepared from DC blank tablet | 89 |
| 3.19 The SEM morphology of the surface of the tablet compressed at 1800 kg CF:- (a) WG blank tablet, (b) CPM-solution-dropping tablet and (c) DZP-solution-dropping tablet prepared from DC blank tablet | 90 |

| Figure | Page |
|---|-------------|
| 3.20 SEM morphology of DCP dihydrate with magnification (a) 100X (b) 500X (c) 1000X | 92 |
| 3.21 SEM morphology of Ac-Di-Sol [®] with magnification (a) 100X (b) 500X (c) 1000X | 93 |
| 3.22 SEM morphology of Pharmatose [®] with magnification (a) 100X (b) 500X (c) 1000X | 94 |
| 3.23 SEM morphology of magnesium stearate with magnification (a) 100X (b) 500X (c) 1000X | 95 |
| 3.24 SEM morphology of talcum with magnification (a) 100X (b) 500X (c) 1000X | 96 |
| 3.25 SEM morphology of CPM with magnification (a) 100X (b) 500X (c) 1000X | 97 |
| 3.26 SEM morphology of diazepam with magnification (a) 100X (b) 500X (c) 1000X | 98 |
| 3.27 SEM morphology of powder scratched from near and deeper surface of CPM-solution-dropping tablet prepared from DC blank tablet compressed at 1000 kg CF | 99 |
| 3.28 SEM morphology of powder scratched from near and deeper surface of CPM-solution-dropping tablet prepared from DC blank tablet compressed at 1400 kg CF | 100 |
| 3.29 SEM morphology of powder scratched from near and deeper surface of CPM-solution-dropping tablet prepared from DC blank tablet compressed at 1800 kg CF | 101 |
| 3.30 SEM morphology of powder scratched from near and deeper surface of CPM-solution-dropping tablet prepared from WG blank tablet compressed at 1000 kg CF | 102 |
| 3.31 SEM morphology of powder scratched from near and deeper surface of CPM-solution-dropping tablet prepared from WG blank tablet compressed at 1400 kg CF | 103 |

| Figure | Page |
|---|-------------|
| 3.32 SEM morphology of powder scratched from near surface of DZP-solution-dropping tablet prepared from DC blank tablet (magnification 1000X) compressed at (a) 1000 kg (b) 1400 kg (c) 1800 kg | 104 |
| 3.33 SEM morphology of powder scratched from near surface of DZP-solution-dropping tablet prepared from WG blank tablet (magnification 1000X) compressed at (a) 1000 kg (b) 1400 kg (c) 1800 kg | 105 |
| 3.34 Powdered X-ray diffraction pattern of CPM | 107 |
| 3.35 Powdered X-ray diffraction pattern of diazepam | 107 |
| 3.36 Powdered X-ray diffraction pattern of DCP dihydrate | 108 |
| 3.37 Powdered X-ray diffraction pattern of Pharmatose [®] | 108 |
| 3.38 Powdered X-ray diffraction pattern of Ac-Di-Sol [®] | 109 |
| 3.39 Powdered X-ray diffraction pattern of magnesium stearate | 109 |
| 3.40 Powdered X-ray diffraction pattern of talcum | 110 |
| 3.41 Powdered XRD pattern of scratched surface powder from CPM-solution-dropping tablet prepared from DC blank tablet of 1000 kg CF | 110 |
| 3.42 Powdered XRD pattern of scratched surface powder from CPM-solution-dropping tablet prepared from DC blank tablet of 1400 kg CF | 111 |
| 3.43 Powdered XRD pattern of scratched surface powder from CPM-solution-dropping tablet prepared from DC blank tablet of 1800 kg CF | 111 |
| 3.44 Powdered XRD pattern of scratched surface powder from CPM-solution-dropping tablet prepared from WG blank tablet of 1000 kg CF | 112 |
| 3.45 Powdered XRD pattern of scratched surface powder from CPM-solution-dropping tablet prepared from WG blank tablet of 1400 kg CF | 112 |
| 3.46 Powdered XRD pattern of scratched surface powder from CPM-solution-dropping tablet prepared from WG blank tablet of 1800 kg CF | 113 |
| 3.47 Powdered XRD pattern of scratched surface powder from DZP-solution-dropping tablet prepared from DC blank tablet of 1000 kg CF | 113 |
| 3.48 Powdered XRD pattern of scratched surface powder from DZP-solution-dropping tablet prepared from DC blank tablet of 1400 kg CF | 114 |
| 3.49 Powdered XRD pattern of scratched surface powder from DZP-solution-dropping tablet prepared from DC blank tablet of 1800 kg CF | 114 |

| Figure | Page |
|--|-------------|
| 3.50 Powdered XRD pattern of scratched surface powder from DZP-solution-dropping tablet prepared from WG blank tablet of 1000 kg CF | 115 |
| 3.51 Powdered XRD pattern of scratched surface powder from DZP-solution-dropping tablet prepared from WG blank tablet of 1400 kg CF | 115 |
| 3.52 Powdered XRD pattern of scratched surface powder from DZP-solution-dropping tablet prepared from WG blank tablet of 1800 kg CF | 116 |
| 3.53 Single crystal XRD pattern of powder from the surface of DC blank tablet compressed at 1000 Kg CF | 117 |
| 3.54 Single crystal XRD pattern of powder from the surface of DC blank tablet compressed at 1400 Kg CF | 118 |
| 3.55 Single crystal XRD pattern of powder from the surface of DC blank tablet compressed at 1800 Kg CF | 118 |
| 3.56 Single crystal XRD pattern of powder from the surface of WG blank tablet compressed at 1000 Kg CF | 119 |
| 3.57 Single crystal XRD pattern of powder from the surface of WG blank tablet compressed at 1400 Kg CF | 119 |
| 3.58 Single crystal XRD pattern of powder from the surface of WG blank tablet compressed at 1800 Kg CF | 120 |
| 3.59 Single crystal XRD pattern of the powder from the surface of CPM-solution-dropping tablet prepared from DC blank tablet compressed at 1000 Kg of CF | 120 |
| 3.60 Single crystal XRD pattern of the powder from the surface of CPM-solution-dropping tablet prepared from DC blank tablet compressed at 1400 Kg of CF | 121 |
| 3.61 Single crystal XRD pattern of the powder from the surface of CPM-solution-dropping tablet prepared from DC blank tablet compressed at 1800 Kg of CF | 121 |
| 3.62 Single crystal XRD pattern of the powder from the surface of DZP-solution-dropping tablet prepared from DC blank tablet compressed at 1000 Kg of CF | 122 |

| Figure | Page |
|---|-------------|
| 3.63 Single crystal XRD pattern of the powder from the surface of DZP- solution-dropping tablet prepared from DC blank tablet compressed at 1400 Kg of CF | 122 |
| 3.64 Single crystal XRD pattern of the powder from the surface of DZP- solution-dropping tablet prepared from DC blank tablet compressed at 1800 Kg of CF | 123 |
| 3.65 Single crystal XRD pattern of the powder from the surface of CPM- solution-dropping tablet prepared from WG blank tablet compressed at 1000 Kg of CF | 123 |
| 3.66 Single crystal XRD pattern of the powder from the surface of CPM- solution-dropping tablet prepared from WG blank tablet compressed at 1400 Kg of CF | 124 |
| 3.67 Single crystal XRD pattern of the powder from the surface of CPM- solution-dropping tablet prepared from WG blank tablet compressed at 1800 Kg of CF | 124 |
| 3.68 Single crystal XRD pattern of the powder from the surface of DZP- solution-dropping tablet prepared from WG blank tablet compressed at 1000 Kg of CF | 125 |
| 3.69 Single crystal XRD pattern of the powder from the surface of DZP- solution-dropping tablet prepared from WG blank tablet compressed at 1400 Kg of CF | 125 |
| 3.70 Single crystal XRD pattern of the powder from the surface of DZP- solution-dropping tablet prepared from WG blank tablet compressed at 1800 Kg of CF | 126 |
| 3.71 DSC thermogram of CPM-solution-dropping tablet prepared from DC blank tablet compacted at 1000 kg CF | 127 |
| 3.72 DSC thermogram of CPM | 128 |
| 3.73 DSC thermogram of DCP dihydrate | 128 |
| 3.74 DSC thermogram of CPM-solution-dropping tablet prepared from WG blank tablet compacted at 1000 kg CF | 129 |
| 3.75 DSC thermogram of Pharmatose® | 129 |

| Figure | Page |
|---|-------------|
| 3.76 DSC thermogram of DZP-solution-dropping tablet prepared from DC blank tablet compressed at 1000 kg CF | 130 |
| 3.77 DSC thermogram of diazepam | 130 |
| 3.78 DSC thermogram of DZP-solution-dropping tablet prepared from WG blank tablet compressed at 1000 kg CF | 131 |
| 3.79 CPM dissolution profiles of the CPM-solution dropping tablet prepared from DC blank tablet at the same 1000 kg CF after repeat blank solvent dropping 0, 2, 4 or 6 times | 133 |
| 3.80 CPM dissolution profiles of the CPM-solution dropping tablet prepared from WG blank tablet at the same 1000 kg CF after repeat blank solvent dropping 0, 2, 4 or 6 times | 134 |
| 3.81 DZP dissolution profiles of the DZP-solution dropping tablet prepared from DC blank tablet at the same 1000 kg CF after repeat solvent dropping 0, 2, 4 or 6 times | 134 |
| 3.82 DZP dissolution profiles of the DZP-solution dropping tablet prepared from WG blank tablet at the same 1000 kg CF after repeat blank solvent dropping 0, 2, 4 or 6 times | 135 |
| 3.83 CPM dissolution profiles of CPM-solution-dropping tablet prepared from DC blank tablet compared with CPM DC tablet at the same 1000 kg CF and commercial CPM tablet | 136 |
| 3.84 CPM dissolution profiles of CPM-solution-dropping tablet prepared from DC blank tablet compared with CPM DC tablet at the same 1400 kg CF and commercial CPM tablet | 136 |
| 3.85 CPM dissolution profiles of CPM-solution-dropping tablet prepared from DC blank tablet compared with CPM DC tablet at the same 1800 kg CF and commercial CPM tablet | 137 |
| 3.86 CPM dissolution profiles of CPM-solution-dropping tablet prepared from WG blank tablet compared with CPM WG tablet at the same 1000 kg CF and commercial CPM tablet | 138 |

| Figure | Page |
|---|-------------|
| 3.87 CPM dissolution profiles of CPM-solution-dropping tablet prepared from WG blank tablet compared with CPM WG tablet at the same 1400 kg CF and commercial CPM tablet | 138 |
| 3.88 CPM dissolution profiles of CPM-solution-dropping tablet prepared from WG blank tablet compared with CPM WG tablet at the same 1800 kg CF and commercial CPM tablet | 139 |
| 3.89 DZP dissolution profiles of DZP-solution-dropping tablet prepared from DC blank tablet compared with DZP DC tablet at the same 1000 kg CF and commercial diazepam tablet | 140 |
| 3.90 DZP dissolution profiles of DZP-solution-dropping tablet prepared from DC blank tablet compared with DZP DC tablet at the same 1400 kg CF and commercial diazepam tablet | 141 |
| 3.91 DZP dissolution profiles of DZP-solution-dropping tablet prepared from DC blank tablet compared with DZP DC tablet at the same 1800 kg CF and commercial diazepam tablet | 141 |
| 3.92 DZP dissolution profiles of DZP-solution-dropping tablet prepared from WG blank tablet compared with DZP WG tablet at the same 1000 kg CF and commercial diazepam tablet | 142 |
| 3.93 DZP dissolution profiles of DZP-solution-dropping tablet prepared from WG blank tablet compared with DZP WG tablet at the same 1400 kg CF and commercial diazepam tablet | 143 |
| 3.94 DZP dissolution profiles of DZP-solution-dropping tablet prepared from WG blank tablet compared with DZP WG tablet at the same 1800 kg CF and commercial diazepam CPM tablet | 143 |
| 3.95 The effect of CFs on CPM dissolution profiles of CPM-solution-dropping tablet prepared from DC blank tablet | 144 |
| 3.96 The effect of CFs on CPM dissolution profiles of CPM-solution-dropping tablet prepared from WG blank tablet | 145 |
| 3.97 The effect of CFs on DZP dissolution profiles of DZP-solution-dropping tablet prepared from DC blank tablet | 146 |


| Figure | Page |
|---|-------------|
| 3.98 The effect of CFs on DZP dissolution profiles of DZP-solution-dropping tablet prepared from WG blank tablet | 146 |
| 3.99 CPM dissolution profiles of CPM-solution-dropping tablet prepared from DC blank tablet and WG blank tablet at all tested CF | 148 |
| 3.100 DZP dissolution profiles of DZP-solution-dropping tablet prepared from DC blank tablet and WG blank tablet at all tested CF | 149 |
| 3.101 Dissolution profiles of CPM-solution dropping tablet prepared from DC blank tablet at all tested CF after storage 0, 1, 2 and 3 months | 152 |
| 3.102 Dissolution profiles of CPM DC tablet at all tested CF after storage 0, 1, 2 and 3 months | 153 |
| 3.103 Dissolution profiles of CPM-solution-dropping tablet prepared from WG blank tablet at 1000 and 1400 kg of CF after storage 0, 1, 2 and 3 months | 154 |
| 3.104 Dissolution profiles of DZP-solution-dropping tablet prepared from DC blank tablet at all tested CF after storage 0, 1, 2 and 3 months | 156 |
| 3.105 Dissolution profiles of DZP DC tablet at all tested CF after storage 0, 1, 2 and 3 months | 157 |
| 3.106 Dissolution profiles of DZP-solution-dropping tablet prepared from WG blank tablet of all tested CF after storage 0, 1, 2 and 3 months | 158 |
| 4.1 Disintegration and dissolution pathways of solid dosage forms for absorption of drug | 162 |

ABBREVIATIONS AND SYMBOLS

| | |
|--------------------|---|
| μL | microlitre |
| $^{\circ}\text{C}$ | degree celsius |
| % | percent |
| RSD | relative standard deviation (%) |
| AV | average |
| BP | British Pharmacopoeia |
| CF | compression force |
| CPM | chlorpheniramine maleate |
| CPM-SDT | chlorpheniramine maleate-solution-dropping tablet |
| DC | direct compression |
| DCP | dicalcium phosphate dihydrate |
| DSC | differential scanning calorimetry |
| DSDT | drug-solution-dropping tablet |
| DZP | diazepam |
| DZP-SDT | diazepam-solution-dropping tablet |
| g | gram |
| GPO | Government Pharmaceutical Organization |
| kg | kilogram |
| L | liter |
| mg | milligram |
| mL | milliliter |
| mm | millimeter |
| PEG | polyethylene glycol |

ลิขสิทธิ์มหาวิทยาลัยเชียงใหม่
Copyright © by Chiang Mai University
All rights reserved

| | |
|------|---|
| rpm | round per minute |
| SE | standard error of the mean or error bar |
| SEM | scanning electron microscopy |
| SD | standard deviation |
| WG | wet granulation |
| USA | United State of America |
| USP | United State Pharmacopoeia |
| XRD | X-ray diffraction |
| XRPD | X-ray powder diffraction |



ลิขสิทธิ์มหาวิทยาลัยเชียงใหม่
Copyright© by Chiang Mai University
All rights reserved