

	Page
Chapter 3 Materials and methods	29
3.1 Materials	29
3.1.1 Chemicals	29
3.1.2 Animals	29
3.1.3 Microorganism	29
3.1.4 Equipments	30
3.1.5 Other Materials	30
3.2 Methods	30
3.2.1 Determination of loss on drying	30
3.2.2 Determination of viscosity-average molecular weight (\bar{M}_v)	31
3.2.3 Determination of degree of deacetylation	37
3.2.4 Preparation and characterization of chitosan solution	37
3.2.5 Preparation of ophthalmic formulations	40
3.2.5.1 Design of compatibility and stability studies	40
3.2.5.2 Validation of the UV spectrophotometer	41
3.2.5.3 Determination of vancomycin concentration and pH	41
3.2.6 Minimum inhibitory concentration analysis	41
3.2.7 Kinematic viscosity measurements	43
3.2.8 Pharmacokinatics of topically applied vancomycin in rabbit eyes	44
3.2.8.1 Topical administration and sampling	44
3.2.8.2 Determination of vancomycin concentration in tears	46
Chapter 4 Results and discussion	48
4.1 Loss on drying of chitosan	48
4.2 Viscosity-average molecular weight (\bar{M}_v) of chitosan	48
4.3 Degree of deacetylation of chitosan	50

	Page
4.4 Preparation of chitosan solution	51
4.4.1 Viscosity stability of 0.1% chitosan solution	54
4.4.2 Viscosity stability of 0.3% chitosan solution	56
4.5 Preparation ophthalmic formulations	61
4.5.1 Compatibility and stability studies	61
4.5.2 Validation of the UV spectrophotometer	62
4.5.3 Stability of vancomycin eye drops at 50 mg/ml in each vehicle	72
4.6 Minimum inhibitory concentrations of vancomycin at 50 mg/ml in Tears Naturale II™, sodium chloride and chitosan solutions stored at different temperatures over time	75
4.7 Kinematic viscosity measurements	79
4.8 Pharmacokinetics of topically applied vancomycin in rabbit eyes	81
Chapter 5 Summary and conclusion	93
References	95
Appendices	105
Appendix A The raw data of viscosity-average molecular weight (\overline{M}_v) of chitosan were determined by dilute-solution no 2 and 3.	106
Appendix B The raw data of assay validation reports	110
Appendix C Storage Temperature (United States Pharmacopeia)	126
Appendix D Terms describe of the pharmacokinetics	128
Appendix E Certificate of approval for use of animals	131
Faculty of Medicine, Chiang Mai University	
Curriculum Vitae	133

LIST OF TABLES

Table		Page
2.1	Forms of chitosan investigated in ophthalmology	14
2.2	Micro-organisms isolated from corneal specimens	17
2.3	Risk factors for corneal ulcers and typical organisms involved	18
2.4	Frequency of predisposing risk factors in microbial keratitis	19
2.5	Preparing fortified topical antibiotics	24
3.1	Definitions and nomenclature of dilute-solution viscosity quantities	34
3.2	Feldman buffer for ophthalmic preparations	39
3.3	Composition of each formulation	40
3.4	Scheme for preparing dilutions of vancomycin to be used in Mueller-Hinton broth (MHB) dilution susceptibility tests	42
4.1	Weight data and calculated loss on drying of chitosan	48
4.2	Dilute-solution viscosity data and calculated parameters for chitosan solution at $25.0 \pm 0.1^\circ\text{C}$ (Run No.1)	49
4.3	Experimental results and calculated degrees of deacetylation (DD)	50
4.4	Preparation of 0.1% and 0.3% chitosan solutions by using Feldman buffer for ophthalmic preparations	52
4.5	pH Stability of 0.1% and 0.3% chitosan solutions	53
4.6	Viscosity stability of 0.1% chitosan solution	54
4.7	Viscosity stability of 0.3% chitosan solution	56

Table	Page
4.8	61
Compatibility and stability studies of vancomycin eye drops at 50 mg/ml in Tears Naturale II TM , 0.9% sodium chloride and the 0.1% and 0.3% chitosan solutions	
4.9	63
UV spectrophotometric scan of standard vancomycin in Tears Naturale II TM and Tears Naturale II TM	
4.10	64
UV spectrophotometric scan of standard vancomycin in 0.1% chitosan and 0.1% chitosan	
4.11	65
UV spectrophotometric scan of standard vancomycin in 0.9% sodium chloride and 0.9% sodium chloride	
4.12	66
Calibration of standard vancomycin in Tears Naturale II TM	
4.13	67
Calibration of standard vancomycin in 0.1 % chitosan	
4.14	68
Calibration of standard vancomycin in 0.3 % chitosan	
4.15	69
Calibration of standard vancomycin in 0.9 % sodium chloride	
4.16	70
Interday assay validation of standard vancomycin in each vehicle	
4.17	71
Intraday assay validation of standard vancomycin in each vehicle	
4.18	71
Recovery validation of standard vancomycin in each vehicle	
4.19	72
Lower limit of standard vancomycin in each vehicle	
4.20	73
The pH of vancomycin eye drops at 50 mg/ml in each vehicle	
4.21	76
Percentage of the labeled amounts of vancomycin eye drops at 50 mg/ml in each vehicle	
4.22	78
Minimum inhibitory concentrations of vancomycin at 50 mg/ml in Tears Naturale II TM , sodium chloride and chitosan solutions stored at different temperatures over time	

Table		Page
4.23	Comparison of the kinematic viscosity, ν , of various solvents and vancomycin at 50 mg/ml in solutions at 25°C	79
4.24	Control test of vancomycin	81
4.25	Concentration of vancomycin at 50 mg/ml in Tears Naturale II™ in tear film	82
4.26	Concentration of vancomycin at 50 mg/ml in 0.1% chitosan in tear film	83
4.27	Concentration of vancomycin at 50 mg/ml in 0.3% chitosan in tear film	84
4.28	Concentration of vancomycin at 50 mg/ml in normal saline solution (NSS) in tear film	85
4.29	Areas under the curves of vancomycin at 50 mg/ml 0→30 minutes for the range of vehicles used	86
4.30	Areas under the curves of vancomycin at 50 mg/ml 0→120 minutes for the range of vehicles used	87

LIST OF FIGURES

Figure		Page
1.1	Chemical structure of chitosan	2
1.2	Chemical structure of vancomycin	4
2.1	Anatomy of the eye	6
2.2	Nasolachrymal drainage system	9
3.1	Heating vacuum desiccator	31
3.2	Viscometers commonly used in dilute-solution viscometry	32
3.3	Reduced and inherent viscosity-concentration plots (data from this research)	35
3.4	The Schott-Gerate AVS Automatic Viscosity Measuring System used in dilute-solution viscometry showing, from left to right, the Ubbelohde viscometer, main control unit, and constant temperature water bath	36
3.5	Preparation of chitosan solution	38
3.6	Osmomat 030	38
3.7	Growth (not clear) and no growth (clear) of bacteria	43
3.8	The animals were maintained under conventional, standardized conditions in a single cage with free access to pelleted food and drinking water	44
3.9	Preparation of rabbits for applying eye drops	45
3.10	Eye drops instilled into the rabbit's eye	45
3.11	Tear samples were obtained using 2.0 μ l of calibrated glass capillaries (microcaps Drummond™)	46
3.12	Fluorescent polarization immunoassay (TDx-FLx System Abbott, USA)	47

Figure		Page
4.1	Graph of reduced viscosity, η_{red} , and inherent viscosity, η_{inh} , against concentration (Run No. 1)	50
4.2	Variations in intrinsic viscosity, $[\eta]$, of the 0.1% w/v chitosan solution with autoclaving and storage time at different temperatures	55
4.3	Variations in intrinsic viscosity, $[\eta]$, of the 0.1% w/v chitosan solution with storage time only at different temperatures	55
4.4	Variations in intrinsic viscosity, $[\eta]$, of the 0.3% w/v chitosan solution with autoclaving and subsequent storage time at different temperatures	57
4.5	Variations in intrinsic viscosity, $[\eta]$, of the 0.3% w/v chitosan solution with storage time only at different temperatures	57
4.6	UV spectrophotometric scan (in graph) of standard vancomycin in Tears Naturalle II TM and Tears Naturalle II TM	63
4.7	UV spectrophotometric scan (in graph) of standard vancomycin in 0.1% chitosan and 0.1% chitosan	64
4.8	UV spectrophotometric scan (in graph) of standard vancomycin in 0.9% sodium chloride and 0.9% sodium chloride	65
4.9	Calibration curve of standard vancomycin in Tears Naturalle II TM no. 1	66
4.10	Calibration curve of standard vancomycin in 0.1% chitosan no. 1	67
4.11	Calibration curve of standard vancomycin in 0.3% chitosan no.1	68
4.12	Calibration curve of standard vancomycin in 0.9% sodium chloride no. 1	69

Figure		Page
4.13	The pH graph of vancomycin eye drops at 50 mg/ml in each vehicle	74
4.14	Percentage graph of the labeled amounts of vancomycin eye drops at 50 mg/ml in each vehicle	77
4.15	Comparison of the kinematic viscosity, ν , of various solvents and vancomycin at 50 mg/ml in solutions at 25°C by bar graph	80
4.16	Concentration-time profiles of vancomycin at 50 mg/ml in Tears Naturale II in tear film (6 samples)	82
4.17	Concentration-time profiles of vancomycin at 50 mg/ml in 0.1% chitosan solution in tear film (6 samples)	83
4.18	Concentration-time profiles of vancomycin at 50 mg/ml in 0.3% chitosan solution in tear film (6 samples)	84
4.19	Concentration-time profiles of vancomycin at 50 mg/ml in 0.9% sodium chloride solution in tear film (6 samples)	85
4.20	Area under the curve of vancomycin at 50 mg/ml in the various vehicles used	88

LIST OF ABBREVIATIONS

DD	=	degree of deacetylation
AUC _{0,t}	=	area under concentration-time curve from administration and extrapolation to time
%CV	=	percent coefficient of variation
°C	=	degree celsius
°F	=	degree fahrenheit
Conc	=	concentration
Sol	=	solution
HPLC	=	high performance liquid chromatography
h	=	hour
ht	=	height
kg	=	kilogram
L	=	liter
LLQ	=	lower limit of quantitation
mg	=	milligram
µg	=	microgram
dl	=	deciliter
µl	=	microliter
mM	=	millimolar
min	=	minute
ml	=	milliliter
mm	=	millimeter
nm	=	nanometer
PK	=	pharmacokinetics
SD	=	standard deviation

ลิขสิทธิ์มหาวิทยาลัยเชียงใหม่
 Copyright © by Chiang Mai University
 All rights reserved