

TABLE OF CONTENTS

	Page
ACKNOWLEDGEMENT	iii
ABSTRACT (THAI)	v
ABSTRACT (ENGLISH)	vii
LIST OF TABLES	xii
LIST OF FIGURES	xiii
ABBREVIATIONS	xv
CHAPTER 1 INTRODUCTION	1
1.1 Background and Hypothesis	1
1.2 Objectives	3
1.3 Literature Review	3
CHAPTER 2 MATERIALS AND METHODS	20
2.1 Materials	20
2.2 Methods	23
2.2.1 Plant Material and Identification	23
2.2.2 Extraction and Preparations	23
2.2.3 Identifications of HCE with TLC	26
2.2.4 Identification of HCE with HPLC	30
2.2.5 Method validations	31
2.2.6 Development of HCE Tablet Formulations	31

ลิขสิทธิ์มหาวิทยาลัยเชียงใหม่
 Copyright © by Chiang Mai University
 All rights reserved

2.2.7	Quality controls	37
2.2.8	Stability Tests	40
CHAPTER 3 RESULTS AND DISCUSSION		42
3.1	Extraction and Preparations	42
3.1.1	HCE	42
3.1.2	HCE and Fraction for analysis	42
3.2	Identification of HCE	44
3.2.1	Identification of HCE with TLC	44
3.2.2	Identification of HCE with HPLC	48
3.3	Method Validation	49
3.3.1	Calibration curves	49
3.3.2	Precision and Accuracy	51
3.3.3	Limits of Detection (LOD) and Limits of Quantitative (LOQ)	53
3.4	Development of HCE Tablet Formulation	53
3.4.1	Calculation of HCE Tablet Formulation	53
3.4.2	Improvement of HCEP	54
3.4.3	HCE Tablet Formulations	59
3.5	Quality Controls	62
3.5.1	Quality controls of HCEP and bulk powder	62
3.5.2	Quality controls of HCE tablets	65
3.6	Stability tests of HCE	68
3.6.1	Stability test of HCEP and bulk powder	68
3.6.2	Stability of HCE tablets	69

ลิขสิทธิ์มหาวิทยาลัยเชียงใหม่

Copyright © by Chiang Mai University

All rights reserved

CHAPTER 4 CONCLUSIONS	73
REFERENCES	75
VITA	83



ลิขสิทธิ์มหาวิทยาลัยเชียงใหม่
Copyright© by Chiang Mai University
All rights reserved

LIST OF TABLES

Table	Page
1.1 Names of HC depended on origin	4
2.1 Ratios of HCE and absorbent	32
2.2 Development of HCE tablet formulations	34
2.3 HCE tablet formulations	36
3.1 Linear regression analysis of flavonoid standards	51
3.2 Recoveries of flavonoids from HCE over 3 days	52
3.3 Properties of HC granules/powder through adsorption	56
3.4 Hardness of HC powder/granules as tablets for 3 formulations	58
3.5 Hardness, disintegration times and friability of each formulation tablets	61
3.6 Quality controls of HCEP and bulk powder	62
3.7 Quality controls of HCE tablets by hydraulic press	66
3.8 Quality controls of HCE tablets by single stroke tableting machine	67
3.9 Moisture contents of HCEP and bulk powder	68
3.10 Quality controls of HCE tablets after stability test	70

LIST OF FIGURES

Figure	Page
1.1 <i>Houttuynia cordata</i> Thunb. plant	6
1.2 Structures of flavonoids compounds in <i>H. cordata</i> Thunb.; (a) quercetin and (b) rutin	8
2.1 Liquid-liquid partition of concentrated HCE as 4 fraction (F1-F4)	25
2.2 Mobile phase ratios for gradient HPLC	30
3.1 The extraction of HC	43
3.2 TLC chromatograms of HCE and fractions, in comparison with geraniol standard; (a) and (b) were not sprayed, (c) and (d) were sprayed, (a) and (c) were observed under visible light, while (b) and (d) were examined under UV light (365 nm).	45
3.3 TLC chromatograms of HCE and fractions, in comparison with rutin and quercetin standard; (a) and (b) were not sprayed, (c) and (d) were sprayed, (a) and (c) were observed under visible light while (b) and (d) were examined under UV-365 nm.	47
3.4 HCE chromatogram from HPLC-UV (350 nm) analysis for (a) rutin and (b) quercetin	48
3.5 HPLC chromatogram of (a) rutin and (b) quercetin standards on HPLC-UV (350 nm)	49


Figure	Page
3.6 Standard curve of rutin	50
3.7 Standard curve of Quercetin	50
3.8 HCE with absorbent as granules of (a) formulation 4, and powder of (b) formulation 6 and formulation 7 (c)	57
3.9 SEM of Prosolv [®]	63
3.10 SEM of Prosolv [®] :HCE as 1:2	63
3.11 SEM of Prosolv [®] :HCE as 1:8	64
3.12 HCE tablets by hydraulic press	66
3.13 HCE tablets by single stroke tableting machine	67
3.14 TLC chromatograms of HCE tablet extracts, compared with crude extract and standards. Chromatograms are visualized under UV-254 nm (A) and UV-365 nm (B). Lane 1: crude HCE, Lane 2: HCE tablet extract 0 th Mo at 45°C, Lane 3: HCE tablet extract 1 st Mo at 45°C, Lane 4: HCE tablet extract 2 nd Mo at 45°C, Lane 5: HCE tablet extract 3 rd Mo at 45°C, Lane 6: HCE tablet extract 3 rd Mo at RT, Lane 7: rutin standard, Lane 8: quercetin standard.	72

SYMBOLS AND ABBREVIATIONS

%LOD	Percentage of loss on drying
°C	Degree Celsius
µm	Micrometers
AHP	American herbal pharmacopoeia
AR	Analytical grade
CD	Cluster of differentiation
cm	centimeter
CMC	The chemistry, manufacturing and control
CSD	Colloidal silicon dioxide
DI	Deionized water
DNA	Deoxyribonucleic acid
DVS	Developing solvent
g	Gram
GC-MS	Gas chromatography-mass spectrophotometer
GPO	Government pharmaceutical organization
HC	<i>Houttuynia cordata</i> Thunb.
HCE	<i>Houttuynia cordata</i> Thunb. extract
HCEP	<i>Houttuynia cordata</i> Thunb. extract powder
HIV	Human immunodeficiency virus
HPLC	High performance liquid chromatography

HSV	Herpes simplex virus
IL	Interleukin
kg	Kilogram
kv	Kilovolt
LOD	Limits of detection
LOQ	Limits of quantitative
MCC	Microcrystalline cellulose
mAU	Milliabsorbance units
mg	Milligram
min	Minute
ml	Milliliter
mm	Millimeter
Mo	Month
N	Newton
nm	nanometers
R ²	correlation coefficient
R _f	Retard factor
RNA	Ribonucleic acid
rpm	rounds per a minute
RSD	Relative standard deviation
RT	Room temperature
s	second
s.d.	Standard deviation
SARS	Serve acute respiration syndromes

SEM	Scanning electron microscope
SMCC	Silicified microcrystalline cellulose
T	Ton
TLC	Thin layer chromatography
USP/NF	United state pharmacopeia/national formulary
UV	Ultraviolet
w/v	weight by volume



ลิขสิทธิ์มหาวิทยาลัยเชียงใหม่
Copyright© by Chiang Mai University
All rights reserved