

TABLE OF CONTENTS

	Page
ACKNOWLEDGEMENTS	iii
ABSTRACT (ENGLISH)	iv
ABSTRACT (THAI)	v
LIST OF TABLES	ix
LIST OF FIGURES	x
ABBREVIATIONS AND SYMBOLS	xv
CHAPTER 1 INTRODUCTION	1
1.1 Introduction	1
1.2 Tungsten oxide	3
1.2.1 Physical properties	3
1.2.2 Structure of Tungsten Oxide	3
1.2.2.1 General characteristics of WO_3 and transition metal oxides	3
1.2.2.2 Perovskite-like Structure	4
1.2.2.3 Rutile-like Structure	5
1.2.2.4 Tungsten oxide hydrates	7

1.2.3 Application of Tungsten Oxide	9
1.3 Hydrothermal/solvothermal method	10
1.4 Surfactant	20
1.5 Literature Review	25
1.6 Research Objectives	31
CHAPTER 2 EXPERIMENTAL PROCEDURE	32
2.1 Chemical reagents and equipments	32
2.1.1 Chemical reagents	32
2.1.2 Equipments	32
2.2 Synthesis methods	33
2.2.1 Synthesis of tungsten oxide by a hydrothermal method using ammonium metatungstate hydrate as a tungsten source	33
2.2.2 Synthesis of tungsten oxide by a hydrothermal method using sodium tungstate dihydrate as a tungsten source	35
2.3 Characterization	38
2.3.1 X-ray diffraction (XRD)	38
2.3.2 Fourier transform inferred (FTIR) spectroscopy and Raman spectroscopy	39
2.3.3 Field Emission Scanning Electron Microscopy (FE-SEM)	41

2.3.4 Transmission Electron Microscopy (TEM)	42
2.3.6 Photoluminescence Spectroscopy	43
2.3.7 UV–visible Spectroscopy	44
CHAPTER 3 RESULTS AND DISCUSSION	45
3.1 Result of the products synthesized by a hydrothermal method using ammonium metatungstate hydrate as a tungsten source	45
3.2 Result of the products synthesized by a hydrothermal method using sodium tungstate dihydrate as a tungsten source	58
3.2.1 Effect of acidity	58
3.2.2 Effect of reaction temperature	63
3.2.3 Effect of reaction time	67
CHAPTER 4 CONCLUSIONS	77
4.1 Synthesis of tungsten oxide by a hydrothermal method using ammonium metatungstate hydrate as a tungsten source	77
4.2 Synthesis of tungsten oxide by a hydrothermal method using sodium tungstate dihydrate as a tungsten source	78
REFERENCES	79
APPENDICES	83
APPENDIX A The Joint Committee for Powder Diffraction Standards (JCPDS)	84
APPENDIX B Material Safety Data Sheet	95
CURRICULUM VITAE	100

LIST OF TABLES

Table	Page
1.1 Surfactant classifications	23
2.1 Experimental conditions for synthesizing of tungsten oxide by a hydrothermal method using sodium tungstate dihydrate as a tungsten source with different volumes of 3M HCl, length of reaction times and reaction temperature.	37

LIST OF FIGURES


Figure	Page
1.1 (a) Unit cell for the perovskite lattice (Part I) and octahedral symmetries (Part II) in the perovskite structure; (b) One layer of the monoclinic WO_3 structure in the corner-sharing arrangement of octahedra (ReO_3 -type); (c) The monoclinic WO_3 structure (ReO_3 -type); (d) One layer of the monoclinic WO_3 structure in the edge-sharing arrangement of octahedra.	6
1.2 Formation of $\text{WO}_3 \cdot n\text{H}_2\text{O}$ from the neutral precursor $[\text{H}_2\text{WO}_4]^0$	8
1.3 (a) The six membered rings of (WO_6) octahedra sharing their corners in the xy plane or one layer hexagonal WO_3 ; (b) The hexagonal WO_3 structure showing the tunnels along the z axis; the relevant ring structures are three membered ring cases.	9
1.4 Hydrothermal technology in the 21st century.	14
1.5 Hydrothermal tree showing different branches of science and technology.	15
1.6 Pressure temperature map of materials processing techniques.	16
1.7 Difference in particle processing by hydrothermal and conventional technique	19
1.8 General purpose autoclave popularly used for hydrothermal treatment and hydrothermal synthesis.	20
1.9 Schematic illustration of the micelle monomers.	21

1.10	Schematic illustration of the commonly observed geometrical shapes of surfactant micelles in aqueous solution.	25
2.1	Schematic diagram used for preparing tungsten oxide by a hydrothermal method using ammonium metatungstate hydrate as a tungsten source with different volumes of 1M HCl	34
2.2	Schematic diagram used for preparing tungsten oxide by a hydrothermal method using sodium tungstate dihydrate as a tungsten source with different volumes of 3M HCl, length of reaction times and reaction temperature	36
2.3	X-ray diffractometer	38
2.4	Fourier transform infrared spectroscopy	39
2.5	Raman spectroscopy	40
2.6	Field Emission Scanning Electron Microscopy	41
2.7	Transmission Electron Microscope	42
2.8	Luminescence spectroscopy	43
2.9	UV-visible spectroscopy	44
3.1	XRD patterns of the products synthesized by the hydrothermal reaction at 200 °C for 24 h in the solutions containing 0.00, 2.50 ml, 5.00 ml, and 7.50 ml of 1M HCl.	45
3.2	SEM images at low and high magnifications of the product synthesized by the hydrothermal reaction at 200 °C for 24 h in the solution containing 5.00 ml of 1M HCl.	47

3.3	SEM images at low and high magnifications of the product synthesized by the hydrothermal reaction at 200 °C for 24 h in the solution containing 7.50 ml of 1M HCl.	48
3.4	FTIR spectra of (a) CTAB, and (b–d) the products synthesized by the hydrothermal reaction at 200 °C for 24 h in the solutions containing 2.50 ml, 5.00 ml, and 7.50 ml of 1M HCl, respectively.	49
3.5	Raman spectrum of o-WO ₃ microflowers, synthesized by the hydrothermal reaction at 200 °C for 24 h in the solution containing 7.50 ml 1M HCl.	50
3.6	TEM images of o-WO ₃ microflowers, synthesized by the hydrothermal reaction at 200 °C for 24 h in the solution containing 7.50 ml 1M HCl.	51
3.7	(a, b) High magnification TEM images, and (c, d) SAED patterns of o-WO ₃ microflowers of Figure 3.6 (b).	52
3.8	Schematic diagram for the formation of o-WO ₃ microflowers.	54
3.9	(a) UV-visible spectrum, and (b) the plot of $(\alpha h\nu)^2$ versus $h\nu$ of o-WO ₃ microflowers, synthesized by the hydrothermal reaction at 200 °C for 24 h in the solution containing 7.50 ml of 1M HCl.	55
3.10	PL spectrum of o-WO ₃ microflowers.	57
3.11	XRD patterns of the products synthesized by the hydrothermal reaction at 180 °C for 24 h in the respective solutions containing 4.00 ml, 4.50 ml, 5.00 ml, 5.50 ml, 6.00 ml, 6.50 ml and 7.00 ml of 3M HCl.	58
3.12	SEM image of the product synthesized by the hydrothermal reaction at 180 °C for 24 h in the solution containing 4.00 ml 3M HCl.	59
3.13	SEM image of the product synthesized by the hydrothermal reaction at 180 °C for 24 h in the solution containing 4.50 ml 3M HCl.	60

3.14 SEM image of the product synthesized by the hydrothermal reaction at 180 °C for 24 h in the solution containing 5.00 ml 3M HCl.	60
3.15 SEM image of the product synthesized by the hydrothermal reaction at 180 °C for 24 h in the solution containing 5.50 ml 3M HCl.	61
3.16 SEM image of the product synthesized by the hydrothermal reaction at 180 °C for 24 h in the solution containing 6.00 ml 3M HCl.	61
3.17 SEM image of the product synthesized by the hydrothermal reaction at 180 °C for 24 h in the solution containing 6.50 ml 3M HCl.	62
3.18 SEM image of the product synthesized by the hydrothermal reaction at 180 °C for 24 h in the solution containing 7.00 ml 3M HCl.	62
3.19 XRD patterns of the products synthesized by the hydrothermal reaction at 120, 140, 160, 180 and 200 °C for 12 h in the solution containing 5.00 ml 3M HCl.	63
3.20 SEM image of the product synthesized by the hydrothermal reaction at 120 °C for 12 h in the solution containing 5.00 ml 3M HCl.	64
3.21 SEM image of the product synthesized by the hydrothermal reaction at 140 °C for 12 h in the solution containing 5.00 ml 3M HCl.	65
3.22 SEM image of the product synthesized by the hydrothermal reaction at 160 °C for 12 h in the solution containing 5.00 ml 3M HCl.	65
3.23 SEM image of the product synthesized by the hydrothermal reaction at 180 °C for 12 h in the solution containing 5.00 ml 3M HCl.	66
3.24 SEM image of the product synthesized by the hydrothermal reaction at 200 °C for 12 h in the solution containing 5.00 ml 3M HCl.	66

3.25	XRD patterns of the products synthesized by the hydrothermal reaction at 200 °C for 0, 6, 12, 24 and 48 h in the solution containing 5.00 ml 3M HCl.	67
3.26	SEM image of the product synthesized by the hydrothermal reaction at 200 °C for 0 h in the solution containing 5.00 ml 3M HCl.	68
3.27	SEM image of the product synthesized by the hydrothermal reaction at 200 °C for 6 h in the solution containing 5.00 ml 3M HCl.	69
3.28	SEM image of the product synthesized by the hydrothermal reaction at 200 °C for 12 h in the solution containing 5.00 ml 3M HCl.	69
3.29	SEM image of the product synthesized by the hydrothermal reaction at 200 °C for 24 h in the solution containing 5.00 ml 3M HCl.	70
3.30	SEM image of the product synthesized by the hydrothermal reaction at 200 °C for 48 h in the solution containing 5.00 ml 3M HCl.	70
3.31	SEM image of the product synthesized by the hydrothermal reaction at 200 °C for 72 h in the solution containing 5.00 ml 3M HCl.	71
3.32	FTIR spectrum of product synthesized by the hydrothermal reaction at 200 °C for 48 h in the solution containing 5.00 ml 3M HCl.	72
3.33	Raman spectrum of product synthesized by the hydrothermal reaction at 200 °C for 48 h in the solution containing 5.00 ml 3M HCl.	73
3.34	(a) UV-visible spectrum, and (b) the plot of $(\alpha h\nu)^2$ versus $h\nu$ of products synthesized by the hydrothermal reaction at 200 °C for 24 h in the solution containing 7.50 ml of 1M HCl.	75

ABBREVIATIONS AND SYMBOLS

°C	=	Degree Celsius
eV	=	Electron Volt
g	=	Gram
h	=	Hour
ml	=	Milliliter
nm	=	Nanometer
μm	=	Micrometer
JCPDS	=	The Joint Committee for Powder Diffraction Standards
MW	=	Molecular Weight
FT-IR	=	Fourier-Transform Infrared Spectrometer
PL	=	Photoluminescence Spectrometer
SEM	=	Scanning Electron Microscope
TEM	=	Transmission Electron Microscope
XRD	=	X-ray Diffraction Spectrometer

ลิขสิทธิ์มหาวิทยาลัยเชียงใหม่
Copyright © by Chiang Mai University
All rights reserved