

REFERENCES

1. a) Gawley, R. E., Aub, J. (1996). "Principle of asymmetric synthesis". Tetrahedron Organic Chemistry Series Volume 14, 1st ed. Oxford, Tarrytown, N.Y.: Pergamon. b) Procter, G. (1996). "Asymmetric Synthesis". New York : Oxford University press., pp. 1-13.
2. Eliel, E. L. (1974). "Studies in asymmetric synthesis. Highly stereoselective reactions of organosulfur compounds". *Tetrahedron*, 30, 1503-1513.
3. a) Marckwald, W. (1904) "Ueber asymmetrische synthese". *Ber. Dtsch. Chem. Ges.* 37, 349-354. b) Marckwald, W. (1904) "Ueber asymmetrische synthese". *Ber. Dtsch. Chem. Ges.* 37, 1368-1370.
4. Evans, D. A., Ennis, M. D., Mathre, D. J. (1982). "Asymmetric alkylation reactions of chiral imide enolates. A practical approach to the enantioselective synthesis of .alpha.-substituted carboxylic acid derivatives". *J. Am. Chem. Soc.* 104, 1737-1739.
5. www.standardbase.hu/tech/SITechPolar.pdf (date accessed 24 May 2011)
6. Parker, D. (1981). "NMR determination of enantiomeric purity". *Chem. Rev.* 91, 1441-1457.
7. Aboul-Enein, H. Y., Ali, I. (2003). "Chiral Separations by Liquid Chromatography and Related Technologies". New York : M. Dekker, Inc.

8. Cabusas, M. E. Y. (1998). "Chiral separations on HPLC derivatized polysaccharide CSPs: Temperature, mobile phase and chiral recognition mechanism studies". Ph.D. Thesis. Virginia Polytechnic Institute and State University. pp. 10-12.
9. a) Seebach, D., Beck, K.A., and Heckel, A. (2001). "TADDOLs, Their Derivatives, and TADDOL Analogues: Versatile Chiral Auxiliaries". *Angew. Chem. Int., Ed.* 40, 92–138. b) Pellissier, H. (2008). "Use of TADDOLs and their derivatives in asymmetric Synthesis". *Tetrahedron*, 64, 10279–10317.
10. Qian, C.-T., Gao, F.-F., Sun, J. (2000). "First application of amino-TADDOL derivatives addition of diethylzinc to aryl aldehydes: effect of substituents on the nitrogen atom". *Tetrahedron: Asymmetry*, 11, 1733-1740.
11. Ueki, M., Matsumoto, Y., Jodry, J.J., Milkami, K. (2001). "Titanium 3,3'-modified-biphenolate complexes atropisomerically controlled by TADDOLs: novel chiral lewis acid catalysts for asymmetric methylation with an achiral methyl-titanium reagent". *Synlett*, 12, 1889-1892.
12. Adam, W., Alsters, P. L, Neumann, R., Möller, C.R., Seebach, D., Zhang, R. (2003). "Highly efficient catalytic asymmetric epoxidation of allylic alcohols by an oxovanadium-substituted polyoxometalate with a regenerative TADDOL-derived hydroperoxide". *Org. Lett.*, 5, 725-728.
13. Moteki, S.A., Chandra, K. L., Reddy, D. S., Takacs, J. M. (2006). "TADDOL-derives phosphites and phosphoramidites for efficient rhodium-catalyzed asymmetric hydroboration". *Org. Lett.*, 8, 3097-3100.

14. Vinogradov, M. G., Gorshkova, L. S., Chel'tsova, G. V., Pavlov, V. A., Razmanov, I. V., Ferapontov, V.A., Malyshev, O. R., Heise, G. L. (2003). "Enantioselective reduction of C=O and C=N bonds by TADDOL-containing aluminiumhydride reagents based on NaAlH₄ and AlH₃". *Russ. Chem. Bull., Int. Ed.*, 52, 471-479.
15. a) Weber, B., Seebach, D. (1992). "Enantiomerically Pure Tertiary Alcohols by TADDOL-Assisted Additions to Ketones-or How to Make a Grignard Reagent Enantioselective". *Angew. Chem., Int. Ed. Engl.*, 31, 84-86. b) Weber, B., Seebach, D. (1994). "Highly enantioselective addition of primary alkyl *Grignard* reagents to carbocyclic and heterocyclic arylketones in the presence of magnesium TADDOLate preparative and mechanistic aspects". *Tetrahedron*, 50, 6117-6128.
16. Marson, A., Freixa, Z., Kamer, P. C. J., van Leeuwen, P. W. N. M. (2007). "Chiral calix[4]arene-based diphosphites as ligands in the asymmetric hydrogenation of prochiral olefins". *Eur. J. Inorg. Chem.*, 4587-4591.
17. Aoki, M., Seebach, D. (2001). "Preparation of TADDOH, a hydroperoxide from TADDOL, and use in highly enantioface- and enantiomer-differentiating oxidations". *Helv. Chem. Acta.*, 84, 187-207.
18. Du, H., Zhao, D., Ding, K. (2004). "Enantioselective catalysis of the hetero-Diels–Alder reaction between Brassard's Diene and aldehydes by hydrogen-bonding activation: A one-step synthesis of (S)-(+)-Dihydrokawain". *Chem. Eur. J.*, 10, 5946-5970.
19. Mengel, A., Reiser, D. (1999). "Around and beyond Cram's rule". *Chem. Rev.*, 99, 1191-1223.

20. Mukaiyama, T., Soai, K., Sato, T., Shimizu, H., Suzuki, K. (1979). "Enantioface-differentiating (asymmetric) addition of alkyllithium and dialkylmagnesium to aldehydes by using (2S,2'S)-2-hydroxymethyl-1-[(1-alkylpyrrolidin-2-yl)methyl]pyrrolidines as chiral ligands". *J. Am. Chem. Soc.*, 101, 1455-1460.
21. a) Jongkol, R., Choommongkol, R., Tarnchompoo, B., Nimmanpipug, P., Meepowpan, P. (2009). "Syntheses of methylenolactocin and nephrosterinic acid via diastereoselective acylation and chemoselective reduction-lactonization". *Tetrahedron*, 65, 6382-6389. b) Jongkol, R., (2007) "Syntheses of Tetrahydro-4'-Carbomethoxy-5'-Alkyl-2'-Furanone-3'-Spiro-11-9,10-Dihydro-9,10-Ethanoanthracenes as Precursors for Preparation of α -Methylene- γ -Alkyl- β -Carboxylic- γ -Butyrolactones via Acylation and Tandem Reduction-Ring Closure Reactions", M.S. Thesis, Chiang Mai University.
22. Xianming, H., Kellogg, R. M. (1996). "Asymmetric reduction and Meerwein-Porndorf-Verley reaction of prochiral aromatic ketones in the presence of optically pure 1-aryl-2,2-dimethylpropane-1,3-diols". *Recl. Trav. Chim. Pays-Bas*, 115, 410-417.