

TABLE OF CONTENTS

	Page
ACKNOWLEDGEMENTS	iii
ABSTRACT (in English)	iv
ABSTRACT (in Thai)	vii
LIST OF TABLES	xii
LIST OF FIGURES	xiii
ABBREVIATIONS	xiv
CHAPTER I INTRODUCTION	1
CHAPTER II LITERATURE REVIEW	6
CHAPTER III OBJECTIVE	12
CHAPTER IV MATERIALS AND METHODS	13
4.1 Cell culture	13
4.2 B cell stimulation	14
4.3 Isolation of RNA	14
4.4 RT-PCR	15
4.4.1 Synthesis of first strand cDNA for PCR amplification	15
4.4.2 PCR reaction	16
4.4.3 One step RT-PCR	17
4.5 Agarose gel electrophoresis	18
4.6 Analysis of the band intensity	18

4.7 Staining of CD138 and Blimp-1 protein on Raji and LP-1 cells	19
4.7.1 Titration of anti-CD138 and conjugated antibody for staining	19
4.7.2 Determination of the CD138 expression in PWM stimulated Raji cells	19
4.7.3 Determination of different permeabilization agents for intra cellular protein staining	19
4.7.4 Titration of anti-CD 138 and conjugated antibody for staining	20
4.7.5 Determination of Blimp-1 expression in PWM stimulated Raji cells	20
4.8 Detection of antibody production by ELISA	21
4.8.1 Titration of goat IgG-IgM concentration coating to the plate	21
4.8.2 Detection of antibody production in cell culture supernatants by ELISA	21
CHAPTER V RESULTS	23
5.1 Determination of sandwich ELISA optimal conditions used for the detection of antibody in cell supernatants	23
5.2 Detection of antibody production in cell culture supernatants by ELISA	26
5.3 Staining of CD138 in PWN stimulate Raji cells	30
5.4 Determination the optimal condition used to staining of Blimp-1 protein	32
5.5 Quality control in determination of Blimp-1 expression by using GAPDH gene control	36

5.6 Determination of Blimp-1 mRNA isoform expression in PWM stimulated Raji B cells	38
5.7 Determination of Blimp-1 protein expression in PWM stimulated Raji cells	41
CHAPTER VI DISCUSSION	44
CHAPTER VII SUMMARY	50
REFERENCES	53
APPENDIX	59
CURRICULUM VITAE	76

LIST OF TABLES

Table	Page
1 Block titration of anti human IgG+IgM (H+L) coated plate	24
2 Titration of peroxidase conjugated anti human IgG-IgM.	25
3 Antibody standard in the concentration of 50, 25, 12.5, 6.25 and 3.12 were determined by using ELISA	26
4 Antibody secretion from PWM stimulated Raji B cells at different time points	29
5 Kinetic expression of the full length and $\Delta 7$ Blimp-1 mRNA isoforms in PWN stimulated Raji cell	40


LIST OF FIGURES

Figure	Page
1 Blimp-1 controls plasma cell differentiation	2
2 Model for the role of Blimp-1 in repressing B cell fate and Inducing plasma cell fate	7
3 Murine Blimp-1 mRNA and the human Blimp-1, <i>prdm1</i> gene	8
4 RT-PCR analysis of mouse Blimp-1 mRNA	9
5 Antibody secretion from PWM stimulated Raji B cells at Different time points	28
6 Expression of CD138 on Raji cells by flow cytometry	30
7 Comparison of permeability reagents used for intracellular protein staining	32
8 Comparison of FITC conjugated anti goat IgM used for Blimp-1 staining	35
9 Kinetic expression of the full length and $\Delta 7$ Blimp-1 mRNA isoforms in PWM stimulated Raji B cells	37
10 Kinetic expression of the full length and $\Delta 7$ Blimp-1 mRNA isoforms in PWM stimulated Raji B cells	39
11 Blimp-1 protein expression in PWM stimulated Raji cells	41

ABBREVIATIONS

Ab	Antibody
Ag	Antigen
Blimp-1	B lymphocyte-induced maturation protein 1
BSA	bovine serum albumin
cDNA	complementary deoxyribonucleic acid
CD	cluster of differentiation
DNA	deoxyribonucleic acid
EDTA	ethylene diamine tetra acetic acid
FITC	fluorescein isothiocyanate
FBS	fetal bovine serum
GAPDH	glyceraldehydes-3-phosphate dehydrogenase
HRP	horseradish peroxidase enzyme
Ig	immunoglobulin
IL	interleukin
mRNA	messenger ribonucleic acid
OD	optical density
PBS	phosphate buffer saline
PCR	polymerase chain reaction
PWM	pokeweed mitogen
RNA	ribonucleic acid

rpm	revolution per minute
RT	room temperature
RT-PCR	reverse transcription polymerase chain reaction
SD	standard deviation
Std	standard



ลิขสิทธิ์มหาวิทยาลัยเชียงใหม่
Copyright© by Chiang Mai University
All rights reserved