TABLE OF CONTENTS

TABLE OF CONTENTS			
		Page	
	ACKNOWLEDGEMENTS	iii	
	ABSTRACT (IN ENGLISH)	v	
	ABSTRACT (IN THAI)	vii	
	TABLE OF CONTENTS	ix	
	LIST OF TABLES	xiii	
	LIST OF FIGURES	XV	
	ABBREVIATIONS AND SYMBOLS	XX	
	CHAPTER I INTRODUCTION	1	
	1. STATEMENT AND SIGNIFICANCE OF THE PROBLEM	1	
	2. LITERATURE REVIEWS	3	
	3. OBJECTIVE	8	
	4. USEFULNESS OF THE STUDY	8	
	CHAPTER II MATERIALS AND METHODS	10	
	1. MATERIALS	10	
	1.1 For metaphase chromosome preparation	10	
	1.2 For light and scanning electron microscopic studies	10	
	1.3 For polytene chromosome preparation	10 - 15	
	1.4 For molecular study	10	

TABLE OF CONTENTS (continued)

	Page
2. METHODS	11
2.1 Field collection and establishment of isoline colonies	11
2.2 Mosquito rearing	14
2.2.1 Wild-caught An. campestris-like Form E	14
2.2.2 Insectarium condition	14
2.2.3 Egg laying	15
2.2.4 Rearing of larva, pupa and adult	15
2.2.5 Searching for a suitable blood-feeding condition	16
2.2.6 Ability of free mating in a 30 cm cubed cage	17
2.2.7 Male ability to mate artificially	18
2.2.8 Searching for a suitable oviposition-condition	18
2.2.9 Establishment of the laboratory-colony strain	21
2.2.10 Rearing condition for chromosome preparation	21
2.3 Metaphase karyotype preparation	21
2.4 Crossing study	22
2.5 DNA extraction, amplification and sequencing	22
2.6 DNA sequence and phylogenetic analysis	23
2.7 Morphological studies of eggs, larvae, pupae and adults	23
under light and/or scanning electron microscopy	

TABLE OF CONTENTS (continued)

	Page
2.8 Salivary gland polytene chromosomes preparation	24
2.9 Malaria susceptibility test	25
2.10 Statistical analysis	26
CHAPTER III RESULTS	27
1. Mosquito rearing	27
1.1 Observation on the life duration, hatchability, pupation	27
and emergence	
1.2 Feeding ability	29
1.3 Ability of free mating in a 30 cm cubed cage	29
1.4 Ability of male in artificial mating	31
1.5 Ability of eggs deposition	32
1.6 Rearing condition for chromosome preparation	32
2. Metaphase karyotype characters	33
3. Crossing study	36
4. DNA sequences and phylogenetic analysis	42
5. Morphological studies of eggs, larvae, pupae and adults	50
under light microscope	
6. Morphological studies of eggs under scanning electron	50 ersi
microscopy	

TABLE OF CONTENTS (continued)

	Page
7. Salivary gland polytene chromosome investigation	54
8. Malaria susceptibility test	58
CHAPTER IV DISCUSSION	65
CHAPTER V CONCLUSION	75
REFERENCES	77
APPENDIX	96
CURRICULUM VITAE	114

ลิ<mark>ปสิทธิ์มหาวิทยาลัยเชียงใหม่</mark> Copyright[©] by Chiang Mai University All rights reserved

90 910 OF TABLES

Гable		Page
1	Life duration of An. campestris-like Form E from an iso-female	28
	line laboratory-raised colony.	
2	Direct feeding ability on white rat in a 30 x 30 x 30 cm cage and	30
	artificial feeding ability on bovine heparined blood from different	
	ages of An. campestris-like Form E.	
3	Artificial mating ability of An. campestris-like Form E males.	31
4	Crossing experiments among the 12 isolines of An. campestris-	37
	like Forms B, E and F.	
5	Locations, isoline colonies and karyotypic forms of An.	43
	campestris-like and their GenBank accession numbers.	
6	Average genetic distance within and between the An.campestris-	49
	like Form B, E and F for the ITS2, COI and COII regions.	
7	Oocysts and sporozoites detected from An. cracens and An.	61
	campestris-like Form E and F 8 and 14 days post infection with	
	<i>P. falciparum</i> (gametocyte density/200 wbc = 40).	

LIST OF TABLES (continued)

Table

8

Page

Oocysts and sporozoites detected from *An. cracens* and *An.* 62 *campestris*-like Form B, E and F 8 and 14 days post infection with *P. vivax* (gametocyte density/200 wbc = 32, 15 and 20 in experiment I, II and III, respectively).

ลิ<mark>ปสิทธิ์มหาวิทยาลัยเชียงใหม่</mark> Copyright[©] by Chiang Mai University All rights reserved

LIST OF FIGURES

XV

Figure

2

3

Map of Thailand showing distribution of An. campestris-like12Form B, E and F.12

Flow chart for rapid systematic procedure.

Mosquito equipment used in this study. (A) Ovipot derived from a screen-topped plastic cup with the top covered by a black plastic-sheet (left) for single gravid wild-caught adult females to lay eggs, (middle) for 20 gravid adult females to lay eggs, and (right) top view of the plastic cup showing massive egg-batches after 12-hours-oviposition of the 20 gravid adult females; (B) eggs placed in a white plastic tray and exposed to a 40 watt light. (C) White plastic rearing tray containing 2,000 ml of filtered natural water, 15 stems of garden grass and 80 larvae, with the top of the tray covered with a transparent plastic-sheet. (D) Left: adult rearing cage partially covered with a wet towel and a transparent plastic-sheet with plastic container for holding pupae, and two bottles with cotton wicks, one containing 10% sucrose solution and the other 5% multivitamin syrup solution. Right: adult males being kept in a screen-topped paper cup with 13

19

Page

xvi

Figure

4

Page

5% multivitamin syrup solution, and the top covered with a translucent plastic bag to maintain humidity; (E) Artificial feeding system with thin paraffin-membrane at the bottom tip of the feeding apparatus containing bovine heparinized-blood that is in close contact with 50 fasted adult female *An. campestris*-like Form E inside a screen-topped paper cup.

Metaphase karyotypes of *An. campestris*-like Form B, E and F (a-e). Form B: a Kamphaeng Phet strain, showing X_2 , Y_2 chromosomes; Form E: b Chumphon strain, showing X_2 , Y_5 chromosome; c Sa Kaeo strain, showing X_3 , Y_5 chromosome; Form F: d Udon Thani strain, showing X_2 , Y_6 . chromosome; c Sa Kaeo strain, showing X_3 , Y_5 chromosome; Form F: d Udon Thani strain, showing X_2 , Y_6 chromosome; e showing homozygous X_2 , X_2 chromosomes; f Diagrams of representative metaphase karyotypes of *An. campestris*-like Form F.

A phylogenetic trees of *An. campestris*-like Form B, E and F (*CAM*) based on molecular analysis of ITS2. The tree was generated by neighbor-joining analysis. Numbers on the nodes indicate probabilities based on 1,000 bootstrap replicates. A

35

Figure

Page

47

48

52

probability of more than 50% is shown. Branch lengths are proportional to genetic distance (scale bar).

- A phylogenetic trees of *An. campestris*-like Form B, E and F (*CAM*) based on molecular analysis of COI. The tree was generated by neighbor-joining analysis. Numbers on the nodes indicate probabilities based on 1,000 bootstrap replicates. A probability of more than 50% is shown. Branch lengths are proportional to genetic distance (scale bar).
- A phylogenetic tree of *An. campestris*-like Form B, E and F (*CAM*) based on molecular analysis of COII. The tree was generated by neighbor-joining analysis. Numbers on the node indicate probabilities based on 1,000 bootstrap replicates. A probability of more than 50% is shown. Branch lengths are proportional to genetic distance (scale bar).
- Egg surface topography of *An. campestris*-like under scanning electron microscope (SEM); whole eggs: (A) dorsal aspect, anterior end (a), posterior end (p) (x 230). (B) Lateral aspect, anterior end (a), posterior end (p) (x 250). (C) Anterior end, showing irregularly jagged tubercles on the deck and four large,

xvii

xviii

Figure

11

Page

53

56

rosette-shaped tubercles (2,200). (D) A higher magnification of the irregularly jagged tubercles on the deck (x 7,000).

- Egg surface topography of *An. campestris*-like under scanning electron microscope (SEM): (A) the inner surface of the frill (fr), showing its sclerotized, ridge-like texture with picket-like ribs (x 3,500). (B) The outer surface of the frill (fr), showing its smooth surface and parallel brick-like texture along its entire length (x 3,500). (C) The exochorionic sculptering in the middle of the egg, showing more or less regularly hexagonal reticulum with membrane like-wall enclosing many small, less prominent tubercles. (D) The anterior end, showing the micropylar orifice surrounded by a smooth collar with an irregular outer margin (x 3,300).
- Complete set of salivary gland polytene chromosomes of *An.* 55
 campestris-like Form B (C = Centromere, L = Left arms, R = Right arms, X = X-chromosome).
 - Complete set of salivary gland polytene chromosomes of An. campestris-like Form E (C = Centromere, L = Left arms, R =
 - Right arms, X = X-chromosome).

Page Figure 12 Complete set of salivary gland polytene chromosomes of An. 57 campestris-like Form F (C = Centromere, L = Left arms, R = Right arms, X = X-chromosome). 13 Oocysts of P. vivax recovered from midgut of (A) An. cracens 63 and (B) An. campestris-like Form E (Chiang Mai) on day 8 after infection. 14 Free flow regular spindle-shaped sporozoites of P. vivax from 64 the squashed salivary glands (small arrow) (A) An. cracens and (B) An. campestris-like Form E (Chiang Mai) on day 14 after infection.

ลิขสิทธิ์มหาวิทยาลัยเชียงใหม่ Copyright[©] by Chiang Mai University All rights reserved

ABBREVIATIONS AND SYMBOLS

%	percentage
	less than
>	more than
°C	degree Celsius
μΙ	microliter
μΜ	micromolar
bp	base pair
С	centromere
CAM	An. campestrs-like Forms
cm	centimeter
cm ²	square centimeter
COI	cytochome c oxidase subunit I
COII	cytochome c oxidase subunit II
DNA	deoxyribonucleic acid
dNTPs	deoxynucleotide triphosphates
e.g.	exempli gratia
et al	and others
etc.	etcetera
fr Chang	frill
hnts r	houre served
i.e.	id est

ITS2	second internal transcribed spacer
W SIE BO	international unit
IU/Kg	international unit per kilogram
KV	voltage
	left arms
mg/kg	milligram per kilogram
ml	milliliter
min	minute
mtDNA	mitochondrial DNA
No.	number
p	probability
РВ	phosphate buffer
PCR	polymerase chain reaction
рН	potential of hydrogen
R	right arms
rDNA	ribosomal DNA
RH UNI	relative humidity
SEM	scanning electron microscope
s.l.	sensu lato
spp.	species
S.S.	sensu stricto
wbc Chiang	white blood cell