TABLE OF CONTENTS



3.3.1 Sele	ecting of herbal plants		16
3.3.2 Prej	paration of herbal powders 16		
3.3.3 Ver	ification of herbal plants and powders	17	
3.3.	3.1 Identity verification	17	
3.3.4 Ana	lysis of ash content and herbal		
pow	der compositions		18
3.3.	4.1 Ash content		18
3.3.	4.2 Starch Content	18	
3.3.	4.3 Volatile Oil Content	3	19
3.3.	4.4 Fiber Content	19	
3.3.5 Exa	mination of fundamental pharmaceutical		
prop	perties of herbal powder	502	19
3.3.	5.1 Flow properties	19	0
3.3.	5.2 Compressibility	21	
3.3.6 Stat	istical analysis of the relationship between	4	
hert	al powder composition and its fundamental	0/	
pha	maceutical properties		21
3.3.7 Dev	elopment of herbal tablet formulation	21	
3.3.	7.1 Tablet production process selection	21	
-3.3.	7.2 The pressure transmission ratio (R)	23	
3.3.	7.3 Disintegration test	23	
3.3.8 Sca	e-up phase	23	
3.3.9 Qua	lity control of finished products		23
87805113.3	9.1 Tablet appearance	23	
3.3.	9.2 Tablet dimensions		24
Copyright [©] 3.3.	9.3 Weight variation	24	sitv
3.3.	9.4 Strength of tablets	24	/
A I I 3 .3.	9.5 Disintegration test	24	
0			

Chapter 4 Results and Discussion 25

4.1	Examination of herbal powders	25

4.1.1 Macroscopic characteristics

ix

26

	4.1.2 Microscopic characteristics	30
4.2	Herbal powder composition	49
4.3	4.3 Flow properties	
4.4	4.4 Compaction property	
4.5	Formulation of herbal powders	60
	4.5.1 Development Phase I-Effect of binders	
	at a compression force of 15,000 N	64
6	4.5.2 Development Phase II-Effect of binders	
	at a compression force of 20,000 N	68
6	4.5.3 Development Phase III-Effect of adding	
	MCC at a concentration of 20%	71
-3524	4.5.4 Development Phase IV-Effect of adding	3024
Z S	MCC at a concentration of 20% in combination	
106	with a compression force of 20,000 N	74
	4.5.5 Development Phase V-Effect of adding binder	4
	at a higher concentration	78
	4.5.6 Development Phase VI-Effect of adding binder	~
	at a higher concentration in combination with	
	a compression force of 20,000 N	81
	4.5.7 Development Phase VII-Effect of adding MCC	
	at a concentration of 50%	84
	4.5.8 Development Phase VIII-Effect of adding MCC	
	at a concentration of 50% together with	
	a compression force of 20,000 N	87
	4.5.9 Development Phase IX-Effect of adding	
Copyright	superdisintegrant ang Mai Uni	vers ⁹⁰ v
4.6	Quality control of herbal tablet formulation	92
AII r	ignts reser	
Chapter 5 Co	onclusions	103
5.1	Part used	103
	5.1.1 Fruit	103
	5.1.2 Underground part	103

5.1.3 Leaf	104
5.1.4 Stem	104
5.2 Herbal powder composition	104
5.2.1 Plants with large amounts of fiber	104
5.2.2 Plants with large amounts of starch	104
5.2.3 Plants with large amounts of volatile oil	104
rences 105	

References

Appendix	108
1. Starch content (Glucoamylase method)	109
2. Crude fiber content	
Curriculum Vitae	113
	1967
AI UNIVER	

ลิขสิทธิ์มหาวิทยาลัยเชียงใหม่ **Copyright[©]** by Chiang Mai University All rights reserved

LIST OF TABLES

Table	ે અમદામંછ	Page
2.1	The common quantity of the basic component that is found in	
	each part of plant	4
3.1	Flow properties and corresponding angle of repose	19
3.2	Scale of flowability	20
3.3	Production process selection by evaluating the flow	
	and compression properties	22
4.1	Quantitative analysis and basic composition of herbal powder	50
4.2	Flow properties of herbal powder	54
4.3	Compressibility ratio of herbal powder and herbal granules	
	using different binders	56
4.4	Hardness of the herbal tablets compressed with a 15,000 N	
	compression force	59
4.5	The maximum lower and upper punch force and the pressure	
	transmission ratio (R) of herbal tablet formulations using	
	different binders	62
4.6	Hardness, weight and disintegration time of herbal tablets	
	prepared using different binders compressed at $15,000 \text{ N} (n = 6)$	66
4.7	Hardness, weight and disintegration time of herbal tablets	
	prepared using different binders compressed at 20,000 N ($n = 6$)	69
	Hardness, weight and disintegration time of herbal tablets	sitv
	prepared using different binders and 20% MCC compressed at	/
	15,000 N (n = 6)	e 72 0
4.9	Hardness, weight and disintegration time of herbal tablets	
	prepared using different binders and 20% MCC compressed at	
	20,000 N (n = 6)	76

xii

4.10	Hardness, weight and disintegration time of herbal tablets	
	prepared using different binders by double wet granulation	
	with 20% MCC and compressed at 15,000 N ($n = 6$)	79
4.11	Hardness, weight and disintegration time of herbal tablets	
	prepared using different binders by double wet granulation	
	with 20% MCC and compressed at 20,000 N ($n = 6$)	82
4.12	Hardness, weight and disintegration time of herbal tablets	
	prepared using different binders with 50% MCC compressed at	
	15,000 N (n = 6)	85
4.13	Hardness, weight and disintegration time of herbal tablets	
	prepared using different binders with 50% MCC compressed at	
-30	20,000 N (n = 6)	88
4.14	Hardness, weight and disintegration time of C. angustifolia tablets	
100	using different binders with 47% MCC and 3% CS compressed	
	at 20,000 N (n = 6)	91
4.15	Quality control of herbal tablet formulations	93
4.16	Optimal herbal tablet formulations that passed quality control	
	evaluation	97
	MAR TERS	
	UNIVE	

ลิขสิทธิ์มหาวิทยาลัยเชียงใหม่ Copyright[©] by Chiang Mai University All rights reserved

LIST OF FIGURES

Figure

Figure	ે ગેમદાસંધ	Page
4.1	Macroscopic characteristics of crude drugs from herbal fruit 26	
4.2	Macroscopic characteristics of crude drugs from herbal underground par	t 27
4.3	Macroscopic characteristics of crude drugs from herbal leaf 28	
4.4	Macroscopic characteristics of crude drugs from herbal stem 29	
4.5	Powdered drug of the pericarp of <i>P. emblica</i> 30	
4.6	Powdered drug of the fruit of <i>S. trilobatum</i> 31	
4.7	Powdered drug of the fruit of <i>T. chebula</i>	32
4.8	Powdered drug of the fruit of P. nigrum	33
4.9	Powdered drug of the fruit of <i>M. citrifolia</i>	34
4.10	Powdered drug of the rhizome of Z. officinale	35
4.11	Powdered drug of the rhizome of <i>C. longa</i>	36
4.12	Powdered drug of the rhizome of A. calamus	37
4.13	Powdered drug of the root of <i>E. longifolia</i>	38
4.14	Powdered drug of the root of G. glabra	39
4.15	Powdered drug of the leaf of C. angustifolia	40
4.16	Powdered drug of the leaf of A. paniculata	41
4.17	Powdered drug of the leaf of C. asiatica	42
4.18	Powdered drug of the leaf of <i>P. indica</i>	43
4.19	Powdered drug of the leaf of C. hystrix	44
4.20	Powdered drug of the stem of <i>T. crispa</i> 45	sitv
4.21	Powdered drug of the stem bark of C. verum	46
A 4.22	Powdered drug of the stem of <i>C. sappan</i> 47	
4.23	Powdered drug of the stem of D. scandens	48
4.24	Kawakita plot of Z. officinale Powder	53
4.25	Microscopic photograph of herbal powder and granule of	
	C. asiatica leaf	53

4.26 Examplary compression profile of <i>T. crispa</i> powder obtained	
from instrumented tableting machine	61
4.27 Appearance of the most appropriate herbal tablet formulations	
from fruit powder) 9
4.28 Appearance of the most appropriate herbal tablet formulations	
from underground part powder 100	
4.29 Appearance of the most appropriate herbal tablet formulations	
from leaf powder 10	01
4.30 Appearance of the most appropriate herbal tablet formulations	
from stem powder 10	02
REAL AND	

ลิขสิทธิ์มหาวิทยาลัยเชียงใหม่ Copyright[©] by Chiang Mai University All rights reserved

ABBREVIATIONS

a	Compactibility value
AR	Analytical reagent
CS	Croscarmellose sodium
DT	Disintegration time
g	Gram
GT	10% gelatin solution
kg	Kilogram
M	Molar
MCC	Microcrystalline cellulose
mg	Milligram
min	Minute
ml	Milliliter
mm	Millimeter
msec	Millisecond
Ν	Newton
ND	Not Detected
РР	10% polyvinylpyrrolidone
R	Pressure transmission ratio
S	Second
SDETITIO	Standard deviation
SG	5% starch paste + 5% gelatin solution
SP	10% starch paste
USP/NF	United state pharmacopeia/national formulary
w/w	weight by weight e serve o
	a AR CS DT g GT kg M MCC mg min MCC mg min m1 mm msec N ND PP R s SD SC SD SC SP USP/NF w/w

Scientific name of herbal plants

A. calamus	Acorus calamus Linn.
A. paniculata	Andrographis paniculata Wall. Ex Nees.

xvi

C. angustifolia	Cassia angustifolia Vahl.
C. asiatica	Centella asiatica (L.) Urban.
C. hystrix	Citrus hystrix DC.
C. longa	Curcuma longa Linn.
C. sappan	Caesalpinia sappan Linn.
C. verum	Cinnamomum verum J.S.
D. scandens	Derris scandens Benth.
E. longifolia	Eurycoma longifolia Jack.
G. glabra	Glycyrrhiza glabra Linn.
M. citrifolia	Morinda citrifolia L.
P. emblica	Phyllanthus emblica Linn.
P. indica	Pluchea indica (L.) Less.
P. nigrum	Piper nigrum Linn.
S. trilobatum	Solanum trilobatum Linn.
T. chebula	Terminalia chebula Retz.
T. crispa	Tinospora crispa Miers.
Z. officinale	Zingiber officinale Rosc.
N'à L	
	41 UNIVER?

ลิขสิทธิ์มหาวิทยาลัยเชียงใหม่ **Copyright[©]** by Chiang Mai University All rights reserved