TABLE OF CONTENTS

| | Page |
|---|------|
| ACKNOWLEDGEMENT | iii |
| ENGLISH ABSTRACT | iv |
| THAI ABSTRACT | vi |
| LIST OF TABLES | xi |
| LIST OF FIGURES | xii |
| ABBREVIATIONS | xiii |
| CHAPTER 1 INTRODUCTION | |
| 1.1 Herbal medicines | 1 |
| 1.2 Inflammation | 1 |
| 1.3 Drugs for the treatment of inflammation | 5 |
| 1.4 Historical background of Stahlianthus involucratus (King ex | 8 |
| Baker) Craib ex Loes. | |
| 1.5 Hypothesis | 10 |
| 1.6 Purpose of the study | 10 |
| CHAPTER 2 MATERIALS AND METHODS | |
| 2.1 Preparation of the extract | 11 |
| 2.2 Test substance administration | 11 |
| 2.3 Experimental animals | 12 |
| 2.4 Experimental models | 12 |
| 2.4.1 Anti-inflammatory activity | |
| A. Ethyl phenylpropiolate-induced ear edema | 12 |
| B. Carrageenin-induced hind paw edema | 13 |
| C. Arachidonic acid-induced hind paw edema | 15 |
| D. Cotton pellet-induced granuloma formation | 16 |
| 2.4.2 Analgesic activity | |
| A. Acetic acid-induced writhing response | 19 |
| B Tail-flick test | 20 |

| 2.4.3 Antipyretic activity | | | | | |
|---|------|--|--|--|--|
| A. Yeast-induced hyperthermia | 22 | | | | |
| 2.5 Statistical analysis | | | | | |
| 2.6 Drugs and chemicals | 23 | | | | |
| CHAPTER 3 RESULTS | | | | | |
| 3.1 Gas chromatography/mass spectrometry analysis of the extract | 24 | | | | |
| 3.2 Anti-inflammatory activity | | | | | |
| 3.2.1 Effects of <i>S. involucratus</i> extract and diclofenac on ethyl | 24 | | | | |
| phenylpropiolate-induced ear edema | | | | | |
| 3.2.2 Effects of S. involucratus extract and diclofenac on | 24 | | | | |
| carrageenin-induced hind paw edema | | | | | |
| 3.2.3 Effects of S. involucratus extract, diclofenac, and | 25 | | | | |
| prednisolone on arachidonic acid-induced hind paw | | | | | |
| edema | | | | | |
| 3.2.4 Effects of S. involucratus extract, diclofenac, and | 25 | | | | |
| prednisolone on transudative and granuloma components, | | | | | |
| and ulcerogenic effect on cotton pellet-induced | | | | | |
| granuloma formation | | | | | |
| 3.2.5 Effects of S. involucratus extract, diclofenae, and | 25 | | | | |
| prednisolone on body weight and thymus weight of cotton | | | | | |
| pellet-induced granuloma formation | | | | | |
| 3.2.6 Effects of S. involucratus extract, diclofenac, and | 25 | | | | |
| prednisolone on alkaline phosphatase activity of cotton | | | | | |
| pellet-induced granuloma formation | | | | | |
| 3.3 Analgesic activity | | | | | |
| 3.3.1 Effects of S. involucratus extract and diclofenac on acetic | 26 | | | | |
| acid-induced writhing response | / | | | | |
| 3.3.2 Effects of S. involucratus extract, diclofenac, and codeine | e 26 | | | | |
| on tail-flick test | | | | | |

| 3.4 Antipy | retic activity | | | | | | | |
|--|----------------|----------|--------|--------------|---------|-----|----|--|
| 3.4.1 | Antipyretic | activity | of S. | involucratus | extract | and | 26 | |
| diclofenac on yeast-induced hyperthermia | | | | | | | | |
| CHAPTER 4 DISC | CUSSION AN | ID CONC | CLUSIO | N | | | 37 | |
| REFERENCES | 9/19 | | | | | | 41 | |
| VITA | | | | | | | 47 | |
| | MAI | UI UI | | ERSI | | | | |

ลิขสิทธิ์มหาวิทยาลัยเชียงใหม่ Copyright[©] by Chiang Mai University All rights reserved

LIST OF TABLES

| Tab | ole | Page |
|------|---|------|
| 1 | Chemical components of S. involucratus ethanolic extract as analyzed | 27 |
| | by gas chromatography/mass spectrometry analysis | |
| 2 | Percentage of edema inhibition of S. involucratus extract and | 28 |
| | diclofenac on ethyl phenylpropiolate-induced ear edema | |
| 3 | Percentage of edema inhibition of S. involucratus extract and | 29 |
| | diclofenac on carrageenin-induced hind paw edema | |
| 4 | Percentage of edema inhibition of S. involucratus extract, diclofenac, | 30 |
| | and prednisolone on arachidonic acid-induced hind paw edema | 1 |
| 5 | Transudative weight and percentage of granuloma inhibition of | 31 |
| | S. involucratus extract, diclofenac, and prednisolone on | |
| | cotton pellet-induced granuloma formation | |
| 6 | The changes of body weight and dry thymus weight of S. involucratus | 32 |
| | extract, diclofenac, and prednisolone on cotton pellet-induced | |
| | granuloma formation | |
| 7 | Alkaline phosphatase activity of S. involucratus extract, diclofenac, | 33 |
| | and prednisolone on cotton pellet-induced granuloma formation | |
| 8 | Percentage of writhing response inhibition of S. involucratus extract | 34 |
| | and diclofenac on acetic acid-induced writhing response | |
| 9 | Effects of S. involucratus extract, diclofenac, and codeine on tail-flick | 35 |
| | test S11K99ng19ag113gla | |
| 10 | Antipyretic activity of S. involucratus extract and diclofenac on yeast- | 36 |
| Copy | induced hyperthermia Chiang Mai Univer | sity |
| AII | rights reserv | |
| | | |

LIST OF FIGURES

| Figu | ire | Page | | | |
|------|--|------|--|--|--|
| 1 | Interactions between cell-adhesion molecules during the initial binding | | | | |
| | and tight binding of leukocytes to activated endothelial cells | | | | |
| 2 | Steps in the biosynthesis of eicosanoids and sites of drug action | | | | |
| 3 | Stahlianthus involucratus (King ex Baker) Craib ex Loes. | 9 | | | |
| 4 | Method of ethyl phenylpropiolate-induced ear edema in rats | 13 | | | |
| 5 | Method of carrageenin-induced hind paw edema in rats | 15 | | | |
| 6 | Method of arachidonic acid-induced hind paw edema in rats | 16 | | | |
| 7 | Method of cotton pellet-induced granuloma formation in rats | 19 | | | |
| 8 | 8 Method of acetic acid-induced writhing response in mice 2 | | | | |
| 9 | 9 Method of tail-flick test in rats | | | | |
| 10 | Method of yeast-induced hyperthermia in rats | 22 | | | |
| | TO THE WAT UNIVERSITATION OF THE PROPERTY OF T | | | | |

ลิขสิทธิ์มหาวิทยาลัยเชียงใหม่ Copyright[©] by Chiang Mai University All rights reserved

ABBREVIATIONS

| | m 31 | 212 | meter |
|------------------------|------|-----------------|--------------------|
| | cm | = | centimeter |
| | mm | 自心 | millimeter |
| | μm | | micrometer |
| | L | 景(| liter |
| | dL | | deciliter |
| | mL | THILL S | milliliter |
| 20% | μL | = (h | microliter |
| | g | = 2 | gram |
| | mg | | milligram |
| | kg | 4 | kilogram |
| | d | = | day |
| | h | (₹)_{- | hour |
| | min | 1 2 | minute |
| \\ \(\C \) | sec | | second |
| | °C | = | degree Celsius |
| | M | ≠N | molar |
| | W | = | watt |
| | U | = | unit |
| | BW | 18 | body weight |
| | eV | = | electronvolt |
| Copyright [©] | w/w | hia | weight by weight |
| | i.d. | = | inner diameter |
| Allrıg | enz | | enzyme S e r v e o |
| | | | |