

TABLE OF CONTENTS

	Page
ACKNOWLEDGEMENT	iii
ABSTRACT (THAI)	iv
ABSTRACT (ENGLISH)	vi
LIST OF TABLES	x
LIST OF FIGURES	xiii
ABBREVIATIONS AND SYMBOLS	xv
CHAPTER I INTRODUCTION	1
1.1 Statement and significance of the problem	1
1.2 Objectives	3
CHAPTER II THEORY AND LITERATURE REVIEWS	4
2.1 Skin	4
2.2 Free radicals	10
2.3 Antioxidant	12
2.4 Carotenoids	14
2.5 <i>Momordica cochinchinensis</i> (Lour.) Spreng	21
2.6 Nanostructured lipid carriers	27
CHAPTER III MATERIALS AND METHOD	31
3.1 Apparatus and instruments	31
3.2 Chemicals	32
3.3 Plants material and methods	32
3.4 Determination of antioxidant activities of aril oil of <i>Momordica cochinchinensis</i> (Lour.) Spreng	33
3.5 Determination of total carotenoids content of aril oil of <i>Momordica cochinchinensis</i> (Lour.) Spreng	34
3.6 Reversed phase - High Performance Liquid Chromatography (RP-HPLC) analysis	35
3.7 Preparation of nanostructured lipid carriers (NLC)	37

TABLE OF CONTENTS (continued)

	Page
3.8 Formulation cream containing aril oil and cream containing aril oil-loaded NLC	40
3.9 Chemical stability studies β -carotene of cream containing aril oil and cream containing aril oil-loaded NLC	41
3.10 Primary skin irritation test	42
3.11 Wrinkle reducing capacity test	44
CHAPTER IV RESULTS AND DISCUSSION	47
4.1 Morphological characteristics of <i>Momordica cochinchinensis</i> (Lour.) Spreng	47
4.2 The aril oil of <i>Momordica cochinchinensis</i> (Lour.) Spreng	47
4.3 Determination of antioxidant activities of aril oil of <i>Momordica cochinchinensis</i> (Lour.) Spreng	48
4.4 Determination of total carotenoids content of aril oil of <i>Momordica cochinchinensis</i> (Lour.) Spreng	51
4.5 Reversed phase - High Performance Liquid Chromatography (RP-HPLC) analysis	51
4.6 Preparation of nanostructured lipid carriers (NLC)	55
4.7 Formulation cream containing aril oil and cream containing aril oil loaded NLC	62
4.8 Chemical stability studies β -carotene of cream containing aril oil and cream containing aril oil-loaded NLC	65
4.9 Primary skin irritation test	66
4.10 Wrinkle reducing capacity test	67
CHAPTER V CONCLUSIONS	72
REFERENCES	74
APPENDICES	80
Appendix A	81
Appendix B	86
Appendix C	88
CURRICULUM VITAE	89

LIST OF TABLES

Table	Page
1 Role of free radical in various diseases	11
2 Some select antioxidants and their mechanism of action	13
3 Names of <i>Momordica cochinchinensis</i> (Lour.) Spreng in different languages	23
4 Comparison of reported concentrations of carotenes in aril oil of <i>Momordica cochinchinensis</i> (Lour.) Spreng (mg/g of edible portion)	25
5 Fatty acid composition of aril of <i>Momordica cochinchinensis</i> (Lour.) Spreng	25
6 Nutrient compositions of aril oil of <i>Momordica cochinchinensis</i> (Lour.) Spreng	26
7 Composition of the NLC formulation (% , w/w)	38
8 Composition of the aril oil loaded-NLC formulation	39
9 The formula preparation of oil in water cream base	40
10 The percent yield of aril oil of <i>Momordica cochinchinensis</i> (Lour.) Spreng	48
11 The absorbance and total carotenoids concentration of aril oil of <i>Momordica cochinchinensis</i> (Lour.) Spreng	51
12 The precision of HPLC method for standard peak of β -carotene	53
13 The accuracy of HPLC method for standard peak of β -carotene	54
14 The concentration of β -carotene of aril oil of <i>Momordica cochinchinensis</i> (Lour.) Spreng	55
15 Mean particle size, polydispersity index (PI) and zeta potential (ZP) values of the NLC formulation prepared at different homogenization pressure and homogenization cycle 5 cycles. (n = 3)	56
16 Mean particle size, polydispersity index (PI) and zeta potential (ZP) values of the NLC formulation prepared at different homogenization cycle and homogenization pressure 1000 bar (n = 3)	56

LIST OF TABLES (continued)

Table	Page
17 Mean particle size, polydispersity index (PI) and zeta potential (ZP) values of the three different concentrations of aril oil loaded NLC formulation prepared at homogenization pressure 1000 bar and homogenization cycle 5 times (n = 3)	57
18 Mean particle size, polydispersity index (PI) and zeta potential (ZP) Values of NLC formulation before and after 90 days of storage at 4, 25 and 45°C (n = 3)	60
19 Mean particle size, polydispersity index (PI) and zeta potential (ZP) values of 1% aril oil-loaded NLC formulation (1% A-NLC) before and after 90 days of storage at 4, 25 and 45°C (n = 3)	61
20 Mean particle size, polydispersity index (PI) and zeta potential (ZP) values of 3% aril oil-loaded NLC formulation (3% A-NLC) before and after 90 days of storage at 4, 25 and 45°C (n = 3)	61
21 Mean particle size, polydispersity index (PI) and zeta potential (ZP) values of 5 % aril oil-loaded NLC formulation (5% A-NLC) before and after 90 days of storage at 4, 25 and 45°C (n = 3)	61
22 Physical properties of the cream base before and after heating-cooling cycle and after 90 days of storage at 4, 25 and 45°C	62
23 Physical properties of the cream containing aril oil before and after heating-cooling cycle and after 90 days of storage at 4, 25 and 45°C	63
24 Physical properties of the cream containing NLC before and after heating-cooling cycle and after 90 days of storage at 4, 25 and 45°C	64
25 Physical properties of the cream containing aril oil-loaded NLC before and after heating-cooling cycle and after 90 days of storage at 4, 25 and 45°C	64
26 Percent remaining data of β -carotene. Samples were cream containing aril oil at 4, 25, 45°C and light after 90 days of storage	65
27 Percent remaining data of β -carotene. Samples were cream containing aril oil loaded NLC at 4, 25, 45°C and light after 90 days of storage	66

LIST OF TABLES (continued)

Table	Page
28 The value of primary irritation index (PII) and type of skin irritation in volunteers	67
29 The wrinkle reducing property on the R_a , R_z , surface and volume parameters of five treatment sites	69

LIST OF FIGURES

Figure	Page
1 Basic structure of the skin	5
2 The five layers of the epidermis	6
3 Later stages of carotenoids biosynthesis and possible transformations of carotenoids	16
4 Chemical structure of lycopene	17
5 Chemical structure of β -carotene	17
6 Possible scheme for the degradation of carotenoids	20
7 The characterization of <i>Momordica cochinchinensis</i> (Lour.) Spreng	21
8 The three types of NLC: imperfect type concentrations of oil (left), amorphous type (middle), multiple type (right)	
10 The characterization of <i>Momordica cochinchinensis</i> (Lour.) Spreng fruit	29
9 Profile of test sites (N = untreated area; A = cream containing NLC area; B = cream containing aril oil loaded NLC area; C = cream base area; D = cream containing aril oil area)	46
10 The characterization of <i>Momordica cochinchinensis</i> (Lour.) Spreng fruit	47
11 The dried aril with seeds of <i>Momordica cochinchinensis</i> (Lour.) Spreng	48
12 The aril oil of <i>Momordica cochinchinensis</i> (Lour.) Spreng	48
13 Linear regression plot of antioxidant activity against concentration of aril oil by DPPH assay	49
14 Linear regression plot of antioxidant activity against concentration of (\pm)- α -tocopherol standard by DPPH assay	50
15 Linear regression plot absorbance against concentration of $\text{FeSO}_4 \cdot 7\text{H}_2\text{O}$ standard by FRAP assay	50
16 Chromatogram of β -carotene standard using RP-HPLC system	52
17 Standard curve of β -carotene standard	52
18 Chromatogram of aril oil of <i>Momordica cochinchinensis</i> (Lour.) Spreng	55
19 The NLC formulations contained three concentration of aril oil (a = 1% of aril oil; b = 3% of aril oil; c = 5% of aril oil)	58

LIST OF FIGURES (continued)

Figure	Page
20 TEM photography of NLC	59
21 TEM photography of aril oil-loaded NLC	59
22 The physical appearance of four formulations (A = cream base; B = cream containing aril oil; C = cream containing NLC; D = cream containing aril oil-loaded NLC)	64
23 % Efficiency value of R_a parameter of five test sites (A = cream containing NLC area; B = cream containing aril oil-loaded NLC area; N = untreated area; C = cream base area; D = cream containing aril oil area)	69
24 % Efficiency value of R_z parameter of five test sites (A = cream containing NLC area; B = cream containing aril oil-loaded NLC area; N = untreated area; C = cream base area; D = cream containing aril oil area)	70
25 % Efficiency value of volume parameter of five test sites (A = cream containing NLC area; B = cream containing aril oil-loaded NLC area; N = untreated area; C = cream base area; D = cream containing aril oil area)	70
26 % Efficiency value of surface parameter of five test sites (A = cream containing NLC area; B = cream containing aril oil-loaded NLC area; N = untreated area; C = cream base area; D = cream containing aril oil area)	71

ABBREVIATIONS AND SYMBOLS

A-NLC	=	Aril oil-loaded nanostructured lipid carriers
%	=	Percent
%CV	=	Percentage of coefficient of variation
°C	=	Degree celcius
cm	=	Centimeter
DPPH	=	1,1-Diphenyl-2-picrylhydrazyl
EC ₁	=	Concentration of antioxidant having a ferric-TPTZ reducing ability equivalent to that of 1 mM FeSO ₄ .7H ₂ O
FRAP	=	Ferric reducing antioxidant power
g	=	Gram
HPH	=	High pressure homogenizer
hr	=	Hour
IC ₅₀	=	Half maximal inhibitory concentration
mg	=	Milligram
min	=	Minute
mV	=	Millivolt
mL	=	Milliliter
nm	=	Nanometer
No.	=	Number
NLC	=	Nanostructured lipid carriers
O ₂	=	Oxygen
O ₂ ⁻	=	Superoxide
Pas	=	Pascal
pH	=	Power of hydronium ion
PI	=	Polydisperity index
R ²	=	R-squared
R _a	=	Average roughness

RP-HPLC	=	Reversed phase - high performance liquid chromatography
R _z	=	Arithmetic average roughness
SD	=	Standard deviation
SLN	=	Solid lipid nanoparticles
TEM	=	Transmission electron microscope
TPTZ	=	2,4,6-Tris(2-pyridyl)-1,3,5-triazine
UV-Vis	=	Ultraviolet-visible
v/v	=	Volume by volume
w/w	=	Weight by weight
ZP	=	Zeta potential
µg	=	Microgram
µl	=	Microliter
µM	=	Micromolar
µm	=	Micrometer
λ	=	Wavelength