

REFERENCES

- Abbott, W.S. 1925. A method of computing the effectiveness of an insecticide. *J. Econ. Entomol.*, 18: 265-267.
- Aemprapa, S. 2007. Entomopathogenic fungi screening against fruit fly in Thailand. *KMITL Sci. Tech. J.*, 7: 122-126.
- Ahn, Y.J., Park, S.J., Lee, S.G., Shin, S.C., Choi, D.H. 2000. Cordycepin: selective growth inhibitor derived from liquid culture of *Cordyceps militaris* against *Clostridium* spp.. *J. Agric. Food Chem.*, 48: 2744-2748.
- Akhir, S.M., Abd-Aziz, S., Salleh, M.M., Rahman, R.A., Illia, R.M., Hassan, M.A. 2009. Medium optimization of chitinase enzyme production from shrimp waste using *Bacillus licheniformis* TH-1 by response surface methods. *Biotechnology.*, 8(1): 120-125.
- Al-Ahmadi, K.J., Yazdi, M.T., Najafi, M.F., Shahverdi, A.R. 2008. Optimization of medium and cultivation conditions for chitinase production by the newly isolated *Aeromonas* sp. *Biotechnology.*, 7(2): 266-272.
- Ali, S., Wu, J., Huang, Z., Ren, S.X. 2010. Production and regulation of extracellular chitinase from the entomopathogenic fungus *Isaria fumosorosea*. *Biocontrol Sci. Tech.*, 20(7): 723-738.
- Allwood, A.J. 1997. Regional management of fruit flies in the Pacific (53 pp.). Project document. RAS/97/331.

- Allwood, A.J., Chinajariyawong, A., Drew, R.A.I., Hamacek, E.L., Hancock, D.L., Hangsawad, C., Jipanin, J.C., Jirasurat, M., Kong Krong, C., Kritsaneepaiboon, S., Leong, C.T.S., Vijaysegaran, S. 1999. Host plants for fruit flies (Diptera; Tephritidae) in South-East Asia. *Raffles Bulletin of Zoology*, **7**: 1-92.
- Aluja, M., Celedonio-Hurtado, H., Liedo, P., Cabrera, M., Castillo, F., Guillen, J., Rios, E. 1996. Seasonal population fluctuations and ecological implications for management of *Anastrepha* fruit flies. *Department of Agriculture Bulletin*, **640**: 1-44.
- Alves, S.B., Lopes, R.B. Controle Microbiano de Pragas na América Latina: Avanços e Desafios. FEALQ, Piracicaba, São Paulo., 2008.
- Alves, S.B., Rossi, L.S., Lopes, R.B., Tamai, M.A., Pereira, R.M. 2002. *Beauveria bassiana* yeast phase on agar medium and its pathogenicity against *Diatraea saccharalis* (Lepidoptera: Crambidae) and *Tetranychus urticae* (Acari: Tetranychidae). *J Invertebr Pathol.*, **81**: 70-77.
- Anand, R., Prasad, B., Tiwary, B.N. 2009. Relative susceptibility of *Spodoptera litura* pupae to selected entomopathogenic fungi. *Biol Control.*, **54**: 85-92.
- Angel-Sahagun, C.A., Lezama-Gutierrez, R., Molina-Ochoa, J., Galindo- Velasco, E., Lopez-Edwards, M., Rebolledo-Dominguez, O., Cruz-Vazquez, C., Reyes-Velazquez, W.P., Skoda, S.R., Foster, J.E. 2005. Susceptibility of biological stages of the horn fly, *Haematobia irritans*, to entomopathogenic fungi (Hyphomycetes), *J. Insect. Sci.*, **5**: 50.
- Aregger, E. 1992. Conidia production of the fungus *Beauveria brongniartii* on barley and quality evaluation during storage at 2°C. *J Invertebr Pathol.*, **59**: 2-10.

- Armstrong, J.W., Jang, E.B. 1997. An overview of present and future fruit fly research in Hawaii and the U.S. mainland (pp. 30-42). In: Allwood, A.J., Drew, R.A.I. (Eds.), *Management of fruit flies in the Pacific*. ACIAR Proceedings No. 76. *Aust. Centre for Int. Ag. Res.* 267pp. (Conference Proceedings)
- Asaff, A., Cerda-Garcia-Rojas, C., de la Torre, M. 2005. Isolation of dipicolinic acid as an insecticidal toxin from *Paecilomyces fumosoroseus*. *Appl. Microbiol. Biotechnol.*, 68: 542-547.
- Aung, O.M. Biodiversity and molecular phylogeny of entomopathogenic fungi in Chiang Mai province, Thailand. Ph.D Thesis, King Mongkut's University of Technology Ladkrabang (KMUTL), 2008.
- Azevedo, J.L., Maccheroni, J.W., Pereira, J.O., Araújo, W.L. 2000. Endophytic microorganisms: A review on insect control and recent advances on tropical plants. *Eletron. J. Biotechnol.*, 3(1): 1-4.
- Batta, Y.A. 2003. Production and testing of novel formulations of the entomopathogenic fungus *Metarhizium anisopliae* (Metschnikoff) Sorokin (Deuteromycotina: Hyphomycetes). *Crop Prot.*, 22: 415-422.
- Benjamin, R.K., Blackwell, M., Chapela, I.H., Humber, R.A., Jones, K.G., Klepzig, K.A., Lichtwardt, R.W., Malloch, D., Noda, H., Roeper, R.A., Spatafora, J.W., Weir, A. 2004. Insects and other arthropod-associated fungi. In: Mueller, G.M., Bills, G.F., Mercedes, S. (Eds.), *Biodiversity of fungi inventory and monitoring methods* (pp. 395-434). Foster: Elsevier academic press.

- Benz, S. 1987. Environment. In: Fuxa, J.R., Tanada, Y. (Eds.), *Epizootiology of insect diseases* (pp. 177-214), New York: Wiley.
- Bernardine, D.M., Pinto, J.S., do, N., Rineiro, B., da Silva, C.L. 2006. Efeito dos fungos entomopatogênicos *Metarhizium anisopliae* e *Beauveria bassiana* sobre o desenvolvimento da *Musca domestica* L. (Diptera: Muscidae) em laboratório. *Arq. Inst. Biol.*, 73: 127-129.
- Bernstein, I.L., Bernstein, J.A., Miller, M., Tierzieva, S., Bernstein, D.I., Lummus, Z., Selgrade, M.K., Doerfler, D.L., Seligy, V.L. 1999. Immune responses in farm workers after exposure to *Bacillus thuringiensis* pesticides. *Environ Health Perspect.*, 107: 575-582.
- Blackwell, M., Vega, E. Insect-fungal associations: ecology and evolution (352 pp.). Oxford: Oxford University Press. 2005.
- Bokonon-Ganta, A.H., McQuate, G.T., Messing, R.H. 2007. Natural establishment of a parasitoid complex on *Bactrocera latifrons* (Diptera: Tephritidae) in Hawaii. *Biol Control.*, 42: 365-373.
- Boudreaux, H. B. 1963. Biological aspects of some phytophagous mites. *Annu. Rev. Entomol.*, 8: 137-154.
- Bugeme, D.M., Knapp, M., Boga, H.I., Wanjoya, A.K., Maniania, N.K. 2009. Influence of temperature on virulence of fungal isolates of *Metarhizium anisopliae* and *Beauveria bassiana* to the two-spotted spider mite *Tetranychus urticae*. *Mycopathologia.*, 167: 221-227.
- Burtet, M.J.G., Silva, M.E., Diehl-Fleig, E. 1997. Produção de conídios e micélio seco de *Beauveria bassiana* (Bals.) Vuill. para controle de formigas cortadeiras. *Congresso Bras Entomol.*, 16 : 101.

- Butt, T.M., Goettel, M.S. 2000. Bioassays of entomogenous fungi. In : Navon, A., Ascher, K.R.S. (Eds.), *Bioassays of entomopathogenic microbes and nematodes* (pp. 141-195). Oxon, UK : CABI publishing.
- Cagle, L. R. 1949. Life history of the twospotted spider mite. *Virginia Agr. Exp. Stn. Tech. Bull.*, 113: 1-31.
- Calderon, A., Fraga, M., Carreras, B. 1995. Production of *Beauveria bassiana* by solid-state fermentation. *Rev Protec Veg.*, 10: 269- 273.
- Calo, L., Fornelli, F., Ramires, R., Nenna, S., Tursi, A., Caiaffa, M.F., Macchia, L. 2004. Cytotoxic effects of the mycotoxin beauvericin to human cell lines of myeloid origin. *Pharmacol. Res.*, 49: 73-77.
- Carey, J., D'Amico, R., Sutton, D.A., Rinaldi, M.G. 2003. *Paecilomyces lilacinus* vaginitis in an immunocompetent patient. *Emerg Infect Dis.*, 9(9): 1155-1158.
- Castillo, M., Moya, P., Hernandez, E., Primo-Yufera, E. 2000. Susceptibility of *Ceratitis capitata* Wiedemann (Diptera: Tephritidae) to entomopathogenic fungi and their extracts. *Biol. Control.*, 19: 274-282.
- Chandler, D. Fungal control of insect and mite pests. School of life science. Warwick.UK. 2009. Online available: <http://www2.warwick.ac.uk/fac/sci/>
- Chandler, D., Davidson, G., Jacobson, R.J. 2005. Laboratory and glasshouse evaluation of entomopathogenic fungi against the two-spotted spider mite, *Tetranychus urticae* (Acari: Tetranychidae), on tomato, *Lycopersicon esculentum*. *Biocontrol Sci. Techn.*, 15(1): 37-54.
- Chandler, D., Davidson, G., Pell, J.K., Ball, B.V., Shaw, K., Sunderland, K.D. 2000. Fungal biocontrol in Acari. *Biological Sci. Techn.*, 10: 357-384.

- Charnley, A.K. 1989. Mechanisms of fungal pathogenesis in insects. In: Whipps, J.M., Lumsden, R.D. (Eds.), *Biotechnology for improving plant growth*. Cambridge, Cambridge University Press. 5: 5-125.
- Charnley, A.K. 2003. Fungal pathogens of insects: cuticle degrading enzymes and toxins. *Adv. Bot. Res.*, 40: 241-321.
- Charnley, A.K., Collins, S.A. 2007. Entomopathogenic fungi and their role in pest control. In: Kubicek, C.P., Druzhinina, I.S. (Eds.), *The Mycota IV: Environmental and Microbial Relationships (2nd edition)* (pp. 159-187). Berlin: Springer-Verlag.
- Christenson, L.D., Foote, R.H. 1960. Biology of fruit flies. *Ann. Rev. Entomol.*, 5: 171-192.
- Chutima, R., Dell, B., Vessabutr, S., Bussaban, B., Lumyong, S. 2010. Endophytic fungi from *Pecteilis susannae* (L.) Rafin (Orchidaceae), a threatened terrestrial orchid in Thailand. *Mycorrhiza*, 21(3): 221-229.
- Clarkson, J.M., Charnley, A.K. 1996. New insights into the mechanisms of fungal pathogenesis in insects. *Trends Microbiol.*, 4: 197-203.
- Claydon, N., Grove, J. 1982. Insecticidal secondary metabolic products from the entomogenous fungus *Verticillium lecanii*. *J Invertebr Pathol.*, 40: 413-418.
- Claydon, N., Grove, J.F., Pople, M. 1979. Insecticidal secondary metabolic products from the entomogenous fungus *Fusarium larvarum*. *J Invertebr Pathol.*, 33: 364-367.
- Cordon, T.C., Schwatz, J.H. 1962. The fungus *Beauveria tenella*. *Science.*, 138: 1265-1266.

- Cranshaw, W.S., Sclar, D.C. Spider mites. Colorado State University Extension. 2006. See: <http://www.ext.colostate.edu/pubs/insect/05507.html>.
- Cruz, B.P.B., Abreu, O.C., Oliveira, A.D., Chiba, S. 1993. Crescimento de *Metarhizium anisopliae* (Metsch.) Sorokin em meios de cultura naturais líquidos. *O. Biológico.*, 49: 111-116.
- Cunningham, K., Manson, W., Spring, F., Hutchinson, S. 1995. Cordycepin, a metabolic product isolated from cultures of *Cordyceps militaris*. *Nature.*, 166: 949.
- Dalla Santa, H.S., Dalla Santa, O.R., Brand, D., Vandenberghe, L.P.S., Soccol, C.R. 2005. Spore production of *Beauveria bassiana* from agro-industrial residues. *Braz Arch Biol Biotech.*, 48: 51-60.
- Dancer, B.N., Chantawannakul, P. 1997. The proteases of American foulbrood scales. *J Invertebr Pathol.*, 70: 79-87.
- de la Rosa, W., Lopez, F.L., Liedo, P. 2002. *Beauveria bassiana* as a pathogen of the Mexican fruit fly (Diptera: Tephritidae) under laboratory conditions, *J. Econ. Entomol.*, 95: 36-43.
- de la Torre, M., Cardenas-Cota, H.M. 1996. Production of *Paecilomyces fumosoroseus* conidia in submerged culture. *Entomophaga.*, 41: 443-453.
- de Moraes, K.C., Schrank, A., Vainstein, M.H. 2003. Regulation of extracellular chitinases and proteases in the entomopathogen and acaricide *Metarhizium anisopliae*. *Curr. Microbiol.*, 46: 205-210.
- de Oliveira, F.Q. 2010. Biological control of fruit fly. *Revista Verde.*, (Mossoró- RN - Brasil) 5(1): 23 – 30; see <http://revista.gvaa.com.br>

- Deshpande, M.V. 1999. Mycopesticide production by fermentation: Potential and challenges. *Crit Rev Microbiol.*, 25: 229-243.
- Dhar, P., Kaur, G. 2010. Production of cuticle - degrading proteases by *Beauveria bassiana* and their induction in different media. *African Journal of Biochemistry Research*. 4(3), pp. 65-72
- Dhillon, M.K., Singh, R., Naresh, J.S., Sharma, H.C. 2005. The melon fruit fly, *Bactrocera cucurbitae*: a review of its biology and management. *Journal of Insect Science.*, 5:40.
- Diamantidis, A.D., Carey, J.R., Papadopoulus, N.T. Life- history of an invasive tephritid. Department of agriculture, crop production and rural environmental, University of Thessaly, Phytokou St. N, Ionia 384 46, Magnisias, Greece. 2008.
- Dimbi, S., Maniania, N.K., Lux, A.S., Ekesi, S., Mueke, K.J. 2003. Pathogenicity of *Metarhizium anisopliae* (Metsch.) Sorokin and *Beauveria bassiana* (Balsamo) Vuillemin, to three adult fruit fly species: *Ceratitis capitata* (Wiedemann), *C. rosa* var. *Fasciventris* Karsch and *C. cosyra* (Walker) (Diptera: Tephritidae). *Mycopathologia.*, 156: 375-382.
- Drew, R.A.I., Romig, M.C., 1997. Overview - Tephritidae in the Pacific and South-East Asia. In: Allwood, A.J., Drew, R.A.I. (Eds.), Management of Fruit Flies in the Pacific (pp. 43-46). ACIAR Proceedings No. 76. Australian Center for International Agricultural Research, Canberra.

- Duarte, V.S., Silva, R.A., Wekesa, V.W., Rizzato, F.B., Dias, C.T.S., Delalibera, I. Jr. 2009. Impact of natural epizootics of the fungal pathogen *Neozygites floridana* (Zygomycetes: Entomophthorales) on population dynamics of *Tetranychus evansi* (Acari: Tetranychidae) in tomato and nightshade. *Biol Control.*, 51: 81-90.
- Dumas, C., Matha, V., Quiot, J.M., Vey, A. 1996. Effects of destruxins, cyclic depsipeptide mycotoxins, on calcium balance, and phosphorylation of intracellular proteins in lepidopteran cell lines. *Comp. Biochem. Physiol. Pharmacol. Toxicol. Endocrinol.*, 114: 213-219.
- Easterbrook, M. A. 1992. The possibilities for control of two-spotted spider mite *Tetranychus urticae* on field-grown strawberries in the UK by predatory mites. *Biocontrol. Sci. Technol.*, 2: 235-245.
- Eisenbrand, G., Pool-Zobel, B., Baker, V., Balls, M., Blaauboer, B.J., Boobis, A., Carere, A., Kevekordes, S., Lhuguenot, J.-C., Pieters, R., Kleiner, J., 2002. Methods on in vitro toxicology. *Food Chem. Toxicol.*, 40: 193-236.
- Ekesi, S., Maniania, N.K., Lux, S.A. 2002. Mortality in three African tephritid fruit fly puparia and adults caused by the entomopathogenic fungi *Metarhizium anisopliae* and *Beauveria bassiana*. *Biocontrol Sci Technol.*, 12: 7-17.
- El Damir, M. 2006. Effect of growing media and water volume on conidial production of *Beauveria bassiana* and *Metarhizium anisopliae*. *J Biol Sci.*, 6(2): 269-274.

- Escudero, L.A., Ferragut, F. 2005. Life-history of predatory mites *Neoseiulus californicus* and *Phytoseiulus persimilis* (Acari: Phytoseiidae) on four spider mite species as prey, with special reference to *Tetranychus evansi* (Acari: Tetranychidae). *Biol Control.*, 32: 378-384.
- Evans, H.C. 1982. Entomogenous fungi in tropical forest ecosystems: an appraisal. *Ecol Entomol.*, 7: 47-60.
- Faifer, G.C., Zabal, O., Godoy, H.M. 1992. Further studies on the hematopoietic damage produced by a single dose of T-2 toxin in mice. *Toxicology.*, 75: 169-174.
- Faramarzi, M.A., Fazeli, M., Yazdi, M.T., Adrangi, S., Al-Ahmadi, K.J., Tasharrofi, N., Mohseni, F.A. 2009. Optimization of cultural conditions for production of chitinase by a soil isolate *Massilia timonae*. *Biotechnol. Bioseparation.*, 8(1): 93-99.
- Faria, M.R., Wraight, S.P. 2007. Mycoinsecticides and mycoacaricides: a comprehensive list with worldwide coverage and international classification of formulation types. *Biol Control.*, 43:237-256.
- Ferron, P. 1978. Biological control of insect pests by entomogenous fungi. *Annu Rev Entomol.*, 23: 409-442.
- Ferron, P. 1985. Fungal control. In: Kerkut, G.A., Gilbert, L.I. (Eds.), *Comparative Insect Physiology, Biochemistry and Pharmacology*, Vol. 12 (pp. 313-346), Oxford: Pergamon Press.
- Fiedler, Z., Sosnowska, D. 2007. Nematophagous fungus *Paecilomyces lilacinus* (Thom) Samson is also a biological agent for control of greenhouse insects and mite pests. *Biol Control.*, 57: 547-558.

- Furtado, I.P., Moraes, G.J., Kreiter, S., Tixier, M.S., Knapp, M. 2007. Potential of a Brazilian population of the predatory mite *Phytoseiulus longipes* as a biological control agent of *Tetranychus evansi* (Acari: Phytoseiidae, Tetranychidae). *Biol Control.*, 42: 139-147.
- Gabriel, B.P. 1968. Histochemical studies of the insect cuticle infected by the fungus *Entomophthora coronata*. *J. Invertebr. Pathol.*, 11: 82-89.
- Gallo, J. 2007. Integrated plant control: System and management. *Encyclopedia of Pest Management.*, 2: 279-287.
- Garcia, A.S., Messias, C.L., De Souza, H.M.L., Piedramuena, A.E. 1984. Patogenicidade de *Metarhizium anisopliae* var. *anisopliae* a *Ceratitis capitata* (Wied) (Diptera, Tephritidae). *Rev. Bras. Entomol.*, 28: 421-424.
- Geden, C.J., Rutz, D.A., Steinkraus, D.C. 1995. Virulence of different isolates and formulations of *Beauveria bassiana* for house flies and the parasitoid *Muscidifurax raptor*. *Biol. Control.*, 5: 615-621.
- Gelernter, W.D., Lomer, C.J. 2000. Success in biological control of above-ground insects by pathogens. In: Gurr, G., Wratten, S. (Eds.), *Biological control: measures of success* (pp. 297-322). Dordrecht, The Netherlands: Kluwer Academic.
- Genthner, F.J., Chancy, C.A., Couch, J.A., Foss, S.S., Middaugh, D.P., George, S.E., Warren, M.A., Bantle, J.A. 1998. Toxicity and pathogenicity testing of the insect pest control fungus *Metarhizium anisopliae*. *Arch. Environ. Contam. Toxicol.*, 35: 317-324.
- Gillespie, A.T., Claydon, N. 1989. The use of entomogenous fungi for pest control and the role of toxins in pathogenesis. *Pesticide Sci.*, 27: 203-215.

- Gillespie, J., Burnett, P.C., Charnley, A.K. 2000. The immune response of the desert locust *Schistocerca gregaria* during mycosis of the entomopathogenic fungus *Metarhizium anisopliae* during penetration of host (*Manduca sexta*) cuticle. *J Gen Microbiol.*, 135: 2233–2239.
- Glare, T.R., Inwood, A.J. 1998. Morphological and genetic characterisation of *Beauveria* spp. from New Zealand. *Mycol Res.*, 102: 250-256.
- Goettel, M.S., Hajek, A.E. 2000. Evaluation of non-target effects of pathogens used for management of arthropods. In: Wajnberg, E., Scott, J.K., Quimby, P.C. (Eds.), *Evaluating indirect ecological effects of biological control* (pp. 81-97). Wallingford: CABI Publishing.
- Goettel, M.S., Inglis, G.D. 1997. Fungi: Hyphomycetes. In: Lacey, L.A. (Ed.), *Manual of techniques in insect pathology* (pp. 211-249), San Diego: Academic press.
- Goettel, M.S., Roberts, D.W. 1992. Mass production, formulation and field application of entomopathogenic fungi. In: Lomer, C.J., Prior, C. (Eds.), *Biological Control of Locusts and Grasshopper* (pp. 230-238), CAB International, Wallingford: Oxon.
- Gopalakrishnan, C., Anusuya, D., Narayanan, K. 1999. *In vitro* production of conidia of entomopathogenic fungus *Paecilomyces farinosus*. *Entomology.*, 24: 389-392.
- Granham, J.E., 1985. Spider mites. In: Helle, W., Sabelis, M.W. (Eds.), *Their biology, natural enemies and control* (pp. 367-370). Amsterdam, the Netherlands.
- Grove, J.F., Pople, M. 1980. The insecticidal effect of beauvericin and the enniatin complex. *Mycopathologia.*, 70:103-105.

- Gutleb, A., Morrison, E., Murk, A.J. 2002. Cytotoxicity assay for mycotoxins produced by *Fusarium* strains: a review. *Environ. Toxicol. Pharmacol.*, 11: 309-320.
- Ha, N.G., Kim, S.Y., Kang, J.H., Kang, P.D., Sung, G.B., Hong, I.P. 2005. Biological activities and cultural characteristics of an entomogenous fungus, *Paecilomyces tenuipes* (Peck) Samson. *Korean J. Seric Sci.*, 47: 12-17.
- Hajek, A.E. 1997. Ecology of terrestrial fungal entomopathogens. *Adv Microb Ecol.*, 15: 193-249.
- Hajek, A.E. Natural Enemies: An Introduction to Biological Control (378 pp.). Cambridge, UK: Cambridge University Press. 2004.
- Hajek, A.E., Delalibera, I. Jr. 2010. Fungal pathogens as classical biological control agents against arthropods. *BioControl.*, 55: 147-158.
- Hajek, A.E., McManus, M.L., Delalibera, I. Jr. 2007a. A review of introductions of pathogens and nematodes for classical biological control of insects and mites. *Biol Control.*, 41:1-13.
- Hajek, A.E., St. Leger, R.J. 1994. Interactions between fungal pathogens and insect hosts. *Annu. Rev. Entomol.*, 39: 293-322.
- Hall, R.A., Papierok, B. 1982. Fungi as biological agents of arthropods of agricultural and medical importance. *Parasitology.*, 84: 205-240.
- Hanelt, M., Gareis, M., Kollarzik, B. 1994. Cytotoxicity of mycotoxins evaluated by the MTT-cell culture assay. *Mycopathologia.*, 128: 167-174.
- Harada, Y., Akiyama, N., Yamamoto, K., Shiota, Y. 1995. Production of *Cordyceps militaris* fruit body on artificially inoculated pupae of *Mamestra brassicae* in the laboratory. *Nippon Kingakukai Kaiho.*, 36: 63-72.

- Hardy, D. 1973. The fruit flies (Tephritidae-Diptera) of Thailand and bordering countries. *Pac. Insects Monogr.*, 31: 1-353.
- Hedgecock, S., Moore, D., Higgins, P.M., Prior, C. 1995. Influence of moisture content on temperature tolerance and storage of *Metarhizium flavoviride* in an oil formulation. *Biocontrol Sci Technol.*, 5: 371-377.
- Helyer, N., Gill, G., Bywater, A., Chambers. 1992. Elevated humidities for control of chrysanthemum pests with *Verticillium lecanii*. *Pesticide Science.*, 36:373-378.
- Higgins, D.G., Bleasby, A.J., Fuchs, R. 1992. CLUSTAL V: improved software for multiple sequence alignment. *Comput Appl Biosci.*, 8: 189-191.
- Hoddle, M. S. The biology and management of perseae mite, *Oligonychus perseae* Tuttle, Baker and Abbatiello (Acari: Tetranychidae). Department of Entomology, University of California, USA. 2004. See: <http://www.biocontrol.ucr.edu/mitel.htm>
- Hollingsworth, R., Allwood, A.J. Melon fly (2pp). SPC Pest Advisory Leaflet. Draft. 2000.
- Hong, T.D., Gunn, J., Ellis, R.H., Jenkins, N.E., Moore, D. 2001. The effect of storage environment on the longevity of conidia of *Beauveria bassiana*. *Mycol Res.*, 105: 597-602.
- Howlett, F.M. 1912. The effects of oil of citronella on two species of *Dacus*. *Trans. R. Entomol. Soc.*, 60: 412-418.

- Htar Htar Naing, Wint War Kyaw, Si Thu Zaw, Sai Than Zaw Tun, San Li Seng, Khin Htar Nge, Sandar Hlaing, Kyaw Min Han, Chit Darli, Win Min Thant, Aung Kyaw Oo and Ye Min Oo. "Exploration of species composition of fruit fly and monitoring its population on Jujube (*Ziziphus jujube* Mill) in Yezin area". Final Year Special Problem, Yezin Agricultural University, Myanmar, 2008.
- Humber, R.A. 2008. Evolution of entomopathogenicity in fungi. *J. Invertebr Pathol.*, 98(3): 262-266.
- Humber, R.A. Entomopathogenic fungal identification (26 pp.), APS/ESA workshop, USDA-ARS plant protection research unit, Ithaca, NY. 1998.
- Hussey, N.W., Scopes, N.E.A. 1985. Greenhouse vegetables. In: Helle, W., Sabelis, M.W. (Eds.), *Spider mites, their biology, natural enemies and control*. Amsterdam: Elsevier. 1: 285-296.
- Hywel-Jones, N.L. (2001) A review of invertebrate pathogenic Clavicipitaceae of Thailand. *4th BRT Annual Meeting*: 34-41.
- IAEA. Trapping Guidelines For Area-wide fruit fly programmes (48 pp.). Insect pest control section, Joint FAO/IAEA Division of Nuclear Techniques in Food and Agriculture, International Atomic Energy Agency, Wagramer Strasse 5. A-1400 Vienna, Austria. 2003.
- Ibrahim, Y.B., Low, W. 1993. Potential of mass production and field efficacy of isolates of the entomopathogenic fungi *Beauveria bassiana* and *Paecilomyces fumosoroseus* on *Plutella xylostella*. *J. Invertebr Pathol.*, 39: 222-232.

- Ignoffo, C.M. 1973. Effects of entomopathogens on vertebrates. *Ann NY Acad Sci.*, 217: 141-172.
- Ignoffo, C.M. 1985. Manipulating enzootic-epizootic diseases of arthropods, p.243-262. In: Hoy, M.A., Herzog, D.C. (Eds.), *Biological control in agricultural IPM systems*. San Diego: Academic Press.
- Inglis, D.G., Goettel, S.M., Butt, T.M., Strasser, H. 2001. Use of Hyphomycetes fungi for managing insect pests. In: Butt T.M., Jackson, C., Magan, N. (Eds.), *Fungi as biocontrol agents: progress, problems and potential* (pp. 23-69). Wallingford: CABI International.
- Irigaray, F.J.S.C., Marco-Mancebon, V., Perez-Moreno, I. 2003. The entomopathogenic fungus *Beauveria bassiana* and its compatibility with triflumuron: effect on the two-spotted spider mite *Tetranychus urticae*. *Biol Control.*, 26: 168-173.
- Jackson, M.A. 1997. Optimizing nutritional conditions for the liquid culture production of effective fungal biological control agents. *J. Ind Microbiol Biotechnol.*, 19: 180-187.
- Jackson, M.A. 2007. The biotechnology of producing and stabilizing living, microbial biological control agents for insect and weed control. In: Hou, C.T., Shaw, F.J. (Eds.), *Biocatalysis and biotechnology: functional foods and industrial products* (pp 533-543). Boca Raton: CRC Press.
- Janes, D., Umek, A., Kreft, S. 2006. Evaluation of antibacterial activity of extracts of five species of woodcolonizing fungi. *J. Basic Microbiol.*, 46(3): 203-207.
- Jeppson, L.R., Keifer, H.H., Baker, E.W. Mites injurious to economic plants. Berkeley, CA: University of California Press. 1975.

- Jimenez, J.I., Mercier, J. 2005. Optimization of volatile organic compound production from rye grain culture of *Muscodor albus* for postharvest fumigation. *Phytopathology.*, 95: S48.
- Jones, E.B.G. 2004. Fungi on arthropods, crustaceans and fish. In: Jones, E.B.G., Tanticharoen, M., Hyde, K.D. (Eds.), *Thai Fungal Diversity*. Thailand. 227-239.
- Jones, E.B.G., Hyde, K.D. 2004. Introduction to Thai fungal diversity. In: Jones, E.B.G., Tanticharoen, M., Hyde, K.D. (Eds.), *Thai fungal diversity* (pp. 7-35.), BIOTEC, Thailand.
- Jung, E.C., Kim, K.D., Bae, C.H., Kim, J.C., Kim, D.K., Kim, H.H. 2007. A mushroom lectin from ascomycete *Cordyceps militaris*. *Biochim Biophys Acta.*, 1770: 833-838.
- Kamata, N., Sato, H., Shimazu, M. 1997. Seasonal changes in the infection of pupae of the beech caterpillar, *Quadricalcarifera punctatella* (Motsch.) (Lepidoptera: Notodontidae), by *Cordyceps militaris* Link (Clavicipitales, Clavicipitaceae) in the soil of the Japanese beech forest. *J. Appl. Entomol.*, 121: 17-21.
- Karanja, L.W., Phiri, N.A., Oduor, G.I. Effect of different solid substrates on mass production of *Beauveria bassiana* and *Metarhizium anisopliae* entomopathogens. CABI Africa, ICRAF Complex, United Nations Avenue, Gigiri, Nairobi: Kenya. 2010.
- Kaya, H.K., Lacey, L.A. 2007. Introduction to microbial control. In: Lacey, L.A., Kaya, H.K. (Eds.), *Field manual of techniques in invertebrate pathology*. 3-7.

- Keller, S. 1997. The genus *Neozygites* (Zygomycetes, Entomophthorales) with special reference to species found in tropical regions. *Sydowia.*, 49: 118-146.
- Kerry, B.R. 1990. An assessment of progress toward microbial control of plant parasitic nematode. *J. Nematol.*, 22: 621-631.
- Khachatourians, G.G. 1991. Physiology and genetics of entomopathogenic fungi. In: Arora, D.K., Ajello, L., Mukerji, K.G. (Eds.), *Handbook of Applied Mycology. vol ~. Humans. Animals and Insects* (pp. 613-663). New York: Marcel Dekker.
- Kirkland, B.H., Eisa, A., Keyhani, N.O. 2005. Oxalic acid as a fungal acaracidal virulence factor. *J Med Entomol.*, 42(3): 346-351.
- Kneifel, H., Konig, W.A., Loeffler, W., Muller, R. 1977. Ophiocordin, an antifungal antibiotic of *Cordyceps ophiglossides*. *Arch. Microbial.*, 113: 121-130.
- Koh, J.H., Yu, K.W., Suh, H.J., Choi, Y.M., Ahn, T.S. 2002. Activation of macrophages and the intestinal immune system by orally administered decoction from cultured mycelium of *Cordyceps sinensis*. *Biosci Biotechnol Biochem.*, 66: 407-411.
- Kostiainen, T., Hoy, M.A. The Phytoseiidae as biological control agents of pest mites. University of Florida, Florida Agricultural Experiment Station, IFAS Publication, Monograph 17. 1996.
- Kulkarni, N.S. 1999. Utilization of fungal pathogen *Nomuraea rileyi* (Farlow) Samson for the management of lepidopterous pests. Ph.D. Thesis, University of Agricultural Sciences, Dharwad. 1999.

- Kuo, Y., Lin, C., Tsai, W., Wu, C., Chen, C., Shiao, M. 1994. Growth inhibitors against tumor cells in *Cordyceps sinensis* other than cordycepin and polysaccharides. *Cancer Invest.*, 12: 611-615.
- Lacey, L.A., Frutos, R., Kaya, H.K., Vail, P. 2001. Insect pathogens as biological control agents: Do they have a future? *Biol Control.*, 21: 230-248.
- Lacey, L.A., Neven, L.G. 2006. The potential of the fungus, *Muscodor albus*, as a microbial control agent of potato tuber moth (Lepidoptera: Gelechiidae) in stored potatoes. *J Invertebr Pathol.*, 91: 195-198.
- Lee, S.Y., Kinoshita, H., Ihara, F., Igarashi, Y., Nihira, T. 2008. Identification of novel derivative of helvolic acid from *Metarhizium anisopliae* grown in medium with insect component. *J. Biosci Bioeng.*, 105:476-480.
- Lee, S.Y., Nakajima, I., Ihara, F., Kinoshita, H., Nihira, T. 2005. Cultivation of entomopathogenic fungi for the search of antibacterial compounds. *Mycopathologia.*, 160: 321-325.
- Leopold, J., Samsina'kova, A., Misikova, S. 1973. Enzymic character of toxic substances released by the entomophagous fungus *Beauveria bassiana* and their stimulation of their production. *Zentralbl. Bakteriol. Parasitenkd. Infektionskr.*, 128: 31-41.
- Lezama-Cutierrez, R., Trujillo-De-La Luz, A., Molina-Ochoa, J., Rebolledo-Dominguez, O., Pescador, A.R., Lopez- Edwards, M., Aluja, M. 2000. Virulence of *Metarhizium anisopliae* (deuteromycotina: hyphomycetes) on *Anastrepha ludens* (Diptera: Tephritidae): laboratory and field trials. *J. Econ Entomol.*, 93: 1080-1084.

- Li, Z., Alves, S.B., Roberts, D.W., Fan M., Delalibera, I.Jr., Tang, J., Lopes, R.B., Faria, M., Rangel, D.E.N. 2010. Biological control of insects in Brazil and China: history, current programs and reasons for their successes using entomopathogenic fungi. *Biocontrol Sci Tech.*, 20 (2): 117-136
- Li, S.P., Yang, F.Q., Tsim, K.W.K. 2006. Quality control of *Cordyceps sinensis*, a valued traditional Chinese medicine. *J. Pharm. Biomed. Anal.*, 41: 1571-1584.
- Liebhold, A., Zhou, G., Gribko, L., Hohn, M. 1996. Models that predict the spatial dynamics of gypsy moth, *Lymantria dispar* (L.), populations (pp. 491-498). In: Hain, F.P., Salom, S.M., Ravlin, W.F., Payne, T.L., Raffa, K.F. (Eds.), Behavior, population dynamics and control of forest insects. Ohio State University, Wooster. lifesci/research/entomopathogenicfungi/fungal/
- Liu, H., Skinner, M., Parker, B.L., Brownbridge, M. 2002. Pathogenicity of *Beauveria bassiana*, *Metarhizium anisopliae* (Deuteromycotina: Hyphomycetes), and other entomopathogenic fungi against *Lygus lineolaris* (Hemiptera: Miridae). *J. Econ Entomol.*, 95:675-681.
- Liu, J.C., Boucias, D.G., Pendland, J.C., Liu, W.Z., Maruniak, J. 1996. The mode of action of Hirsutellin A on eukaryotic cells. *J. Invertebr. Pathol.*, 67: 224-228.
- Liu, X. Z., Chen, S. Y. 2002. Nutritional requirement of the nematophagous fungus *Hirsutella rhossiliensis*. *Biocontrol Sci. Techn.*, 12: 381-393.

- Loutelier, C., Cherton, J.C., Lange, C., Traris, M., Vey, A. 1996. Studies on the dynamics of the production of destruxins by *Metarhizium anisopliae*: direct high-performance liquid chromatographic and fast atom bombardment mass spectrometric analysis correlated with biological activity tests. *J. Chromatogr. A.*, 738: 181-189.
- Luangsa-Ard, J.J., Tسانathai, K., Mongkolsamrit, S., Hywel-Jones, N.L. Atlas of invertebrate-pathogenic fungi of Thailand (volume 1). BIOTEC, NSTDA, Bangkok, Thailand. 2007.
- Madelin, M.F. 1966. Fungal parasites of insects. *Ann Rev Entomol.*, 11: 423-448.
- Maia, M.M.D., Heasley, A., Camargo de Moraes, M.M., Melo, E.H.M., Moraes, M.A., Ledingham, W.M. 2001. Effect of culture conditions on lipase production by *Fusarium solani* in batch fermentation. *Bioresour. Technol.*, 76: 23-27.
- Malavasi, A., Zucchi, R.A. Moscas-das-frutas de importância econômica no Brasil. Conhecimento básico e aplicado. Holos, Riberão Preto. 327 p. Brazil. 2000.
- Maniania, N.K. 1991. Potential of some fungal pathogens for the control of pests in the tropics. *Insect Sci Applied.*, 12: 63-76.
- Maniania, N.K., Bugeme, D.M., Wekesa, V.W., Delalibera, I. Jr., Knapp, M. 2008. Role of entomopathogenic fungi in the control of *Tetranychus evansi* and *Tetranychus urticae* (Acari: Tetranychidae), pests of horticultural crops. *Exp. Appl. Acrol.*, 46: 259-274.
- Maniania, N.K., Takasu, K. 2006. Development of microbial control agents at the international centre of insect physiology and ecology. *Bulletin of the Institute of Tropical Agriculture Kyushu University.*, 29:1-9.

- Mascarin, G.M., Alves, S.B., Lopes, R.B. 2010. Culture media selection for mass production of *Isaria fumosorosea* and *Isaria farinose*. *Braz Arch Biol Technol.*, 53(4): 753-761.
- McCoy, C.W. 1990. Entomogenous fungi as microbial pesticides. In: Baker, R.R., Dunn, P.E. (Eds.), *New directions in biological control* (pp. 139-159). A.R. Liss, New York.
- McCoy, C.W., Beavers, G.M., Tarrant, C.A. 1985. Susceptibility of *Artipus floridanus* to different isolates of *Beauveria bassiana*. *Flo. Entomol.*, 68(3): 402-409.
- McCoy, C.W., Samson, R.A., Boucias, D.G. 1988. Entomogenous fungi. In: Ignoffo, C.M. (Ed.), *CRC Handbook of Natural Pesticides. Microbial Insecticides. Part A: Entomogenous protozoa and fungi Vol 5* (pp. 151-236). Boca Raton, USA: CRC Press.
- McMurtry, J.A. 1982. The use of phytoseiids for biological control: progress and future prospects. In: Hoy, M.A. (Ed.), *Recent advances in knowledge of the phytoseiidae* (pp. 23-48). Berkeley: University of California Press.
- McMurtry, J.A., Croft, B.A. 1997. Life-styles of phytoseiid mites and their roles in biological control. *Annual Review of Entomology* 42: 291-321.
- Meyling, N.V. 2008. Ecology of entomopathogenic fungi in agroecosystems. Organic eprint. Available online: http://topics.scirus.com/Ecology_of_entomopathogenic_fungi_in_agroecosystems.html
- Meyling, N.V., Eilenberg, J. 2007. Ecology of the entomopathogenic fungi *Beauveria bassiana* and *Metarhizium anisopliae* in temperate agroecosystems: potential for conservation biological control. *Biol Control.*, 43: 145-155.

- Mikami, Y., Yazawa, K., Fukushima, K., Arai, T., Udagawa, S., Samon, R.A. 1989. Paecilotoxin production in clinical or terrestrial isolates of *Paecilomyces lilacinus* strains. *Mycopathologia* 108, 195-199.
- Miller, G.L. 1959. Use of dinitrosalicylic acid for estimation of reducing sugar. *Anal. Chem.*, 31: 426-428.
- Mochi, D.A., Monteiro, A.C., Bortoli, S.A., Dória, H.O.S., Barbosa, J.C. 2006. Pathogenicity of *Metarhizium anisopliae* for *Ceratitis capitata* (Wied.) (Diptera: Tephritidae) in soil with different pesticides. *Neotrop Entomol.*, 35: 382-389.
- Mustafa, U., Kaur, G. 2009. Effects of carbon and nitrogen sources and ratio on the germination, growth and sporulation characteristics of *Metarhizium anisopliae* and *Beauveria bassiana* isolates. *Afr J Agric Res.*, 3(10): 922-930.
- Mwamburi, L.A., Laing, M.D., Miller, R.M. 2010. Laboratory screening of insecticidal activities of *Beauveria bassiana* and *Paecilomyces lilacinus* against larval and adult house fly (*Musca domestica* L.). *Afr Entomol.*, 18(1):38-46.
- Nakamura, K., Yamaguchi, Y., Kagota, S., Kwon, Y.M., Shinozuka, K., Kunitomo, M. 1999. Inhibitory effect of *Cordyceps sinensis* on spontaneous liver metastasis of Lewis lung carcinoma and B16 melanoma cells in syngeneic mice. *Jpn. J. Pharmacol.*, 79: 335-341.
- Nan, J.X., Park, E.J., Yang, B.K., Song, C.H., Ko, G., Sohn, D.H. 2001. Antifibrotic effect of extracellular biopolymer from submerged mycelial cultures of *Cordyceps militaris* on liver fibrosis induced by bile duct ligation and scission in rats. *Arch Pharm Res.*, 24: 327-332.

- Nimnoi, P., Pongsilp, N., Lumyong, S. 2010. Endophytic actinomycetes isolated from *Aquilaria crassna* Pierre ex Lec and screening of plant growth promoters production. *World J Microbiol Biotechnol.*, 26: 193-203.
- Oduor, G.I. Abiotic factors and the epizootiology of *Neozygites* cf. *floridana*, a fungus pathogenic to the cassava green mite. PhD Thesis, University of Amsterdam, The Netherlands. 1995.
- Ojcius, D.M., Zychlinsky, A., Zheng, L.M., Young, J.D-E. 1991. Ionophore-induced apoptosis: role of DNA fragmentation and calcium fluxes. *Exp Cell Res.*, 197: 43-49.
- Olechowski, H. 1989. Spider mites in field crops. Ontario, Ministry of Agriculture and Food.
- Olechowski, H., Schaafsma, A. Spider mites in field crops. Ministry of Agriculture Food and Rural Affairs, Ontario. 1989.
- Oliveira, H., Fadini, M.A.M., Venzon, M., Rezenda, D., Rezende, F., Pallini, A. 2009. Evaluation of the predatory mite *Phytoseiulus macropilis* (Acari: Phytoseiidae) as a biological control agent of the two-spotted spider mite on strawberry plants under greenhouse conditions. *Exp. Appl. Acarol.*, 47: 275-283.
- Oliveira, H., Janssen, A., Pallini, A., Venzon, M., Fadini, M.A.M., Duarte, V. 2007. A phytoseiid predator from the tropics as potential biological control agent for the spider mite *Tetranychus urticae* Koch (Acari: Tetranychidae). *Biol Control.*, 42: 105-109.

- Opit, G.P., Chen, Y., Williams, K.A., Nechols, J.R., Margolies, D.C. 2005. Plant age, fertilization, and biological control affect damage caused by two-spotted spider mite on ivy geranium: development of an action threshold. *J. Am. Soc. Hortic. Sci.*, 130: 159-166.
- Osborne, L.S., Ehler, L.E., Nechols, J.R. Biological control of the two-spotted spider mite in greenhouses. Bulletin 853, University of Florida Central Florida Research and Education Center. 2807 Binion Road. Apopka, FL. 32703. 1999. See: <http://mrec.ifas.ufl.edu/lso/SpMite/b853a1.htm>
- Ovruski, S., Aluja, M., Sivinski, J., Wharton, R. 2000. Hymenopteran parasitoids of fruit-infesting Tephritidae (Diptera) in Latin America and the southern United States: Diversity, distribution, taxonomic status and their use in fruit fly biological control. *Integrated Pest Management Reviews.*, 5: 81-107.
- Panda, N., Khush, G.S. Host plant resistance to insects. CAB International. 1995.
- Pandey, A.K., Kanaujia, K.R. 2008. Effect of different grains as solid substrate on sporulation, viability and pathogenicity of *Metarhizium anisopliae* (Metschnikoff) Sorokin. *Biol Control.*, 22(2): 369-374.
- Pandey, A., Azmi, W., Singh, J., Banerjee, U.C. 1999. Types of fermentation and factors affecting it. In: Joshi, V.K., Pandey, A. (Eds.), *Biotechnology food fermentation* (pp. 383-426). New Delhi, Educational Publishers.
- Pangrahi, S. 1993. Bioassay of mycotoxins using terrestrial and aquatic, animal and plant species. *Food Chem. Toxic.*, 31: 767-790.

- Parine, N.R., Pathan, A.A.K., Sarayu, B, Nishanth, V.S., Bobbarala, V. 2010. Antibacterial efficacy of secondary metabolites from entomopathogenic fungi *Beauveria bassiana*. *International Journal of Chemical and Analytical Science.*, 1(5): 94-96.
- Paris, S., Segretain, G. 1975. Caracteres physiologiques de *Beuveria tenella* en rapport avec la virulence de souches de ce champignon pour la larve du Hanneton commun, *Melolontha melolontha* L. *Entomophaga.*, 20: 135-138.
- Park, C., Hong, S.H., Lee, J.Y., Kim, G.Y., Choi, B.T., Lee, Y.T., Park, D.I., Park, Y.M., Jeong, Y.K., Choi, Y.H. 2005. Growth inhibition of U937 leukemia cells by aqueous extract of *Cordyceps militaris* through induction of apoptosis. *Oncol Rep.*, 13: 1211-1216.
- Parvin, M. M., Asgar, M.D., Haque, M.M. 2010. Voracity of three predators on two-spotted spider mite, *Tetranychus Urticae* Koch (Acari: Tetranychidae) and their developmental stages. *Res. J. Agric. Biol. Sci.*, 6(1): 77-83.
- Paterson, R.R.M. 2008. Cordyceps: a traditional Chinese medicine and another fungal therapeutic biofactory? *Phytochemistry.*, 69: 1469-1495.
- Pedras, M.S.C., Irina, Z.L., Ward, D.E. 2002. The destruxins: synthesis, biosynthesis, biotransformation, and biological activity. *Phytochemistry.*, 59: 579-596.
- Pegram, R.A., Wyatt, R.D., Smith, T. 1982. Oosporein-toxicosis in the turkey poultry. *Avian Dis.*, 26: 47-59.
- Pell, J.K., Eilenberg, J., Hajek, A.E., Steinkraus, D.C. 2001. Biology, ecology and pest management potential of Entomophthorales. In: Butt, T.M., Jackson, C., Magan, N. (Eds.), *Fungi as biocontrol agents: progress, problems and potential* (pp. 71-153). Wallingford: CAB International.

- Pell, J.K., Wilding, N., Player, A.L., Clark, S.J. 1993. Selection of an isolate of *Zoophthora radicans* (Zygomycetes: Entomophthorales) for biocontrol of the diamondback moth, *Plutella xylostella* L. (Lepidoptera: Yponomeutidae). *J. Invertebr Pathol.*, 61:75-80.
- Price, P.W. 1997. Insect ecology (868pp.). New York, John Wiley and Sons, Inc.
- Purcell, M.F. 1998. Contribution of biological control of integrated pest management of Tephritid fruit flies in the Tropics and Subtropics. *Integrated Pest Management Reviews.*, 3: 63-83.
- Quesada-Moraga, E., Navas-Cortés, J.A., Maranhao, E.A.A., Ortiz-Urquiza, A., Santiago-Álvarez, C. 2007. Factors affecting the occurrence and distribution of entomopathogenic fungi in natural and cultivated soils. *Mycol Res.*, 111: 947-966.
- Quesada-Moraga, E., Ruiz-Garcia, A., Santiago-Alvarez, C. 2006. Laboratory evaluation of entomopathogenic fungi *Beauveria bassiana* and *Mearhizium anisopliae* against puparia and adults of *Ceratitis capitata* (Diptera: Tephritidae). *J. Econ Entomol.*, 99: 1955-1966.
- Quiot, J.M., Vey, A., Vago, C. 1985. Effects of mycotoxins on invertebrate cells in vitro. *Adv. Cell Cult.*, 4: 199-212.
- Ramadan, M.M., Messing, R.H. 2003. A survey for potential biocontrol agents of *Bactrocera cucurbitae* (Diptera: Tephritidae) in Thailand. *Proc Hawaiian Entomol Soc.*, 36: 115-122.
- Reed, J., Muench, H. 1938. A simple method of estimating fifty percent endpoints. *Am J Hyg.*, 27: 493-497.

- Rehner, S.A., Buckley, E.P. 2005. A *Beauveria* phylogeny inferred from nuclear ITS and EF1-a sequences: evidence for cryptic diversification and links to *Cordyceps* teleomorphs. *Mycologia.*, 97: 84-98.
- Rehner, S.A., De Muro, M.A., Bischoff, J.F. 2006a. Description and phylogenetic placement of *Beauveria malawiensis* sp. nov. (Clavicipitaceae, Hypocreales). *Mycotaxon.*, 98: 137-145.
- Rehner, S.A., Posada, F., Buckley, E.P., Infante, F., Castillo, A., Vega, F.E. 2006b. Phylogenetic origins of African and Neotropical *Beauveria bassiana* s.l. pathogens of the coffee berry borer, *Hypothenemus hampei*. *J Invertebr Pathol.*, 93: 11-21.
- Reubel, G.H., Gareis, M., Amselgruber, W.M. 1987. Cytotoxicity evaluation of mycotoxins by an MTT-Bioassay. *Mycotoxin Res.*, 3: 85-95.
- Ribeiro, A.E.L., Gondim, M.G.C.Jr., Calderan E. 2009. Host range of *Neozygites floridana* isolates (Zygomycetes: Entomophthorales) to spider mites. *J. Invertebr Pathol.*, 102: 196-202.
- Riddell, R.W. 1950. Permanent stained mycological preparations obtained by slide culture. *Mycologia.*, 42: 265-270.
- Roberts, D.R. 1981. Toxins of entomopathogenic fungi. In: Burges, H.D. (Ed.), *Microbial control of pests and diseases 1970-1980* (pp 441-464). London: Academic Press.
- Roberts, D.R., Hajek, A.E. 1992. Entomopathogenic fungi as bioinsecticides. In: Leatham, G.F. (Ed.), *Frontiers in industrial mycology* (pp 144-159). Chapman and Hall, New York,

- Robl, D., Sung, L.B., Novakovich, J.H., Marangoni, P.R.D., Zawadneak, M.A.C., Dalzoto, P.R., Gabardo, J., Pimentel, I.C. 2009. Spore production in *Paecilomyces lilacinus* (Thom.) Samson strains on agro-industrial residues. *Braz J Microbiol.*, 40: 296-300.
- Rombach, M.C. 1988. Production of *Beauveria bassiana* (Deuteromycotina: Hyphomycetes) sympoduloconidia in submerged culture. *Entomophaga.*, 34: 45-52.
- Roy, H.E., Brodie, E.L., Chandler, D., Goettel, M.S., Pell, J.K., Wajnberg, E., Vega, F. 2010. Deep space and hidden depths: understanding the evolution and ecology of fungal entomopathogens. *Biocontrol.*, 55: 1-6.
- Sabelis, M.W. 1981. Biological control of two spotted spider mites using phytoseiid predators. Pudoc, Wageningen, Netherlands: I. Agric. Res. Report 910.
- Saberhagen, C., Klotz, S.A., Bartholomew, W., Drews, D., Dixon, A. 1997. Infection due to *Paecilomyces lilacinus*: a challenging clinical identification. *Clin. Infect. Dis.*, 25(6): 1411-1413.
- Safdar, A. 2002. Progressive cutaneous hyalohyphomycosis due to *Paecilomyces lilacinus*: rapid response to treatment with caspofungin and itraconazole. *Clin. Infect. Dis.*, 34 (10): 1415-1417.
- Saitou, N., Nei, M. 1987. The neighbor-joining method: a new method for reconstructing phylogenetic tree. *Mol Biol Evol.*, 4(4): 406-425.
- Salles, L.A.B. Bioecologia e controle da moscas-dasfrutas sul-americana. CNPFT - Embrapa. 58p. 1995.
- Salles, L.A.B. Moscas-das-frutas, *Anastrepha fraterculus* (Wied.): bioecologia e controle. CNPFT - Embrapa. 16p. (Documentos 41). 1991.

- Samsinakova, A., Misikova, S. 1973. Enzyme activities in certain entomophagous representatives of Deuteromycetes in relationship to their virulence. *Ceska Mykol.*, 27: 55-60.
- Samson, R.A., Evans, H.C., Latg, J.P. 1988. Atlas of entomopathogenic fungi. Berlin Heidelberg. New York. Springer.
- Samuels, R.I., Paterson, I.C. 1995. Cuticle degrading proteases from insect moulting fluid and culture filtrates of entomopathogenic fungi. *Comp Biochem Physiol.*, 10: 661-669.
- Sarango, V.M.G. Monitoring and pest control of fruit flies in Thailand: new knowledge for integrated pest management. SLU, Institutionen för ekologi, Uppsala, Sweden. 2009.
- Sato, H., Kamata, N., Shimazu, M. 1997. Aerial infection of *Cordyceps militaris* Link (Clavicipitales: Clavicipitaceae) against larvae of *Quadricalcarifella punctatella* (Motschulsky) (Lepidoptera: Notodondidae). *Appl. Entomol. Zool.*, 32: 249-252.
- Sato, H., Shimazu, M. 2002. Stromata production for *Cordyceps militaris* (Clavicipitales: Clavicipitaceae) by injection of hyphal bodies to alternative host insects. *Appl Entomol Zool.*, 37: 185-192.
- Satoh, D., Horri, A., Ochiai, M., Ashida, M. 1999. Prophenoloxidase-activation enzyme of the silkworm, *Bombyx mori*. *J Biol Chem.*, 274: 7441-7453.
- Schachter, B. 2003. Slimy business-the biotechnology of biofilms. *Nat. Biotechnol.*, 2: 361-365.
- Schenck, S. 2004. Control of nematodes in tomato with *Paecilomyces lilacinus* strain 251. *Haw. Agricul. Res. Cent. Veg.*, 5: 1-5.

- Schneider, P., Misiak, M., Hoffmeister, D. 2008. *In vivo* and *in vitro* production options for fungal secondary metabolites. *Mol. Pharm.*, 234-242.
- Schmidt, R., 1989. The application of *Artemia salina* L. bioassay for screening of Fusaria toxins. In: Chelkowski, J. (Ed.), *Fusarium: mycotoxins, taxonomy and pathogenicity* (pp. 121-130). Elsevier, Amsterdam.
- Shah, F.A., Cheng, S.W., Tariq, M.B. 2005. Nutrition influences growth and virulence of the insect-pathogenic fungus *Metarhizium anisopliae*. *FEMS Microbiol Lett.*, 251(2): 259-266.
- Shah, P.A., Pell, J.K. 2003. Entomopathogenic fungi as biological control agents. *Appl Microbiol Biotech.*, 61: 413-423.
- Shelly, T.E. 1995. Methyl eugenol and the mating competitiveness of irradiated male *Bactrocera dorsalis* (Diptera: Tephritidae). *Ann. Entomol. Soc. America.*, 88: 883-886.
- Shelton, A.M., Vandenberg, J.D., Ramos, M., Wilsey, W.T. 1998. Efficacy and persistence of *Beauveria bassiana* and other fungi for control of diamondback moth (Lepidoptera: Plutellidae) on cabbage seedlings. *J. Entomol. Sci.*, 33: 142-151.
- Shi, W.B., Feng, M.G. 2006. Field efficacy of application of *Beauveria bassiana* formulation and low rate pyridaben for sustainable control of citrus red mite *Panonychus citri* (Acari: Tetranychidae) in orchards. *Biol Control.*, 39: 210-217.
- Shi, W.B., Feng, M.G. 2009. Effect of fungal infection on reproductive potential and survival time of *Tetranychus urticae* (Acari: Tetranychidae). *Exp. Appl. Acrol.*, 48: 229-237.

- Shi, W.B., Zhang, L.L., Feng, M.G. 2008a. Field trials of four formulations of *Beauveria bassiana* and *Metarhizium anisopliae* for control of cotton spider mites (Acari: Tetranychidae) in the Tarim Basin of China. *Biol Control.*, 45: 48-55.
- Shima, M. 1955. On the metabolic products of the silkworm muscardines. *Bull Seric Exp Stat.*, 14: 427-449.
- Shimazu, M. 1994. Potential of the cerambycid-parasitic type of *Beauveria brongniartii* (Deuteromycotina: Hyphomycetes) for microbial control of *Monochamus alternates* Hope (Coleoptera: Cerambycidae). *Appl. Entomol. Zool.*, 29: 127-130.
- Shimazu, M., Zhang, B., Liu, Y. 2002. Fungal pathogens of *Anoplophora glabripennis* (Coleoptera: Cerambycidae) and their virulences. *Bull. FFPRI.*, 1(382): 123-130.
- Silva, L., Loch, L.C. 1987. Sporulation of the entomopathogenic fungus *Nomuraea rileyi* on polished rice grain media. *Anais da sociedade entomologica da Brasil.*, 16: 213-222.
- Singh, S. 2003. Effects of aqueous extract of neem seed kernel and azadirachtin on the fecundity, fertility and post-embryonic development of the melonfly, *Bactrocera cucurbitae* and the Oriental fruit fly, *Bactrocera dorsalis* (Diptera: Tephritidae). *J. Appl. Entomol.*, 127: 540-547.
- Smith-Meyer, M.K.P. 1981. Mite pests of crops in southern Africa. *Science Bulletin, Department of Agriculture and Fisheries, Republic of South Africa* No. 397: 65-67.

- Soccol, C.R., Ayala, L.A., Soccol, V.T., Krueger, N., Santos, H.R. 1997. Spore production by entomopathogenic fungus *Beauveria bassiana* from declassified potato by solid-state fermentation. *Rev Microbiol.*, 28: 34-42.
- Soe, K.M., Bhromsiri, A., Karladee, D. 2010. Effects of selected endophytic actinomycetes (*Streptomyces* sp.) and bradyrhizobia from Myanmar on growth, nodulation, nitrogen fixation and yield of different soybean varieties. *J. Nat. Sci.*, 9(1): 95-109.
- Sookar, P., Bhagwant, S., Awuor Ouna, E. 2008. Isolation of entomopathogenic fungi from the soil and their pathogenicity to two fruit fly species (Diptera: Tephritidae). *J. Appl Entomol.*, 132: 778-788.
- Soper, R.S., Ward, M.G. 1981. Production, formulation and application of fungi for insect control. In: Papavizas, G.C. (Ed.), *Biological Control in Crop Production* (pp. 161-180). BARC Symposium No. 5. Allanheld, Osnum, Ottawa, NJ.
- Sosa-Gómez, R.D., Moscardi, F. 1994. Effect of till and no-till soybean cultivation on dynamics of entomopathogenic fungi in the soil. *Florida Entomol.*, 77: 284-287.
- Sosnowska, D. 2003. Możliwość zastosowania *Pochodnia chlamydosporia* (Goddard) Zare et Gams oraz *Paecilomyces lilacinus* (Thom) Samson w biologicznym zwalczaniu małowia burakowego (Heterodera schachtii Schmidt) i guzako'w korzeniowych (Meloidogyne spp.). Rozprawy Naukowe Instytutu Ochrony Ros'lin, zeszyt 9. 95 pp.

- St. Leger, R.J., Joshi, L., Roberts, D. 1998. Ambient pH is a major determinant in the expression of cuticle-degrading enzymes and hydrophobin by *Metarhizium anisopliae*. *Appl. Environ. Microbiol.*, 64:709-713.
- St. Leger, R.J., Charnley, K., Cooper, R.M. 1986a. Cuticle-degrading enzymes of entomopathogenic fungi: mechanisms of interaction between pathogen enzymes and insect cuticle. *J. Invertebr. Pathol.*, 47: 295-302.
- St. Leger, R.J., Joshi, L., Bidochka, M., Rizzo, M.J., Roberts, D.W. 1996a. Characterization and ultrastructural localization of chitinases from *Metarhizium anisopliae*, *M. flavoviride* and *Beauveria bassiana* during fungal invasion of host (*Manduca sexta*) cuticle. *Appl. Environ. Microbiol.*, 62: 907-912.
- St.Leger, R.J. 1995. The role of cuticle-degrading proteases in fungal pathogenesis of insects. *Can. J. Bot.*, 73: 1119-1125.
- Steinkraus, D.C. 2007. Documentation of naturally-occurring pathogens and their impact in agroecosystems, p.267-281. In: Lacey: L.A., Kaya, H.K. (Eds.), *Field manual of techniques in invertebrate pathology: application and evaluation of pathogens for control of insects and other invertebrate pests*, 2nd edition. Dordrecht: Springer Scientific Publishers.
- Steinkraus, D.C., Zawislak, J., Lorenz, G. Spider mites on cotton in the midsouth. University of Arkansas, Fayetteville, AR. 2001.
- Sukhirun, N., Bullangpoti, V., Pluempanupat, W. 2009. The insecticidal studies from *Alpinia galanga* and *Cleome viscosa* extract as alternative control tool to *Bactrocera dorsalis* (Hendel). *KKU Sci. J.*, 37: 71-76.

- Takahashi, S., Uchida, K., Kakinuma, N., Hashimoto, R., Yanagisawa, T., Nakagawa, A. 1998. The structures of pyridovericin and pyridomacrolidin, new metabolites from the entomopathogenic fungus, *Beauveria bassiana*. *J Antibiot.*, 51: 1051-1054.
- Tamai, M.A., Alves, S.B., de Almeida, J.E.M., Faion, M. 2002. Evaluation of entomopathogenic fungi for control of *Tetranychus urticae* Koch. (Acari, Tetranychidae). *Arquivos do Instituto Biologico.*, 69(3): 77-84.
- Tamura, K., Dudley, J., Nei, M., Kumar, S. 2007. MEGA4 : molecular evolutionary genetics analysis (MEGA) software version 4.0. *Mol Biol Evol.*, 24: 1596-1599.
- Tanada, Y., Kaya, H.K. "Insect Pathology." San Diego, CA: Academic Press. 1993.
- Thakur, D., Yadav, A., Gogoi, B.K., Bora, T.C. 2007. Isolation and screening of *Streptomyces* in soil of protected forest areas from the states of Assam and Tripura, India, for antimicrobial metabolites. *Journal de Mycologie Médicale.*, 17: 242-249.
- Thakur, R., Sandhu, S.S. 2010. Distribution, occurrence and natural invertebrate hosts of indigenous entomopathogenic fungi of Central India. *Indian J Microbiol.*, 50(1): 89-96.
- Thapinta, A., Hudak, P.F. 1998. Pesticide use and residual occurrence in Thailand. *Environ Monit Assess.*, 60: 103-114.
- Toledo, A.J. 2002. Evaluacion de algunos agentes entomopatogenos para el control microbiano de tres especies de moscas de la fruta (Diptera: Tephritidae) de importancia economica. Tesis de Doctor en Ciencias. Facultad de Ciencias. Universidad Nacional Autonoma de Mexico. Mexico, D.F. 124 p.

- Tucker, S.L., Talbot, N.J. 2001. Surface attachment and pre-penetration stage development by plant pathogenic fungi. *Annu. Rev. Phytopathol.*, 39:385-417.
- Uziel, A., Levy, K., Yuval, B. 2003. Infection of *Ceratitis capitata* by two species of the *Entomophthora muscae* species complex (Zygomycetes: Entomophthorales) in the field. *Phytoparasitica.*, 31: 1-3.
- Van der Geest, L.P.S., Elliot, S.L., Breeuwer, J.A.J., Beerling, E.A.M. 2000. Diseases of mites. *Exp. Appl. Acarol.*, 24: 497-560.
- Van der Schaaf, D.A., Ravensburg, W.J., Malais, M. 1991. *Verticillium lecanii* as a microbial insecticide against whitefly. *IOBC/ WPRS Bulletin.*, 14: 120-123.
- Vandenberg, J. D., Ramos, M. 1997. Screening of fungal isolates against larvae of the diamondback moth, 1995. *Arthropod Manage. Tests.*, 22: 420-421.
- Vandenburg, J.D., Ramos, M., Altre, J.A., 1998. Dose-response and ageand temperature-related susceptibility of the diamondback moth (Lepidoptera: Plutellidae) to two isolates of *Beauveria bassiana* (Hyphomycetes: Moniliaceae). *Environ. Entomol.*, 27: 1017-1021.
- Vargas, A.B., Mayhé-Nunes, A.J., Queiroz, J.M., Souza, G.O., Folly- Ramos, E. 2007. Efeitos de fatores ambientais sobre a Mirmecofauna em Comunidade de restinga no rio de Janeiro, RJ. *Neotropical Entomology.*, 36(1): 28-37.
- Verghese, A., Tandon, P.L., Stonehouse, J.M. 2004. Economic evaluation of the integrated management of the oriental fruit fly *Bactrocera dorsalis* (Diptera: Tephritidae) in mango in India. *Crop Protection.*, 23: 61-63.

- Vey, A., Fargues, J. 1977. Histological and ultra structural studies of *Beauveria bassiana* infection in *Leptinotarsa decemlineata* Say larvae during ecdysis. *J. Invertebr. Pathol.*, 30: 207-215.
- Vey, A., Hoagland, R., Butt, T.M. 2001. Toxic metabolites of fungal biocontrol agents. In: Butt, T. M., Jackson, C., Magan, N. (Eds.), *Fungi as biocontrol agents: progress, problems and potential* (pp. 311-345). Oxon: CAB International.
- Vey, A., Quiot, J.M. 1989. Effect cytotoxique in vitro et chez l'insecte hôte des dextrusines, toxines cyclodepsipeptidiques produites par le champignon entomopathogène *Metarhizium anisopliae*. *Can. J. Microbiol.*, 35: 1000-1008.
- Vilcinskas, A., Götz, P. 1999. Parasitic Fungi and their Interactions with the Insect Immune System. *Adv Parasit.*, 43: 267-313.
- Vimaladevi, P.S. 1994. Conidia production of the entomopathogenic fungus, *Nomuraea rileyi* and its evaluation for control of *Spodoptera litura* (Fab.) on *Ricinus communis*. *J Invertebr Pathol.*, 63: 145-150.
- Visconti, A., Minervini, F., Lucivero, G., Gambatesa, V. 1991. Cytotoxic and immunotoxic effect of *Fusarium* mycotoxins evaluated by a rapid colorimetric bioassay. *Mycopathologia.*, 113: 181- 186.
- Vrie, M. 1985. Greenhouse ornamentals. In: Helle, W., Sabelis, M.W. (Eds.), *Spider mites. Their biology, natural enemies and control* (pp. 273-283). Amsterdam, the Netherlands: Elsevier.
- Wang, T., Chen, L. 2009. Dynamic complexity of microbial pesticide model. *Nonlinear Dyn.*, 58: 539-552.

- Watanabe, N., Hattori, M., Yokoyama, E., Isomura, S., Ujita, M., Hara, A. 2006. Entomogenous fungi that produce 2,6-Pyridin Dicarboxylic Acid (Dipicolinic Acid). *J Biosci Bioeng.*, 102(4): 365-368.
- Watson, D.W., Geden, C.J., Long, S.J., Rutz, D.A. 1995. Efficacy of *Beauveria bassiana* for controlling the house fly and stable fly (Diptera: Muscidae). *Biol Control.*, 5: 405-411.
- Weems, H.V.Jr., Heppner, J.B. Oriental fruit fly, *Bactrocera* (= *Dacus*) *dorsalis* (Hendel) (Insecta: Diptera: Tephritidae). Florida Department of Agriculture and Consumer Services, Division of plant industry, and T.R. Fasulo, Florida: University of Florida Publication. 1999.
- Weinzierl, R., Henn, T., Koehler, P.G., Tucker, C.L. Microbial insecticide (13pp). IFAS Extension. University of Florida. 2009.
- Wekesa, V.W., Knapp, M., Maniania, N.K., Boga, H.I. 2006. Effects of *Beauveria bassiana* and *Metarhizium anisopliae* on mortality, fecundity and egg fertility of *Tetranychus evanse*. *J. Appl Entomol.*, 130: 155-159.
- Wekesa, V.W., Moraes, G.J., Knapp, M.; Delalibera Jr.I. 2007. Interactions of two natural enemies of *Tetranychus evansi*, the fungal pathogen *Neozygites floridana* (Zygomycetes: Entomophthorales) and the predatory mite, *Phytoseiulus longipes* (Acari: Phytoseiidae). *Biol Control.*, 41: 408-414.
- Westenfeld, F., Alston, W.K., Winn, W.C. 1996. Complicated soft tissue infection with prepatellar bursitis caused by *Paecilomyces lilacinus* in an immunocompetent host: case report and review. *J. Clin. Microbiol.*, 34(6): 1559- 1562.

- White, I. M., Elson-Harris, M. Fruit flies of economic significance: their identification and bionomics (601p.). London: International Institute of Entomology. 1992.
- White, I.M., Elson-Harris, M.M. Fruit flies of economic significance: their identification and bionomics (601 pp). Oxon, UK: CAB International. 1994.
- White, T.J., Bruns, T.D., Lee, S., Taylor, J.W. 1990. Amplification and direct sequencing of fungal ribosomal RNA genes for phylogenetics. In: Innes, M.A., Gelfand, D.H., Sninsky, J.J., White, T.J. (Eds.), *PCR Protocols: a Guide to Methods and Applications* (pp. 315-322). San Diego, CA: Academic Press.
- Widestrand, J., Lundh, T., Petterson, H., Lindberg, J.E. 1999. Cytotoxicity of four trichothecenes evaluated by three colorimetric bioassays. *Mycopathologia.*, 147: 149-155.
- Wildman, H.G. 1997. Potential of tropical microfungi within the pharmaceutical industry. In: Hyde, K.D. (Ed.), *Biodiversity of Tropical Microfungi* (pp. 29-46). Hong Kong SAR: Hong Kong University Press.
- Won, S. Y., Park, E. H. 2005. Anti-inflammatory and related pharmacological activities of cultured mycelia and fruiting bodies of *Cordyceps militaris*. *J Ethnopharmacol.*, 96: 555-561.
- Wraight, S.P., Jackson, M.A., De Kock, S.L. 2001. Production, stabilization and formulation of fungal biocontrol agents. In: Butt, T.M., Jackson, C., Magan, N. (Eds.), *Fungi as biocontrol agents: progress, problems and potential* (pp. 253-288). Wallingford: CABI Publishing.

www.benemite.com/twospot.htm. The two-spotted spider mite.

www.bugs.bio.usyd.edu.au/learning/resources/Mycology/UsesOf_Fungi/primaryProduction/BiologicalControl/insects.html. Mycology: Insect control.

www.entnemdept.ufl.edu/creatures/orn/twospotted_mite.htm. Two-spotted spider mites, *Tetranychus urticae*. University of Florida Institute of Food and Agricultural Sciences.

www.ext.colostate.edu/pubs/insect/05507.htm. Spider mite. Colorado State University.

www.growercentral.com/PDFS/two_spotted_spider_mites.pdf. Two-Spotted Spider Mites (*Tetranychus urticae*)

www.nysaes.cornell.edu/ent/biocontrol/pathogens/fungi.htm. Biological control. A Guide to Natural Enemies in North America.

www.planthealth.info/pdf_docs/mites_MN.pdf. Two-spotted spider mite. Minnesota Department of Agriculture.

www.spc.int/Pacifly/Control/Biocontrol.htm. Biological control against fruit flies in Pacific island countries and territories. 2004.

Ye, H., Liu, J.H. 2005. Population dynamics of the oriental fruit fly, *Bactrocera dorsalis* (Diptera: Tephritidae) in the Kunming area, southwestern China. *Insect Science.*, 12(5): 387-392.

Yu, K.W., Kim, K.M., Suh, H.J. 2003. Pharmacological activities of stromata of *Cordyceps scarabaecola*. *Phytother Res.*, 17: 244-249.

Yu, R., Song, L., Shao, Y., Bin, W., Wang, L., Shang, H., Wo, Y., Ye, W., Yao, X. 2004a. Isolation and biological properties of polysaccharide CPS-1 from cultured *Cordyceps militaris*. *Fitoterapia.*, 75: 465-472.

- Yu, R., Wang, L., Zhang, H., Zhou, C., Zhao, Y. 2004b. Isolation, purification and identification of polysaccharides from cultured *Cordyceps militaris*. *Fitoterapia*, 75: 662-666.
- Zimmerman, G. 1986. Insect pathogenic fungi as pest control agents. *Fortshr Zool.*, 217-231.
- Zimmermann, G. 1993. The entomopathogenic fungus *Metarhizium anisopliae* and its potential as a biocontrol agent. *Pestic Sci.*, 37:375-379.
- Zimmermann, G. 1994. Strategies for the utilization of entomopathogenic fungi. In: Proc VIth Int. Colloquium Invertebrate Pathology and Microbial Control, Montpellier, France, 1: 67-73.
- Zimmermann, G. 2007. Review on safety of the entomopathogenic fungi *Beauveria bassiana* and *Beauveria brongniartii*. *Biocontrol Sci Techn.*, 17: 553-596.