

TABLE OF CONTENTS

	Page
ACKNOWLEDGEMENTS	iii
ABSTRACT (ENGLISH)	v
ABSTRACT (THAI)	viii
LIST OF TABLES	xvii
LIST OF FIGURES	xix
ABBREVIATIONS	xxi
CHAPTER 1 INTRODUCTION	1
CHAPTER 2 LITERATURE REVIEW	7
2.1 Entomopathogenic fungi (EPFs) as biocontrol agents	7
2.2 Pathogenic activity of entomopathogenic fungi	8
2.3 Production of entomopathogenic fungi and their cytotoxic effect	12
2.4 Economic importance of fruit fly (<i>Bactrocera</i> spp.)	16
2.4.1 Fruit fly damage symptom	17
2.4.2 Biology (Life Cycle) of fruit fly	18
2.4.3 Management strategies	19
2.4.3.1 Trapping	19
2.4.3.2 Cultural control	19
2.4.3.2.1 Sanitation measures	19
2.4.3.2.2 Resistant crops	20

TABLE OF CONTENTS (Continued)

	Page
2.4.3.2.3 Bagging	20
2.4.3.2.4 Early harvest	20
2.4.3.3 Chemical control	20
2.4.3.4 Biological control	20
2.5 Economic importance of two-spotted spider mite (<i>Tetranychus urticae</i> (Koch))	21
2.5.1 Two-spotted spider mite damage symptom	22
2.5.2 Biology of two-spotted spider mite	23
2.5.3 Management of two spotted spider mite	24
2.5.3.1 Cultural control	25
2.5.3.2 Biological control	25
2.5.3.3 Chemical control	27

CHAPTER 3 Collection, Isolation and Identification of Entomopathogenic

Fungi

3.1 Introduction	29
3.2 Objective	30
3.3 Methodology	30
3.3.1 Sample site	30
3.3.2 Collection and isolation of collected entomopathogenic fungi	31
3.3.3 Identification of entomopathogenic fungi	32

TABLE OF CONTENTS (Continued)

	Page
3.3.3.1 Morphological examination	32
3.3.3.2 DNA extraction, PCR and sequencing	32
3.3.4 Phylogenetic analysis	33
3.4 Results	34
3.4.1 Isolation and identification of entomopathogenic fungi	34
3.4.2 Molecular identification of fungal isolates	35
3.5 Discussion	36
 CHAPTER 4 Infectivity of Insect Pathogenic Fungal Isolates to Fruit Fly Pupa,	
<i>Bactrocera</i> spp. (Diptera: Tephritidae)	
4.1 Introduction	39
4.2 Objectives	41
4.3 Methodology	42
4.3.1 Fruit fly collection and handling	42
4.3.2 Fungal isolates and preparation of conidial suspensions	42
4.3.3 <i>In vitro</i> screening of virulence isolates against fruit fly pupa	43
4.3.4 Recovery test of fungi from infected pupae	44
4.3.5 Soil bioassays	44
4.3.6 Statistical analysis	45
4.4 Results	45
4.4.1 Virulence of fungal isolates	45

TABLE OF CONTENTS (Continued)

	Page
4.4.2 Pathogenic activity of fungal isolates in soil bioassay	49
4.5 Discussion	51
CHAPTER 5 Selection of the Effective Fungal Pathogens to Two-spotted Spider Mites <i>Tetranychus urticae</i> Koch (Acari: Tetranychidae)	
5.1 Introduction	54
5.2 Objectives	55
5.3 Methodology	55
5.3.1 Culturing two-spotted spider mites	55
5.3.2 Fungal culture and preparing the conidial suspension	56
5.3.3 Screening of virulence fungal isolates to <i>T. urticae</i>	56
5.3.4 Plant growing	57
5.3.5 Pot experiments	57
5.3.6 Statistical analysis	59
5.4 Results	59
5.4.1 Susceptibility of <i>T. urticae</i> to fungal isolates	59
5.4.2 Control of <i>Tetranychus urticae</i> (Koch) in pot experiments	64
5.5 Discussion	65
CHAPTER 6 Spore Production of EPFs in Solid State Fermentation	
6.1 Introduction	69

TABLE OF CONTENTS (Continued)

	Page
6.2 Objective	70
6.3 Methodology	70
6.3.1 Preparation and inoculating of substrates	70
6.3.2 Linear growth	71
6.3.3 Spore production	71
6.3.4 Percent moisture	72
6.3.5 Germination and purity of conidia	72
6.3.6 <i>In vitro</i> pathogenicity	72
6.3.7 Statistical analysis	73
6.4 Results	73
6.4.1 Linear growth of EPFs on grains	73
6.4.2 Spore production and moisture content of fungal inocula	75
6.4.3 Germination and purity of fungal inocula	76
6.4.4 Virulence of fungal inocula against fruit fly pupa	77
6.5 Discussion	77

CHAPTER 7 Screening of Cuticle Degrading Enzymes, Secondary Metabolites and Cytotoxic Effect of Crude extract from EPFs

7.1 Introduction	81
7.2 Objectives	84

TABLE OF CONTENTS (Continued)

	Page
7.3 Methodology	85
7.3.1 Detecting the chitinase activity	85
7.3.1.1 Agar plate method	85
7.3.1.2 Cultivation media	85
(a) Inoculum preparation	85
(b) Fermentation medium	85
7.3.1.3 Enzyme assay	86
7.3.2 Detecting the protease activity	86
7.3.2.1 Agar plate method	86
7.3.2.2 Preparing of inoculum and cultivation	86
7.3.2.3 Enzyme assay	87
7.3.3 Statistical analysis	88
7.3.4 Test organisms for antimicrobial activity	88
7.3.4.1 Cultivation media and methods	88
7.3.4.2 Antifungal and antibacterial assay	89
7.3.5 Preparing of crude extract for detecting DPA from selected strain, <i>M. flavoviride</i>	89
7.3.5.1 Analytical methods	90
7.3.6 Entomopathogens and cell line for <i>in vitro</i> toxicity test	90
7.3.6.1 Culture condition	90
7.3.6.2 Preparing of fungal crude	91

TABLE OF CONTENTS (Continued)

		Page
	7.3.6.3 Cell viability assay	91
7.4	Results	91
	7.4.1 Screening of cuticle-degrading enzymes by agar plate method	91
	7.4.2 Production of chitinase in different composition of medium	92
	7.4.3 Protease activity	94
	7.4.4 Determination of antimicrobial activity	95
	7.4.5 Dipicolinic acid by high performance liquid chromatography	100
	7.4.6 Cytotoxic effect	100
7.5	Discussion	101
CHAPTER 8	GENERAL DISCUSSION	106
CHAPTER 9	CONCLUSION	112
REFERENCES		115
APPENDICES		156
	APPENDIX A	157
	APPENDIX B	162
	APPENDIX C	165
	APPENDIX D	170
CURRICULUM VITAE		171

LIST OF TABLES

Table		Page
2.1	New and bioactive compounds from Thai invertebrate pathogenic fungi	11
2.2	Some pesticides that may be useful for managing spider mites in yards and gardens	28
3.1	Origin of collected entomopathogenic fungi	38
4.1	Percent mortality of dead <i>Bactrocera</i> pupae after 10 days exposure of six fungal isolates with various conidial concentration	46
4.2	Mortality of <i>Bactrocera</i> pupae infected by the fungal suspension of two species of <i>Cordyceps</i> after 10 days exposure	47
4.3	Lethal concentrations and lethal times of each tested isolates infected fruit fly, <i>Bactrocera</i> sp. for 10 days	48
5.1	Percent mortality of adult <i>Tetranychus urticae</i> Koch after 7 days exposure of six fungal isolates with various conidial concentrations	60
5.2	Percent mortality and mean number of <i>Tetranychus urticae</i> Koch after infection with the mycelial suspension of two species of <i>Cordyceps</i> 7 days post inoculation	61
5.3	Lethal concentrations and lethal times of each tested isolates infected two-spotted spider mite, <i>Tetranychus urticae</i> (Koch) 7 days post inoculation	62
5.4	Population of two-spotted spider mite <i>Tetranychus urticae</i> (Koch) 7 days after treated with different conidial concentrations	63

LIST OF TABLES (Continued)

Table	Page
6.1 Mean spore number per gram of fungal isolates incubated on various solid substrates at room temperature.	76
6.2 Percent germination of harvested dry spores on PDA, and the virulence activity of spores against fruit fly pupa (<i>Bactrocera</i> sp).	77
7.1 Screening of protease in 1% skimmed milk agar 4 days after incubation	92
7.2 Positive results of antibacterial activity of fungal extract from different cultured media against tested bacteria	97
7.3 Positive responses of antifungal activities from six entomopathogenic fungal isolates of different cultured media	99
7.4 <i>In vitro</i> cytotoxicity of the entomogenous fungal extracts from cultural broth on Vero cell after 72 hours incubation	100

LIST OF FIGURES

Figure	Page
2.1 Life cycle of fruit fly	18
2.2 Life cycle of two-spotted spider mite (<i>T. urticae</i> (Koch))	23
3.1 Collected entomopathogenic fungi from different hosts in Chiang Mai province, Thailand	31
3.2 Colony characteristic of collected entomopathogenic fungi on half-strength PDA	34
3.3 Purified PCR products of six collected insect pathogenic fungi	35
3.4 Neighbor-joining tree based on an alignment of internal transcribed spacer 1, 5.8S ribosomal RNA gene, and internal transcribed spacer 2	37
4.1 Fruit fly collection and rearing	42
4.2 Fungal mycosis of tested entomopathogenic isolates on infected fruit fly (<i>Bactrocera</i> spp.) pupae and adult	49
4.3 Daily mortality percentage of fruit fly (<i>Bactrocera</i> spp.) infecting different fungal isolates with different conidial concentration incubated for 10 days at 25°C	50
4.4 Mean percent mortality of fruit fly in different soil bioassay at a concentration of 10 ⁸ conidia ml ⁻¹ after 14 days treated	51
5.1 <i>In vitro</i> screening of fungal isolates infecting <i>T. urticae</i> by different conidial concentrations spraying on the surface of soybean leaf	57
5.2 Pot experiment	58


LIST OF FIGURES (Continued)

Figure	Page
5.3 Percent mortality of the growth stages of <i>Tetranychus urticae</i> Koch infected by six insect pathogens 7 days after inoculation with 10^8 conidia ml^{-1} . Bars represent the standard deviation of three replicated means	61
5.4 Fungal mycosis showed on the dead body of two-spotted spider mite (<i>Tetranychus urticae</i> , Koch)	62
5.5 Comparison of the mean number of alive two-spotted spider mite between treated fungal conidial suspensions and untreated control after 7 days exposure of in one-time spray trial	64
5.6 Mean number of alive <i>Tetranychus urticae</i> (Koch) in two-time spray trial 7 days after post inoculation	65
6.1 Mean linear growth of fungal isolates cultured on five grains at $25\pm 2^\circ\text{C}$ under normal day and night at 30 days after inoculation	74
6.2 Mean daily growth rate (cm) of entomopathogenic fungi on different solid substrates at room temperature ($25\pm 2^\circ\text{C}$) for 30 days	75
7.1 Chitinase activity of insect pathogenic fungi in different concentration of colloidal chitin and glucose	93
7.2 Protease activity of insect pathogenic fungi in different concentration of colloidal chitin and glucose	94
7.3 Agar well diffusion method for screening of antimicrobial activity in insect pathogenic fungi	96

ABBREVIATIONS

BCAs	Biocontrol agents/ biological control agents
BCC70	(BCC19370) <i>Cordyceps pseudomilitaris</i>
BCC91	(BCC17791) <i>Cordyceps militaris</i>
BLAST	Basic local alignment search tool
CFU	Colony forming unit
CT01	(CMUCDCT01) <i>Metarhizium flavoviride</i>
DNA	Deoxyribonucleic acid
DPA	Dipicolinic acid
EPFs	Entomopathogenic fungi
GSSM	Galetin semi-solid medium
HPLC	High performance liquid chromatography
IPM	Integrated pest management
MEGA	Molecular evolutionary genetics analysis
MF02	(CMUCDMF02) <i>Isaria tenuipes</i>
MF03	(CMUCDMF03) <i>Beauveria bassiana</i>
MF04	(CMUCDMF04) <i>Metarhizium anisopliae</i>
MG03	(CMUCDMG03) <i>Beauveria bassiana</i>
MT02	(CMUCDMT02) <i>Paecilomyces lilacinus</i> (<i>Purpureocillium lilacinum</i>)
PCR	Polymerase chain reaction
PDA	Potato dextrose agar

RH	Relative humidity
SMY	Sabouraud malt yeast extract medium
TSSM	Two -spotted spider mites



ลิขสิทธิ์มหาวิทยาลัยเชียงใหม่
Copyright© by Chiang Mai University
All rights reserved