

TABLE OF CONTENTS

	Page
ACKNOWLEDGEMENTS	iii
ABSTRACT (ENGLISH)	iv
ABSTRACT (THAI)	vi
LIST OF TABLES	ix
LIST OF FIGURES	xi
CHAPTER 1 INTRODUCTION	1
CHAPTER 2 LITERATURE REVIEW	3
CHAPTER 3 MATERIALS AND METHODS	7
CHAPTER 4 RESULTS	19
CHAPTER 5 DISCUSSION	50
CHAPTER 6 CONCLUSION	57
REFERENCES	58
CURRICULUM VITAE	66

ลิขสิทธิ์มหาวิทยาลัยเชียงใหม่
 Copyright[©] by Chiang Mai University
 All rights reserved

LIST OF TABLES

Table		Page
1	Measurements of the fine structural features of the songs of six species of <i>Pycnonotus</i> bulbuls recorded in northern Thailand. Values for each species are given as means \pm SE.	23
2	Discriminant function analysis of the songs of six species of <i>Pycnonotus</i> bulbuls produced three factors summarizing nine fine structural features of songs, represented here as the correlation coefficients between the original data and the three factors; variables with the strongest relationships (correlations > 0.60) are indicated in bold font.	27
3	Encountering rates between species showing that visually-similar species were routinely encountered based on the proportion of locations where species were encountered; highest proportions of encounter rate between paired visual similar speceis are indicated in bold font.	29
4	Discriminant function analysis of the songs of pair-wise comparisons of <i>Pycnonotus</i> bulbuls that share similar plumage features, based on three separate discriminant analyses of nine structural features of songs; variables with the strongest relationships (correlations > 0.60) are indicated in bold font.	30
5	Sexual comparisons using morphometric data of the six <i>Pycnonotus</i> bulbuls in Chiang Dao. Data shows the average \pm SE.	32
6	Discriminant function analysis of the morphometric data of six species of <i>Pycnonotus</i> bulbuls produced three factors summarizing five morphometric features, represented here as the correlation coefficients between the original data and the three factors; variables with the strongest relationships (correlations > 0.60) are indicated in bold font.	34

LIST OF TABLES (CONTINUED)

Table		
7 Nucleotide comparison shows nucleotides composition bias of excess adenine and cytosine bases, which is consistent for all the six <i>Pycnonotus</i> species. Codon positions included in this analysis were 1 st +2 nd +3 rd included gaps and missing positions.	36	
8 Pair-wise Euclidean distances of acoustic variables for all six species.	47	
9 Pair-wise Euclidean distances of morphometric variables between all six species. Euclidean distances within sex were calculated independently; males (lower diagonal) and females (upper diagonal).	47	
10 Pair-wise nucleotide composition distance using 16s mtDNA of all six species. The distance scores indicate a relatively close genetic relationship between all six species.	47	

LIST OF FIGURES

Figure	Page
1 Illustrated map showing the two study sites in Chiang Mai, northern Thailand; a) Chiang Dao Wildlife Research station, and b) Chiang Mai University (Adapted from Royal Thai Survey Department, 2002).	8
2 Climatic variations at Chiang Dao Wildlife Research Station in 2009 showing monthly rainfall (mm) and average daily minimum and maximum temperature (°C).	20
3 Sound spectrograms showing the typical song recorded from each of the six species of <i>Pycnonotus</i> bulbul in northern Thailand: a) Black-headed Bulbul, b) Black-crested Bulbul, c) Streak-eared Bulbul, d) Stripe-throated Bulbul, e) Red-whiskered Bulbul, and f) Sooty-headed Bulbul.	24
4 A scatterplot of the first three canonical discriminant functions of Acoustic features resulting from the discriminant analysis explained correctly 94.0% of variances.	26
5 Histograms showing the discriminant scores for pair-wise comparisons of <i>Pycnonotus</i> bulbuls that share similar plumage features, based on three separate discriminant analyses of nine structural features of songs.	29
6 Gel electrophoresis showing sex identification using the primers P2/P8. Two different PCR fragments were amplified from females (double bands) and males (single band).	31
7 Scatterplot of the first three canonical discriminant functions shows the results from the discriminant analysis; a) cross-validated DFA correctly explained 98.7% of variances from female morphometric features, and b) cross-validated DFA correctly explained 96.4% of variances from male morphometric features.	35

LIST OF FIGURES (CONTINUED)

Figure	Page
8 Alignment of 16s mtDNA gene sequences of all six species of the <i>Pycnonotus</i> Bulbuls. The consensus sequence is shown above the sequence alignment with nucleotide identities depicted as dots (.) and sequence deletions as dashes (-). The numbers refer to the nucleotide positions of each amplified fragment.	41
9 Hierarchical cluster analyses of the six <i>Pycnonotus</i> based on Euclidean distance of acoustic features. Spectrograms show typical song structure of each species. Dendrogram shows two discrete species' song; <i>trilled songs</i> i.e. Black-headed Bulbul and Streak-eared Bulbul, and <i>frequency-modulated tones</i> i.e. Sooty-headed Bulbul, Stripe-throated Bulbul, Red-whiskered Bulbul, and Black-crested Bulbul.	43
10 Hierarchical cluster analyses of the six <i>Pycnonotus</i> based on Euclidean distance of female morphometric features (a) and male morphometric features (b). Species clustering indicate monophyly of all six <i>Pycnonotus</i> Bulbuls with similar dendograms of both sexes.	44
11 Molecular phylogeny based on 16s mtDNA (487bp) indicated two separated groups of the <i>Pycnonotus</i> Bulbuls. Numbers above nodes indicate bootstrap (replications = 1000) of the maximum composition likelihood.	45
12 Scatter plots showing the positive correlation of acoustic distance compared with females morphometric distance (a), male morphometric distance (b), and genetic distance (c).	49