TABLE OF CONTENTS

2/52/53

ทยาลัยเชียอให

Chiang Mai Universit

Page

iii

ix

xxi

XXV

XXXV

Acknowledgements

Abstract (English)

Abstract (Thai)

List of tables

List of figures

Abbreviations and symbols

Chapter 1 General introduction and thesis outline

Chapter 2 Literature review

2.1 Orchid

2.2 Terrestrial orchid

2.3 Orchid seed

2.4 Orchid in Thailand S P e S e 13

Page

| 2.5 Mycorrhiza | 15 |
|---|----|
| 2.6 Orchid mycorrhiza | 19 |
| 2.7 Orchid mycorrhizal interactrions | 21 |
| 2.8 Isolation and identity of orchid mycorrhizal fungi | 21 |
| 2.9 Roles of mycorrhizal associations | 28 |
| 2.10 Siderophores production | 30 |
| 2.11 Plant growth hormone, Indole-3-acetic acid (IAA), production | 34 |

Chapter 3 Isolation and identification of orchid mycorrhizal fungi

| 3.1 Introduction | 39 |
|---|----------------|
| 3.2 Materials and methods | 40 |
| 3.2.1 Orchid source and root sampling | 40 19 19 |
| 3.2.2 Orchid roots observation and mycorrhizal colonization | ⁴¹ |

Page

| 3.2.3 Fungal isolation | 41 |
|---|----|
| 3.2.4 Fungal identification | 45 |
| 3.2.4.1 Morphological observation | 45 |
| 3.2.4.2 Molecular analysis | 45 |
| 3.3 Results | 47 |
| 3.3.1 Orchid roots observation and mycorrhizal colonization | 47 |
| 3.3.2 Fungal isolation and identification | 51 |
| 3.4 Discussion | 59 |

Chapter 4 Effects of orchid mycorrhizal fungi on symbiotic seed

ng1988188162

reser⁶⁴

germination

4.1 Introduction

4.2 Materials and methods

Chiang Mai Uni⁶⁴ersit 4.2.1 Orchid seed collection

Page

| 4.2.2 | Orchid seed viability test | 64 |
|-------------|---|----|
| 4.2.3 | Symbiotic seed germination study of <i>P.susannae</i> | 65 |
| | 4.2.3.1 Effects of agar media on symbiotic seed | 65 |
| | germination | |
| | 4.2.3.2 Effects of endophytic fungi recovered from P. | 66 |
| | susannae on symbiotic seed germination | |
| | 4.2.3.3 Effects of mycorrhizal fungi on symbiotic seed | 67 |
| | germination | |
| 4.3 Results | | 71 |
| 4.3.1 | Symbiotic seed germination of <i>P. susannae</i> | 71 |
| | 4.3.1.1 Effects of agar media on symbiotic seed | 71 |
| | germination | |
| | 4.3.1.2 Effects of endophytic fungi recovered from <i>P</i> . | 73 |
| | susannae on symbiotic seed germination | |
| | | |

| | Page |
|--|---------------------------|
| | |
| 4.3.1.3 Effects of mycorrhizal fungi on symbiotic seed | 76 |
| germination | |
| 4.4 Discussion | 80 |
| | |
| Chapter 5 Screening for indole-3-acetic acid (IAA) and | |
| siderophores production by endophytic fungi | |
| 5.1 Introduction | 85 |
| 5.2 Materials and methods | 87 |
| 5.2.1 Colorimetric assay for IAA determination | 87 |
| 5.2.2 Effect of L-tryptophan concentration on IAA production | 88 |
| 5.2.3 Effect of incubation period on IAA production | 88 |
| | |
| 5.2.4 Extraction of fungal IAA and thin layer chromatography | 88 |
| 5.2.5 Evaluation of IAA biological activity | ⁸⁹ niversit |
| 5.2.5.1 Kidney bean root cutting | 89 |
| | |

Page

| 5.2.5.2 Corn seed germination | 90 |
|--|-----------------------------------|
| 5.2.5.3 Elongation of rice coleoptiles | 90 |
| 5.2.6 Siderophores production | 91 |
| 5.2.6.1 Screening for siderophores production | 91 |
| 5.2.6.2 Chemical determination of produced siderophores | 91 |
| 5.2.7 Statistic analysis | 92 |
| 5.3 Results | 92 |
| 5.3.1 Determination of fungal IAA | 92 |
| 5.3.2 Effect of L-tryptophan concentration on IAA production | 94 |
| 5.3.3 IAA extraction and thin layer chromatography | 97 |
| 5.3.4 Evaluation of IAA biological activities of fungal IAA | 97 |
| 5.3.4.1 Kidney bean root cutting | 8 97 In 1 |
| 5.3.4.2 Corn seed germination | ni 99 ersit |
| 5.3.4.3 Elongation of rice coleoptiles | $\mathbf{\Gamma}^{101}\mathbf{O}$ |

| | Page |
|---|------|
| | |
| 5.3.4 Detection for siderophores production | 102 |
| 5.4 Discussion | 105 |
| | |
| | |
| Chapter 6 Optimization of orchid mycorrhizal fungi on media | |
| 6.1 Introduction | 109 |
| 6.2 Materials and methods | 110 |
| 6.2.1 Effects of temperature on fungal growth on PDA | 110 |
| 6.2.2 Effects of pH on fungal growth on PDA | 110 |
| 6.2.3 Evaluation of suitable grain and potting media for fungal | 112 |
| inoculum production | |
| 6.3 Results | 113 |
| 6.3.1 Effects of temperature on fungal growth on PDA | 113 |
| | 118 |

| Pa | ge |
|----|-----|
| | SC. |

| | Page |
|---|------|
| 6.3.3 Evaluation of suitable grain and potting media for fungal | 122 |
| inoculum production | |
| 6.4 Discussion | 124 |
| | |
| Chapter 7 Mycorrhizal colonization and effects of mycorrhizal | |
| fungi on growth of orchid seedlings | |
| 7.1 Introduction | 127 |
| 7.2 Materials and methods | 129 |
| 7.2.1 Fungal inoculum | 129 |
| 7.2.2 Symbiotic cultivation of orchid seedlings | 129 |
| 7.2.3 Evaluation of symbiotic orchid seedlings | 130 |
| 7.2.4 Determination of mycorrhizal colonization | -131 |
| 7.2.5 Statistic analysis | 132 |
| 7.3 Results | 132 |
| rights rese | rv |

| | Page |
|---|-----------------------|
| | |
| 7.3.1 Evaluation of symbiotic orchid seedlings | 132 |
| 7.3.2 Determination of mycorrhizal colonization | 132 |
| 7.4 Discussion | 135 |
| | |
| Chapter 8 General discussion | 138 |
| | A |
| | |
| References | 146 |
| Appendices | 166 |
| Appendix A | 167 |
| Appendix B | 171 |
| Appendix C | 173 |
| Appendix D | 175 |
| Appendix E by Chiang Mai U | n ¹⁷⁷ ersi |
| Curriculum Vitae | 1 ⁷⁹ e |
| | |

LIST OF TABLES

Table

| 2.1 Seed germination and protocorm developmental stages use for | 12 |
|--|-----|
| determination of growth and development of terrestrial orchid in | |
| vitro | |
| | |
| 2.2 Some terrestrial orchid species in Thailand | 14 |
| 2.3 Types of mycorrhizal associatuions | 18 |
| | |
| 2.4 The major members of orchid mycorrhizal fungi | 20 |
| 2.5 Classification of the Rhizoctonia-forming heterobasidiomycetes | 26 |
| | |
| 2.6 Types of interspecies antagonisms leading to biological control | 33 |
| of plant pathogens | |
| 2.7 Predominant siderophore types in major mycorrhizal types | 34 |
| 2.7 Tredominant siderophore types in major mycormizar types | 54 |
| 2.8 Several microorganisms which are able to converted L- | 37 |
| tryptophan to indole-3-acetic acid (IAA) | |
| | 912 |
| 3.1 Collection site and time of collection of six terrestrial orchid | 4/ |
| samples | |
| 3.2 Endophytic fungi isolated from roots of six terrestrial orchids | 53 |
| rights rese | |

LIST OF TABLES (CONTINUED)

Table

Page

68

68

69

70

73

78

4.1 Mycorrhizal fungi isolated from roots of *Pecteilis susannae* and used to determined effects of agar media on symbiotic seed germination

4.2 Seed germination and protocorm developmental stages use for determination of growth and development of *Pecteilis susannae in vitro*

4.3 Endophytic fungi isolated from roots of *Pecteilis susannae* and used in symbiotic seed germination

4.4 Mycorrhizal fungi isolated from roots of *Pecteilis susannae*,*Eulophia spectabilis*, *Paphiopedilum*, and *Spathoglottis affinis* andused in symbiotic seed germination

4.5 Effects of 3 agar media (Oatmeal agar, Cellulose agar and Vacin and Went agar) on symbiotic seed germination and protocorm development of *Pecteilis susannae* 70 days after sowing

4.6 Effects of 11 endophytic fungi on seed germination and protocorm development of *Pecteilis susannae* 133 days after sowing

LIST OF TABLES (CONTINUED)

Table

112

5.1 Selected endophytic fungi isolated from root of Thai native
92 terrestrial orchids, *Spathoglottis affinis, Paphiopedelum bellatulum*and *Phaius tankervilleae*5.2 IAA production by endophytic fungi isolated from root of six
93 terrestrial orchids
5.3 Effect of IAA produced by *Colletotrichum gloeosporioides*101
CMU-AU 006, *Epulorhiza* sp. CMU-SLP 007 and *Epulorhiza* sp.
CMU-NUT 013 on root to shoot ratio of corn
5.4 Siderophores production by endophytic fungi isolated from root

of six terrestrial orchids

6.1 Orchid mycorrhizal fungi isolated from roots of six terrestrial111orchids and used in optimization of fungal growth temperature andpH

6.2 Grain and potting media used for evaluation of suitable material for fungal inoculum production

LIST OF TABLES (CONTINUED)

Table

119

133

176

178

6.3 The optimal growth pH evaluated from fungal dry weight and fungal colony diameter of orchid mycorrhizal fungi isolated from roots of six terrestrial orchids after cultured on PDA for 10 days

7.1 Effects of mycorrhizal inoculum of fungal isolates CMU-DP 506, *Epulorhiza* sp., and CMU-DP 514, *Tulasnella* sp., used for potting media on growth and survival rate of *Doritis pulcherrima* seedlings after cultivation for 90 days

D Absorbance at 530 nm of various concentration of standard IAA

E Absorbance at 480 nm of various concentration of deferoxamine mesyilate by iron-perchlorate assay

ลิ<mark>ปสิทธิ์มหาวิทยาลัยเชียงใหม่</mark> Copyright[©] by Chiang Mai University All rights reserved

LIST OF FIGURES

Page

Figure

| 1.1 Structure of the thesis | 4 |
|--|--------------|
| 2.1 The schematic overview of epiphytic orchid and terrestrial orchid | 7 |
| 2.2 Orchid capsule and seed, the mature capsule of <i>Calanthe tricarina</i> , seed of <i>Calanthe tricarina</i> , and seed of <i>Liparis loeselii</i> | |
| 2.3 Growth stage used to determine germination and development of orchid seed <i>in vitro</i> | 12 |
| 2.4 Mycorrhizal associations; the interaction of plant roots, soil (nutrients in soil) and fungi (Brundrett <i>et al.</i> 1996) | 16 |
| 2.5 A schematic overview of the different forms of mycorrhizas | 17 |
| 2.6 The coils of fungal hyphae, pelotons, of orchid mycorrhizal fungi within cortex cells of orchid roots | 20 |
| 2.7 Drawing of a typical orchid mycorrhizal fungi, <i>Rhizoctonia</i> repen (Tulasnella calospora), isolated in pure culture | 8 23 nive |
| | |

Figure Page 2.8 Internal transcribed spacer (ITS) regions and recognized sites 27 of ITS primers and Large subunit (LSU) RNA and recognized sites of LSU primers 2.9 Mechanism of iron acquisition by microbial siderophore 32 2.10 Pathways for the microbial biosynthesis of indole-3-acetic 36 acid (IAA) from tryptophan 3.1 Collection sites of terrestrial orchid samples 43 3.2 Location of Chiang Mai province, in northern part 44 of Thailand 3.3 Doritis pulcherrima 48 48 3.4 Eulophia spectabilis 3.5 Paphiopedilum bellatulum 49 3.6 Pecteilis susannae 50 3.7 Phaius tankervilleae 3.8 Spathoglottis affinis

Figure

Page

52

56

57

58

65

72

3.9 Isolation of fungal endophytes from roots of *Pecteilis susannae*3.10 Percentages of the occurrence of endophytic fungal taxaisolated from roots of six terrestrial orchids

3.11 Epulorhiza isolates

3.12 Maximum-parsimony trees based on an alignment of internal transcribed spacer1, 5.8S ribosomal RNA gene and internal transcribed spacer 2 of 19 fungal endophytes isolated from roots of six terrestrial orchids

4.1 *Pecteilis susannae* in natural habitat (site 2: Sa-Lunag, Chiang Mai, Thailand)

4.2 Viable seed with the embryo stained red and hyaline embryo of dead seed of *Pecteilis susannae* after being treated with tetrazolium

4.3 Effects of 8 endophytic fungi on seed germination and protocorm development of *Pecteilis susannae* 70 days after sowing

Figure

Page

76

79

95

4.4 Symbiotic seed germination and protocorm developmental stages of *Pecteilis susannae* using fungal endophyte *Epulorhiza* sp., CMU-AUG 013, cultured on Oatmeal agar 70 days after sowing

4.5 Symbiotic seed germination and protocorm developmental stages of *Pecteilis susannae* using mycorrhizal fungi *Epulorhiza* sp., CMU-AUG 007, cultured on Oatmeal agar 133 days after sowing

4.6 Effects of 11 endophytic fungi on seed germination and protocorm development of *Pecteilis susannae* 133 days after sowing

5.1 Effect of L-tryptophan concentration on IAA production by*Colletotrichum gloeosporioides* CMU-AU 006, *Epulorhiza* sp.CMU-SLP 007 and *Epulorhiza* sp. CMU-NUT 013

5.2 Effect of incubation period on IAA production by Colletotrichum gloeosporioides CMU-AU 006

| Figure | Page |
|---|------|
| 5.3 Effect of incubation period on IAA production by <i>Epulorhiza</i> sp. CMU-SLP 007 | 96 |
| 5.4 Effect of incubation period on IAA production by <i>Epulorhiza</i> sp. CMU-NUT 013 | 97 |
| 5.5 Effect of IAA produced by <i>Colletotrichum gloeosporioides</i> CMU-AU 006, <i>Epulorhiza</i> sp. CMU-SLP 007 and <i>Epulorhiza</i> sp. | 98 |
| CMU-NUT 013 on root formation of kidney bean root cutting | 6 |
| 5.6 Effect of IAA produced by <i>Colletotrichum gloeosporioides</i> CMU-AU 006, <i>Epulorhiza</i> sp. CMU-SLP 007 and <i>Epulorhiza</i> sp. | 100 |
| CMU-NUT 013 on corn seed germination and the length of root5.7 Effect of IAA produced by <i>Colletotrichum gloeosporioides</i> | 102 |
| CMU-AU 006, <i>Epulorhiza</i> sp. CMU-SLP 007 and <i>Epulorhiza</i> sp. CMU-NUT 013 on elongation of rice coleoptiles | |
| 5.8 The Chrome Azurol S (CAS) agar plate assay showing the | 105 |
| purplish-red and yellow halo after inoculated with the active fungal isolates for 14 days | |
| | |

Figure

Page

113

114

114

6.1 Effects of temperature on growth of fungal isolate CMU-AUG028, *Epulorhiza* sp., after culturing on PDA and incubation at 20°Cfor 10 days

6.2 Effects of temperature on growth (dry weight) of orchid mycorrhizal fungi isolated from roots of *Pecteilis susannae* after cultured on PDA for 10 days

6.3 Effects of temperature on growth (colony diameter) of orchid mycorrhizal fungi isolated from roots of *Pecteilis susannae* after cultured on PDA for 10 days

6.4 Effects of temperature on growth (dry weight) of orchid115mycorrhizal fungi isolated from roots of *Eulophia spectabilis* aftercultured on PDA for 10 days

6.5 Effects of temperature on growth (colony diameter) of orchid115mycorrhizal fungi isolated from roots of *Eulophia spectabilis* aftercultured on PDA for 10 days

Figure

Page

116

116

118

6.6 Effects of temperature on growth (dry weight) of orchid mycorrhizal fungi isolated from roots of *Paphiopedilum bellatullum* and *Phaius tankervilleae* after cultured on PDA for 10 days

6.7 Effects of temperature on growth (colony diameter) of orchid mycorrhizal fungi isolated from roots of *Paphiopedilum bellatullum* and *Phaius tankervilleae* after cultured on PDA for 10 days

6.8 Effects of temperature on growth (dry weight) of orchid 117 mycorrhizal fungi isolated from roots of *Doritis pulcherrima* and *Spathoglottis affinis* after cultured on PDA for 10 days

6.9 Effects of temperature on growth (colony diameter) of orchid117mycorrhizal fungi isolated from roots of *Doritis pulcherrima* andSpathoglottis affinis after cultured on PDA for 10 days

6.10 Effects of pH on growth of fungal isolate CMU-AUG 028, *Epulorhiza* sp.

Figure

Page

120

121

122

123

6.11 Effects of pH on growth (dry weight) of orchid mycorrhizalfungi isolated from roots of six terrestrial orchids after culturing onPDA for 10 days

6.12 Effects of pH on growth (colony diameter) of orchid mycorrhizal fungi isolated from roots of six terrestrial orchids after culturing on PDA for 10 days

6.13 Mycelium growth of orchid mycorhizal fungi, *Epulorhiza* sp. (CMU-AUG 007), isolated from roots of *Pecteilis susanne* in grain and potting media after incubation at 30°C for 15 days

6.14 Mycelium growth of orchid mycorhizal fungi, *Epulorhiza* sp.
123 (CMU-AUG 028), isolated from roots of *Pectelis susannae* in grain and potting media after incubation at 30°C for 15 days

6.15 Mycelium growth of orchid mycorhizal fungi, *Epulorhiza* sp. (CMU-AUG 014), isolated from roots of *Eulophia spectabilis* in grain and potting media after incubation at 30°C for 15 days

Figure

Page

124

130

133

6.16 Growth of mycelium of orchid mycorrhizal fungi, *Epulorhiza* sp. (CMU-AUG 028), isolated from roots of *Pectelis susannae*, in grain and potting media

7.1 Orchid mycorrhizal inocula of fungal isolates CMU-DP 506, *Epulorhiza* sp. and CMU-DP 514, *Tulasnella* sp.

7.2 Seedlings of *Doritis pulcherrima* after cultivation using non mycorhizal potting medium, potting medium with fungal isolate CMU-DP 506, *Epulorhiza* sp. and CMU-DP 514, *Tulasnella* sp. for 90 days

7.3 Peloton structures, coiled hyphae, in cortical cells of Doritis134pulcherrima seedlings

7.4 Root and crop colonization by mycorhizal fungi of Doritis135pulcherrima seedlings after cultivation using potting media mixed135with mycorrhizal inoculum of CMU-DP 506, Epulorhiza sp., andCMU-DP 514, Tulasnella sp., for 90 days

D Standard cruve of IAA concentration against absorbance at 530 nm

Figure

Page

178

E Standard curve for hydroxamate concentration determination

using deferoxamine mesylate as standard

ลิ<mark>ปสิทธิ์มหาวิทยาลัยเชียงใหม่</mark> Copyright[©] by Chiang Mai University All rights reserved

xxxiv

ABBREVIATIONS AND SYMBOLS

| | NHEIKA ? |
|--------------------|---|
| АМ | arbuscular mycorrhiza |
| anam. | anamorph |
| Asco- | Ascomycetes |
| Basidio- | Basidiomycetes |
| cm | centimeter |
| DNA | deoxyribonucleic acid |
| Е | embryo |
| ECM | ectomycorrhiza |
| g | gram |
| IAA | indole-3-acetic acid |
| ITS | internal transcribed spacer |
| L | liter |
| LSU | large subunit |
| DDVP ^{mg} | O b ^{milligram} hiang Mai University |
| min | ignts reserved |

xxxvi

| ml | milliliter |
|--------------|---|
| mm | millimeter |
| М | molar (mol per liter) |
| mM | milli Molar |
| MW | molecular weight |
| n | number |
| nm | nanometer |
| No. | number |
| Р | probability |
| PCR | Polymerase Chain Reaction |
| PDA | Potato Dextrose Agar |
| PDB | Potato Dextrose Broth |
| ppm | part per million |
| rh | rhizosphere soil |
| rtSUK | root ng |
| sec | second Chiang Mai University |
| TLC | thin layer chromatography |
| | |

xxxvii

| rpm | rotation per minute |
|------|---------------------------------|
| UV | ultraviolet |
| VAM | vesicular-arbuscular mycorrhiza |
| v/v | volume by volume |
| w/v | weight by volume |
| w/w | weight by weight |
| °C | degree Celsius |
| μg | microgram |
| μΙ | microlitre |
| μm | micrometer |
| μmol | micromole |
| | |

ลิ<mark>ปสิทธิ์มหาวิทยาลัยเชียงใหม่</mark> Copyright[©] by Chiang Mai University All rights reserved