

TABLE OF CONTENTS

	Page
ACKNOWLEDGMENTS	iii
ABSTRACT (ENGLISH)	v
ABSTRACT (THAI)	viii
ABBREVIATIONS AND SYMBOLS	xx
CHAPTER 1 INTRODUCTION	1
1.1 The statement and significance of the problem	1
1.1.1 Pesticides in Thailand	1
1.1.2 Organophosphate and carbamate pesticide statement and significance	3
1.1.3 Synthetic pyrethroid pesticide statement and significance	6
1.2 Research Objectives	9
1.3 Definitions	11
CHAPTER 2 LITERATURE REVIEWS	13
2.1 Classification of pesticides	13
2.2 Non-persistent pesticides	13
2.2.1 Organophosphate pesticides poisoning	14
2.2.2 Carbamate pesticides poisoning	17
2.2.3 The synthetic pyrethroid characterize	17

TABLE OF CONTENTS (CONT.)

	Page
2.3 Biological monitoring of pesticide exposure	24
2.3.1 Blood and salivary cholinesterase enzyme	24
2.3.2 Plasma 3-PBA and urine 3-PBA	28
2.4 Method for measurement of biomarkers	33
2.4.1 Cholinesterase enzyme activity measurement	33
2.4.2 Enzyme-linked immunosorbent assay (ELISA)	34
2.4.2.1 The step of ELISA development	35
2.4.2.2 Design and synthesis of hapten	35
2.4.2.3 Preparation of immunogen and coating antigen	36
2.4.2.4 Production of antibody	37
2.4.2.5 Immunoassay formats	41
CHAPTER 3 EXPERIMENTAL	45
3.1 Materials and Chemicals	45
3.1.1 Chemicals and reagents	45
3.1.1.1 Chemicals	45
3.1.1.2 Immunological reagents	46
3.1.1.3 Disposable products	46
3.1.2 Equipments	47
3.2 Methods	47

TABLE OF CONTENTS (CONT.)

	Page
3.2.2.2.4.2 Competitive indirect enzyme-linked immunosorbent assay	59
3.2.2.3 Quality assurance and control	60
3.2.2.4 Statistic analysis	60
3.2.2.5 High-performance liquid chromatography /time-of-flight mass spectrometry (HPLC/TOFMS)	61
3.2.2.5.1 High resolution mass spectrometry	61
3.2.2.5.2 Ion source conditions	61
3.2.2.5.3 TOF calibration and lock-mass set up	62
3.2.2.5.4 HPLC separation	63
3.2.2.5.5 Mobile phase, gradient	63
3.2.2.5.6 Column	63
3.2.2.5.7 UV-VIS detector	63
3.2.2.5.8 Data processing	63
CHAPTERS 4 RESULTS AND DISCUSSION	64
4.1 Organophosphate and carbamate pesticide results and discussion	64
4.1.1 Developing of human salivary cholinesterase assay	64
4.1.2 Application of developed methods to detect activities of BChE and AChE in saliva	64

TABLE OF CONTENTS (CONT.)

	Page
4.1.3 Examination of the relationship between activities of BChE and AChE in saliva and blood	67
4.2 Synthetic pyrethroid pesticide	69
4.2.1 Developing of sample preparation step and sensitivity ELISA method	69
4.2.1.1 Sample preparation	69
4.2.1.2 Optimization of the ELISA method	74
4.2.1.3 Quality assurance and control	76
4.2.2 Application developed methods to detect plasma 3-PBA	76
4.2.3 Examination the relationship between plasma 3-PBA and urine 3-PBA	79
CHAPTER 5 CONCLUSION	82
5.1 Organophosphate and Carbamate pesticides	82
5.2 Synthetic pyrethroid pesticides	83
REFERENCES	84
APPENDIX	104
CURRICULUM VITAE	109

LIST OF TABLES

Table	Page
2.1 Classification of major pesticides according to persistence into environment	13
2.2 Symptoms and sign of OP pesticide poisoning	14
2.3 Acute oral toxicities of pyrethroids to rats	21
3.1 Participant distribution through 4 sub-districts in the Fang district of Chiang Mai, Thailand	48
3.2 The optimal condition for ChE measuring in saliva and blood	54
4.1 The within-day %CV for pooled saliva and blood in each batch (3 controls; intra-assay) and the between-day %CV of control concentrations in all batches (60 controls; intra-assay) for 200 samples	65
4.2 Limit of blank (LOB), limit of detection (LOD) and limit of quantitation (LOQ) of methods for measurement of ChE activities	66
4.3 Factors affecting variability from the status of diseases to activity of all ChE	70
4.4 Recoveries after hydrolysis, LLE, SPE and ELISA analysis of spiked free-3-PBA and 3-PBA-BSA (10 ng/mL) in fetal bovine serum, done in duplicate in two days.	72

LIST OF TABLES (CONT.)

Table	Page
4.5 Retention time (RT), found mass, mass increment and number of 3-PBA to BSA for characterization of 3-PBA-BSA (Molecular mass for calculation in this study of BSA and 3-PBA are 66,433Da and 196Da, respectively)	72
4.6 Recoveries after hydrolysis, extraction and ELISA analysis of 3-PBA spiked in fetal bovine serum and urine containing no detectable 3-PBA by ELISA in duplicate (n=2) on the same day.	73
4.7 Limit of blank (LOB), limit of detection (LOD) and limit of quantitation (LOQ) of methods for measurement of plasma 3-PBA and urinary 3-PBA	74
4.8 Intra- and inter-assay variation of pooled controls for plasma and urine	77
4.9 Descriptive analytical data for plasma 3-PBA and urinary 3-PBA detected in human plasma between consumers (n=100) and farmers (n=100)	78
4.10 The % detection of plasma 3-PBA and urinary 3-PBA in consumer and farmer groups by concentration ranges	80

LIST OF FIGURES

Figure	Page
1.1 Conceptual framework of the present study	10
2.1 Typical structures of OP pesticides	15
2.2 Metabolic pathway of methyl parathion in the body representative of organophosphate pesticide metabolism (Barr and Needham, 2002).	16
2.3 Structures of the six natural pyrethrins (Soderlund <i>et al.</i> , 2002)	18
2.4 Structures of synthetic pyrethroids registered for use in the United States	19
2.5 Schematic structure of cholinesterase active site of human PChE monomer (Ekholm and Konschin, 1999; Masson <i>et al.</i> , 2001; 2003)	26
2.6 Metabolism of permethrin in mammals	30
2.7 Adduct formation of acyl glucuronides with proteins by transacylation mechanism	31
2.8 Adduct formation of acyl glucuronides via glycation mechanism	31
2.9 (A) Structure of IgG antibody and its fragments. V_L is the variable regions of light chain, V_H is the variable regions of heavy chain, C_L is the constant regions on light chain, and C_{H1} , C_{H2} , C_{H3} are the constant regions on heavy chain. (B) Fragments contained the antigen binding sites such as F(ab), F(ab') and F(ab') ₂ (Abcam, 2012).	38

LIST OF FIGURES (CONT.)

Figure	Page
2.10 The principle of antibody production. B cell are triggered to mature into plasma cells that produce a specific kind of antibody when the B cell encounters a specific antigen (www.niaid.nih.gov)	40
2.11 ELISA format (www.thermoscientific.com)	42
3.1 Bivariate scattergram of SChE and SBChE activities between using the spitted samples and the cotton wool-collected samples were significantly correlated for SChE ($r = 0.763$, $P < 0.01$, $n = 20$) and SBChE ($r = 0.803$, $P < 0.01$, $n = 20$).	51
3.2 Forms of 3-PBA in the plasma sample during alkaline hydrolysis and the LLE method	56
3.3 The target molecule (3-PBA) linked to dual mode SPE (C8 and anionic exchange) cartridge	57
3.4 Comparison between 3-PBA (A) and hapten (B) in this study (Shan <i>et al.</i> , 2004)	58
4.1 Bivariate scattergram of SBChE activities (mU/mL) and PBChE activities (U/mL) found correlation ($r = 0.232$, $P = 0.020$) between activities of SBChE and PBChE in consumers.	68
4.2 ELISA inhibition curve for plasma 3-PBA (A) and urine 3-PBA (B) using antiserum 294 (diluted 1 : 7,000, final dilution in well), coating antigen 3-PBA-BSA (0.5 $\mu\text{g}/\text{mL}$), and GAR-HRP (1 : 10,000)	75

LIST OF FIGURES (CONT.)

Figure	Page
4.3 The concentrations of urinary 3-PBA in farmers in Chiang Mai province at Mae Rim district (N = 40; Panuwet <i>et al.</i> (2004) and Mae Taeng District (N = 69; Panuwet <i>et al.</i> (2008) at selected percentile values compared to the present study (Fang district).	81

ABBREVIATIONS AND SYMBOLS

ng/mL	Nanogram
°C	Degree celsius
mL	Milliliter
g	Gram
Geo-mean	Geometric mean
Mg	Milligram
Kg	Kilogram
Max	Maximum
Min	Minimum
N	Sample size
SD	Standard deviation
U.S.A.	United States of America
WHO	World Health Organization
U.S.EPA	United States Environmental Protection Agency