

TABLE OF CONTENTS

	Page
ACKNOWLEDGEMENT	iii
ABSTRACT	v
TABLE OF CONTENTS	x
LIST OF TABLES	xiv
LIST OF FIGURES	xv
ABBREVIATIONS	xvii
CHAPTER 1 INTRODUCTION	
1.1 Statement of problems	1
1.2 Literature review	4
1.2.1 Cancer	4
1.2.2 Lung cancer	6
1.2.3 p53 tumor suppressor gene	8
1.2.4 Response of p53 to stress signals	11
1.2.5 The p53 tumor suppressor gene family	15
1.2.6 Antibodies and the analysis of p53 protein	16
1.2.7 Our previous study involved searching for a p53 structurally-related protein	18

1.3	Objectives	21
CHAPTER 2 MATERIALS AND METHODS		
2.1	Materials	22
2.1.1	Specimens	22
2.1.2	Instruments	22
2.1.3	Chemical and Reagents	22
2.1.4	Other materials	22
2.2	Methods	23
2.2.1	Preparation of cell lysates from lung tissues	23
2.2.2	Identification of the protein by immunoprecipitation	24
2.2.2.1	Purification of protein by immunoprecipitation	24
2.2.2.2	Protein identification by LC-MS/MS (Liquid chromatography-tandem mass spectrometry)	25
2.2.2.3	Protein determination by Western blot analysis	25
2.2.3	Characterization the CM5pAb reactive protein	31
2.2.3.1	Cell and culture condition	31
2.2.3.2	Investigation of genotoxic stress response	31
2.2.3.3	Investigation of ER stress response	32
2.2.3.4	Investigation of protein sub-cellular localization in response to stress signal	33

CHAPTER 3 RESULTS

3.1	Identification the protein reacted to CM5pAb	34
3.1.1	Protein purification by immunoprecipitation with CM5pAb	34
3.1.2	Identification of immunoprecipitated proteins by LC-MS/MS (Liquid chromatography-tandem mass spectrometry)	35
3.1.3	Verification and confirmation of the identification result from LC-MS/MS	42
3.2	Expression pattern of CM5pAb reactive protein and glucosidase II in human lung tumor tissues	45
3.3	Characterization the CM5pAb reactive protein (glucosidase II) in response to stress signals compared with p53 protein expression when response to stress in cell culture model	47
3.3.1	Expression and cellular localization of p53 and glucosidase II in response to UV irradiation	47
3.3.2	Expression and cellular localization of p53 and glucosidase II in response to tunicamycin-induced ER stress	53

CHAPTER 4 DISCUSSION AND CONCLUSION

4.1	Discussion	58
-----	------------	----

4.2 Conclusion	62
REFERENCES	63
APPENDICES	73
APPENDIX A	74
APPENDIX B	75
APPENDIX C	78
CURRICULUM VITAE	83



ลิขสิทธิ์มหาวิทยาลัยเชียงใหม่

Copyright© by Chiang Mai University
All rights reserved

LIST OF TABLES

Table		Page
2.1	List of antibodies and dilution used in this study	29
3.1	LC-MS/MS result of immunoprecipitated protein band A	36
3.2	LC-MS/MS result of immunoprecipitated protein band B	38
3.3	LC-MS/MS result of immunoprecipitated protein band C	40
3.4	Expression of glucosidase II in relation to clinicopathologic features of tumor tissues	46

LIST OF FIGURES

Figure	Page
1.1 Regulation of p53	13
1.2 Expression of CM-5 reactive proteins and p53 protein in lung tissues	20
3.1 Coomassie blue stained gel showing the CM5Ab immunoprecipitated protein bands	34
3.2 Comparison of protein molecular weight of the CM5 reactive protein with GRP78 and Glucosidase II	43
3.3 Western blot analysis showing the reactivity of the immunoprecipitated glucosidase II to CM5pAb	44
3.4 Western blotting results showing expression similarity of CM5 reactive protein and glucosidase II in lung tumor and normal corresponding tissues	45
3.5 A549 cell morphology after UV treatment	49
3.6 Western blot showing protein levels of p53 and glucosidase II after irradiated with various dose of UV	50
3.7 Western blot analysis and quantification value showing changes of various protein levels in A549 lung adenocarcinoma cells at various time points following UV irradiation	51

3.8	Western blot analysis showing sub-cellular localization of p53 and glucosidase II in response to UV irradiation	52
3.9	A549 cell morphology after tunicamycin treatment	54
3.10	Western blot analysis showing protein levels of p53 and glucosidase II after tunicamycin treatment	55
3.11	Western blot analysis and quantification value showing changes of various protein levels in A549 lung adenocarcinoma cells at various time points following tunicamycin treatment	56
3.12	Western blot analysis showing sub-cellular localization of p53 and glucosidase II in response to tunicamycin-induced ER stress	57