

REFERENCES

- Abdelkader, H., Ismail, S., Kamal, A., Alany, R. G. 2011. Design and evaluation of controlled-release niosomes and discomes for naltrexone hydrochloride ocular delivery. *J. Pharm. Sci.*, 100(5): 1833-1846.
- Abramson, H. A., Gorin, M. H. 1940. Skin Reactions. IX. The electrophoretic demonstration of the patent pores of the living human skin; its relation to the charge of the skin. *J. Phys. Chem.*, 44(9): 1094-1102.
- Aburai, K., Yoshino, S., Sakai, K., Sakai, H., Abe, M., Loiseau, N., Holleran, W., Uchida, Y. and Sakamoto, K. 2011. Physicochemical analysis of liposome membranes consisting of model lipids in the stratum corneum. *J. Oleo Sci.*, 60: 197-202.
- Abuzaytoun, R., Shahidi, F. 2006. Oxidative stability of flax and hemp oils. *J. Am. Oil Chem. Soc.*, 83: 855-861.
- Adhirajan, N., Ravi Kumar, T., Shanmugasundaram, N., Babu, M. 2003. *In vivo* and *in vitro* evaluation of hair growth potential of *Hibiscus rosa-sinensis* Linn. *J. Ethnopharmacol.*, 88: 235-239.
- Agarwal, R., Katare, O. P., Vyas, S. P. 2000. The pilosebaceous unit: A pivotal route for topical drug delivery. *Methods Find. Exp. Clin. Pharmacol.*, 22: 129-133.
- Aggarwal, D., Kaur, I. P. 2005. Improved pharmacodynamics of timolol maleate from a mucoadhesive niosomal ophthalmic drug delivery system. *Int. J. Pharm.*, 290: 155-159.
- Ajayi, I. A., Ajibade, O., Oderinde, R. A. 2011. Preliminary phytochemical analysis of some plant seeds. *Res. J. Chem. Sci.*, 1(3): 58-62.

Akihisa, T., Yasukawa, K., Oinuma, H., Kasahara, Y., Yamanouchi, S., Takido, M., Kumaki, K., Tamura, T. 1996. Triterpene alcohols from the flowers of compositae and their anti-inflammatory effects. *Phytochemistry*, 43: 1255-1260

Alencar, J., Gosset, G. l., Robin, M., Pique, V. r., Culcasi, M., Clement, J.-L., Mercier, A., Pietri, S. 2009. Improving the stability and antioxidant properties of sesame oil: water-soluble spray-dried emulsions from new transesterified phenolic derivatives. *J. Agric. Food Chem.*, 57(16): 7311-7323.

Alexeev, V., Igoucheva, O., Domashenko, A., Cotsarelis, G., Yoon, K. 2000. Localized *in vivo* genotypic and phenotypic correction of the albino mutation in skin by RNA-DNA oligonucleotide. *Nat. Biotechnol.*, 18: 43-47.

Allen, S. T. 1974. Chemical analysis of ecological material. Blackwell scientific publication, New York.

Alvarez-Roman, R., Naik, A., Kalia, Y. N., Guy, R. H., Fessi, H. 2004. Skin penetration and distribution of polymeric nanoparticles. *J. Controlled Release*, 99(1): 53-62.

Andjelkovic, M., Van Camp, J., De Meulenaer, B., Depaemelaere, G., Socaciu, C., Verloo, M., Verhe, R. 2006. Iron-chelation properties of phenolic acids bearing catechol and galloyl groups. *Food Chem.*, 98(1): 23-31.

AOAC 2000. (996.01) Fat (total, saturated, unsaturated, and monounsaturated) in cereal products Official methods of analysis. vol. 17th ed AOAC international, Maryland, U.S.A.

- Arunothayanun, P., Sooksawate, T., Florence, A. T. 1999. Extrusion of niosomes from capillaries: approaches to a pulsed delivery device. *J. Controlled Release*, 60: 391-397.
- Attia, I., El-Gizawy, S., Fouda, M., Donia, A. 2007. Influence of a niosomal formulation on the oral bioavailability of acyclovir in rabbits. *AAPS Pharm. Sci.*, 8(4): 206-212.
- Aungst, B. J., Blake, J. A., Hussain, M. A. 1990. Contributions of drug solubilization, partitioning, barrier disruption, and solvent permeation to the enhancement of skin permeation of various compounds with fatty acids and amines. *Pharm. Res.*, 7(7): 712-718.
- Avinash, H. H., Thorat, Y. S., Kasture, P. V. 2006. Carbopol and its pharmaceutical significance: a review. *Pharmaceutical Rev.*, 4(1).
- Azmin, M. N., Florence, A. T., Handjani-Vila, R. M., Stuart, J. F. B., Vanlerberghe, G., Whittaker, J. S. 1985. The effect of non-ionic surfactant vesicle (niosome) entrapment on the absorption and distribution of methotrexate in mice. *J. Pharm. Pharmacol.*, 37(4): 237-242.
- Baden, H., Kubilus, J. 1983. Effect of minoxidil on cultured keratinocytes. *J. Invest. Dermatol.*, 81: 558-560.
- Baillie, A. J., Florence, A. T., Hume, L. R., Muirhead, G. T., Rogerson, A. 1985. The preparation and properties of niosomes-non-ionic surfactant vesicles. *J. Pharm. Pharmacol.*, 37: 863-868.
- Bajaj, A., Desai, M. 2006. Challenges and strategies in novel drug delivery technologies. *Pharm. Times*, 38: 12-16.

- Balakrishnan, P., Shanmugam, S., Lee, W. S., Lee, W. M., Kim, J. O., Oh, D. H., Kim, D.-D., Kim, J. S., Yoo, B. K., Choi, H.-G., Woo, J. S., Yong, C. S. 2009. Formulation and *in vitro* assessment of minoxidil niosomes for enhanced skin delivery. *Int. J. Pharm.*, 377(1-2): 1-8.
- Barry, B. 2002. Drug delivery routes in skin: a novel approach. *Adv. Drug Del. Rev.*, 54: S31-S40.
- Barry, B. W. 1983. Dermatological formulations: Percutaneous absorption. Marcel Dekker, New York.
- Bayindir, Z. S., Yuksel, N. 2010. Characterization of niosomes prepared with various nonionic surfactants for paclitaxel oral delivery. *J. Pharm. Sci.*, 99(4): 2049-2060.
- Bent, S., Kane, C., Shinohara, K., Neuhaus, J., Hudes, E. S., Goldberg, H., Avins, A. L. 2006. Saw palmetto for benign prostatic hyperplasia. *N. Engl. J. Med.*, 354: 557-566.
- Bertolino, A. P., Klein, L. M., Freedberg, I. M. 1993. Biology of hair follicles. In: Fitzpatrick, T. B., Eisen, A. Z., Wolff, K., Freedberg, I. M., Austen, K. F. *Dermatology in general medicine*. McGraw-Hill, New York, pp. 289-293.
- Bhagavan, N. V. 2002. Lipid I: Fatty acids and eicodanoids. *Medical biochemistry* Harcourt Academic Press, Florida, pp. 1016.
- Biswal, S., Murthy, P., Sahu, J., Sahoo, P., Amir, F. 2008. Vesicles of non-ionic surfactants (niosomes) and drug delivery potential. *Int. J. Pharm. Sci. Nanotech.*, 1: 1-8.

- Blume-Peytavi, U., Massoudy, L., Patzelt, A., Lademann, J. r., Dietz, E., Rasulev, U., Garcia Bartels, N. 2010. Follicular and percutaneous penetration pathways of topically applied minoxidil foam. *Eur. J. Pharm. Biopharm.*, 76(3): 450-453.
- Bramson, H., Hermann, D., Batchelor, K., Lee, F., James, M., Frye, S. 1997. Unique preclinical characteristics of GG745, a potent dual inhibitor of 5-AR. *J. Pharmacol. Exp. Ther.*, 282: 1496-1502.
- Brimer, L., Lorentzen, B., Smitt, V. 1989. Evelser I farmakognosi K-25/9. Royal Danish School of Pharmacy, Copenhagen.
- Brunetti, L., Daghetta, A., Fedeli, E., Kikic, I., Zanderighi, L. 1989. Deacidification of olive oils by supercritical carbon dioxide. *J. Am. Oil Chem. Soc.*, 66: 209-217.
- Burdette, A., Garner, P. L., Mayer, E. P., Hargrove, J. L., Hartle, D. K., Greenspan, P. 2010. Anti-inflammatory activity of select sorghum (*Sorghum bicolor*) brans. *J. Med. Food*, 13: 879-887.
- Carafa, M., Santucci, E., Alhaique, F., Coviello, T., Murtas, E., Riccieri, F. M., Lucania, G., Torrisi, M. R. 1998. Preparation and properties of new unilamellar non-ionic/ionic surfactant vesicles. *Int. J. Pharm.*, 160(1): 51-59.
- Cevc, G., Blume, G., Schützlein, A., Gebauer, D., Paul, A. 1996. The skin: a pathway for systemic treatment with patches and lipid-based agent carriers. *Adv. Drug Del. Rev.*, 18(3): 349-378.
- Chakrabarty, M. M. 1989. Rice bran- a new source for edible and industrial oil. In: *The world conference on edible fats and oils processing- basic principles and modern practices*. Maastricht, the Netherlands.

- Chandraprakash, K., Udupa, N., Umadevi, P., Pillai, G. 1993. Effect of macrophage activation on plasma disposition of niosomal 3H-methotrexate in sarcoma-180 bearing mice. *J. Drug Targeting*, 1(2): 143-145.
- Chase, H. B. 1954. Growth of the hair. *Physiol. Rev.*, 34(1): 113-126.
- Chatterjee, A., Dutta, P. K., Chowdhury, R. 2007. Effect of fatty acids and cholesterol present in bile on expression of virulence factors and motility of *Vibrio cholerae*. *Infect. Immun.*, 75(4): 1946-1953.
- Chen, H., Chang, X., Du, D., Liu, W., Liu, J., Weng, T., Yang, Y., Xu, H., Yang, X. 2006. Podophyllotoxin-loaded solid lipid nanoparticles for epidermal targeting. *J. Controlled Release*, 110(2): 296-306.
- Cheung, P. C. K., Leung, A. Y. H., Ang, P. O. 1998. Comparison of supercritical carbon dioxide and soxhlet extraction of lipids from a brown seaweed, *Sargassum hemiphyllum* (Turn.) C. *J. Agr. Food Chem.*, 46(10): 4228-4232.
- Chotimarkorn, C., Benjakul, S., Silalai, N. 2008. Antioxidant components and properties of five long-grained rice bran extracts from commercial available cultivars in Thailand. *Food Chem.*, 111(3): 636-641.
- Chrastil, J. 1982. Solubility of solids and liquids in supercritical gases. *J. Phys. Chem.*, 86(15): 3016-3021.
- Chu, Y.-H., Hsu, H.-F. 1999. Effects of antioxidants on peanut oil stability. *Food Chem.*, 66(1): 29-34.
- Ciotti, S. N., Weiner, N. 2002. Follicular liposomal delivery. *J. Liposome Res.*, 12(1-2): 143-148.

- Clark, R. V., Hermann, D. J., Cunningham, G. R., Wilson, T. H., Morrill, B. B., Hobbs, S. 2004. Marked suppression of dihydrotestosterone in men with benign prostatic hyperplasia by dutasteride, a dual 5-alpha-reductase inhibitor. *J. Clin. Endocrinol. Metab.*, 89: 2179-2184.
- Colombo, V., Gerber, F., Bronhofer, M., Floersheim, G. 1990. Treatment of brittle fingernails and onychoschizia with biotin: scanning electron microscopy. *J. Am. Acad. Dermatol.*, 23: 1127-1132.
- Cosco, D., Paolino, D., Muzzalupo, R., Celia, C., Citraro, R., Caponio, D., Picci, N., Fresta, M. 2009. Novel PEG-coated niosomes based on bola-surfactant as drug carriers for 5-fluorouracil. *Bio. Mi.*, 11(5): 1115-1125.
- Cotsarelis, G., Millar, S. 2001. Towards a molecular understanding of hair loss and its treatment. *Trends Mol. Med.*, 7: 293-301.
- Courtois, M., Loussouarn, G., Hourseau, C., Grollier, J. F. 1994. Hair cycle and alopecia. *Skin Pharmacol.*, 7(1-2): 84-89.
- Cury-Boaventura, M. F., Pompeia, C., Curi, R. 2005. Comparative toxicity of oleic acid and linoleic acid on Raji cells. *Nutrition*, 21(3): 395-405.
- Dallob, A. L., Sadick, N. S., Unger, W., Lipert, S., Geissert, L. A., Gregoire, S. L., Nguyen, H. H., Moore, E. C., Tanaka, W. K. 1994. The effect of finasteride, a 5-alpha-reductase inhibitor, on scalp skin testosterone and dihydrotestosterone concentrations in patients with male pattern baldness. *J. Clin. Endocrinol. Metab.*, 79: 703-706.
- Das, P. K., Chaudhuri, A., Kaimal, T. N. B., Bhalerao, U. T. 1998. Isolation of gamma-Oryzanol through calcium ion induced precipitation of anionic micellar aggregates. *J. Agr. Food Chem.*, 46: 3073-3080.

- Das, U. N. 1991. Tumoricidal action of cis-unsaturated fatty acids and its relationship to free radicals and lipid peroxidation. *Cancer Lett.*, 56: 235 - 243.
- Datta, K., Singh, A. T., Mukherjee, A., Bhat, B., Ramesh, B., Burman, A. C. 2009. *Eclipta alba* extract with potential for hair growth promoting activity. *J. Ethnopharmacol.*, 124(3): 450-456.
- de Leeuw, J., de Vijlder, H., Bjerring, P., Neumann, H. 2009. Liposomes in dermatology today. *J. Eur. Acad. Dermatol. Venereol.*, 23: 505-516.
- Decker, E. A., Welch, B. 1990. Role of ferritin as a lipid oxidation catalyst in muscle food. *J. Agr. Food Chem.*, 38(3): 674-677.
- Délos, S., Iehlé, C., Martin, P.-M., Raynaud, J.-P. 1994. Inhibition of the activity of 5[alpha]-reductase (type 1) detected in DU 145 cells and expressed in insect cells. *J. Steroid Biochem. Mol. Biol.*, 48(4): 347-352.
- Deo, N., Somasundaran, P. 2003. Disintegration of Liposomes by Surfactants: mechanism of protein and cholesterol effects. *Langmuir*, 19(6): 2007-2012.
- Desai, A. R., Raghuveer, I., Chitme, H. R., Chandra, R. 2011. Formulation and evaluation of niosomal drug delivery of α -lipoic acid. *Drug Invent. Today*, 2(7): 325-327.
- Domashenko, A., Gupta, S., Cotsarelis, G. 2000. Efficient delivery of transgenes to human hair follicle progenitor cells using topical lipoplex. *Nat. Biotechnol.*, 18: 420-423.
- Draize, J. H., Woodard, G., Calvery, H. O. 1944. Methods for the study of previous termirritationnext term and toxicity of substances applied topically to the previous termskinnext term and mucous membranes. *J. Pharmacol. Exp. Ther.*, 82: 377-390.

Drake, L., Hordinsky, M., Fiedler, V., Swinehart, J., Unger, W. P., Cotterill, P. C., Thiboutot, D. M., Lowe, N., Jacobson, C., Whiting, D., Stieglitz, S., Kraus, S. J., Griffin, E. I., Weiss, D., Carrington, P., Gencheff, C., Cole, G. W., Pariser, D. M., Epstein, E. S., Tanaka, W., Dallob, A., Vandormael, K., Geissler, L., Waldsteicher, J. 1999. The effects of finasteride on scalp skin and serum androgen levels in men with androgenetic alopecia. *J. Am. Acad. Dermatol.*, 41(4): 550-554.

Dufes, C., Schätzlein, A. G., Tetley, L., Gray, A. I., Watson, D. G., Olivier, J.-C., Couet, W., Uchegbu, I. F. 2000. Niosomes and polymeric chitosan based vesicles bearing transferrin and glucose ligands for drug targeting. *Pharm. Res.*, 17(10): 1250-1258.

El Maghraby, G., Barry, B., Williams, A. 2008. Liposomes and skin: From drug delivery to model membranes. *Eur. J. Pharm. Sci.*, 34: 203-222

Ellis, J. A., Sinclair, R., Harrap, S. B. 2002. Androgenetic alopecia: pathogenesis and potential for therapy. *Expet Rev Mol Med*, 4: 1-11.

Erdog˘yan, S., Ozer, A. Y., Bilgili, H. 2005. *In vivo* behaviour of vesicular urokinase. *Int. J. Pharm.*, 295: 1-6.

Fang, J.-Y., Hong, C.-T., Chiu, W.-T., Wang, Y.-Y. 2001a. Effect of liposomes and niosomes on skin permeation of enoxacin. *Int. J. Pharm.*, 219(1-2): 61-72.

Fang, J.-Y., Yu, S.-Y., Wu, P.-C., Huang, Y.-B., Tsai, Y.-H. 2001b. *In vitro* skin permeation of estradiol from various proniosome formulations. *Int. J. Pharm.*, 215(1-2): 91-99.

Fardis, M., Oliyai, R., Stella, V. J., Borchardt, R. T., Hageman, M. J., Maag, H., Tilley, J. W. 2007. Case study: denofovir disoproxil fumarate: an oral prodrug of tenofovir prodrugs. Prodrugs, challenges and rewards Part 1 and Part 2 Springer New York, (pp. 1347-1357).

Feitosa, E., Karlsson, G. r., Edwards, K. 2006. Unilamellar vesicles obtained by simply mixing dioctadecyldimethylammonium chloride and bromide with water. *Chems Phys Lipids*, 140(1-2): 66-74.

Foldvari, M., Kumar, P., King, M., Batta, R., Michel, D., Badea, I., Wloch, M. 2006. Gene delivery into human skin *in vitro* using biphasic lipid vesicles. *Curr. Drug Del.*, 3: 89-93.

Friedrich, J. P., List, G. R. 2002. Characterization of soybean oil extracted by supercritical carbon dioxide and hexane. *J. Agr. Food Chem.*, 30(1): 192-193.

Frum, Y., Bonner, M. C., Eccleston, G. M., Meidan, V. M. 2007. The influence of drug partition coefficient on follicular penetration: *In vitro* human skin studies. *Eur. J. Pharm. Sci.*, 30: 280-287.

Fustier, P., Taherian, A. R., Ramaswamy, H. S. 2010. Emulsion delivery systems for functional foods. In: Smith, J., Charter, E. Functional food product development. Blackwell Publishing, Singapore.

García, A., de Lucas, A., Rincón, J., Alvarez, A., Gracia, I., García, M. 1996. Supercritical carbon dioxide extraction of fatty and waxy material from rice bran. *J. Am. Oil Chem. Soc.*, 73(9): 1127-1131.

Giada, M. D. L. R., Mancini-Filho, J. 2009. Antioxidant capacity of the striped sunflower (*Helianthus annuus* L.) seed extracts evaluated by three *in vitro* methods. *Int. J. Food Sci. Nutr.*, 60(5): 395-401.

- Gibson, N., Shenderova, O., Luo, T. J. M., Moseenkov, S., Bondar, V., Puzyr, A., Purto, K., Fitzgerald, Z., Brenner, D. W. 2009. Colloidal stability of modified nanodiamond particles. *Diam. Relat. Mater.*, 18(4): 620-626.
- Glavas-Dodov, M., Fredro-Kumbaradzi, E., Goracinova, K., Calis, S., Simonoska, M., Hincal, A. A. 2003. 5-Fluorouracil in topical liposome gels for anticancer treatment-formulation and evaluation. *Acta Pharmaceut.*, 53(4): 241-250.
- Golden, G. M., Guzek, D. B., Harris, R. R., McKie, J. E., Potts, R. O. 1986. Lipid thermotropic transitions in human stratum corneum. *J. Invest. Dermatol.*, 86: 255-259.
- Goodrum, J. W., Kilgo, M. B. 1987. Peanut oil extraction with sc-co₂ : solubility and kinetic functions *Trans. Am. Soc. Agric. Eng.*, 30(6): 1865-1868.
- Goyan, J. E., Arbor, A., Wruble, M. 1961. Polyethylene glycol suppository bases. In: U. S. P. Office. United States.
- Grams, Y. Y. 2005. Influence of molecular properties and delivery system design on the transfollicular transport across the skin. Ph. D. thesis, Division of Drug Delivery Technology of the Leiden/Amsterdam Center for Drug Research, Leiden University, Leiden.
- Grams, Y. Y., Alarukka, S., Lashley, L., Caussin, J., Whitehead, L., Bouwstra, J. A. 2003. Permeant lipophilicity and vehicle composition influence accumulation of dyes in hair follicles of human skin. *Eur. J. Pharm. Sci.*, 18(5): 329-336.
- Grams, Y. Y., Whitehead, L., Cornwell, P., Bouwstra, J. A. 2004. Time and depth resolved visualisation of the diffusion of a lipophilic dye into the hair follicle of fresh unfixed human scalp skin. *J. Controlled Release*, 98(3): 367-378.

- Greenrod, W., Fenech, M. 2003. The principal phenolic and alcoholic components of wine protect human lymphocytes against hydrogen peroxide- and ionizing radiation-induced DNA damage *in vitro*. *Mutagenesis*, 18(2): 119-126.
- Gregoriadis, G., McCormack, B., Obrenovich, M., Perrie, Y., Lowrie, D. B., Whalen, R. G. 2000. Entrapment of Plasmid DNA Vaccines into Liposomes by Dehydration/Rehydration DNA Vaccines. vol. 29 Humana Press, (pp. 305-311).
- Grice, J. E., Ciotti, S., Weiner, N., Lockwood, P., Cross, S. E., Roberts, M. S. 2010. Relative uptake of minoxidil into appendages and stratum corneum and permeation through human skin *in vitro*. *J. Pharm. Sci.*, 99(2): 712-718.
- Guinedi, A. S., Mortada, N. D., Mansour, S., Hathout, R. M. 2005. Preparation and evaluation of reverse-phase evaporation and multilamellar niosomes as ophthalmic carriers of acetazolamide. *Int. J. Pharm.*, 306(1-2): 71-82.
- Gulcin, I. 2006. Antioxidant and antiradical activities of L-carnitine. *Life Sci.*, 78: 803-811.
- Gunstone, F. D. 2011. Vegetable oils in food technology composition, properties and uses. Wiley-blackwell, West sussex, UK.
- Habib, F., Ross, M., Ho, C., Lyons, V., Chapman, K. 2005. *Serenoa repens* (Permixon) inhibits the 5-alpha-reductase activity of human prostate cancer cell lines without interfering with PSA expression. *Int. J. Cancer*, 114: 190-194.
- Hadgraft, J. 2001. Skin, the final frontier. *Int. J. Pharm.*, 224: 1-18
- Hagerman, A. E., Butler, L. G. 1989. Choosing appropriate methods and standards for assaying tannin. *J. Chem. Ecol.*, 15: 1795-1810.

- Hamilton, J. B. 1951. Patterned loss of hair in man; types and incidence. *Annals NY Acad. Sci.*, 53(3): 708-728.
- Han, J., Kwon, O., Chung, J., Cho, K., Eun, H., Kim, K. 2004. Effect of minoxidil on proliferation and apoptosis in dermal papilla cells of human hair follicle. *J. Dermatol. Sci.*, 34: 91-98.
- Handratta, V. D., Vasaitis, T. S., Njar, V. C., Gediya, L. K., Kataria, R., Chopra, P., Newman, D., Jr., Farquhar, R., Guo, Z., Qiu, Y., Brodie, A. M. 2005. Novel C-17-heteroaryl steroidal CYP17 inhibitors/antiandrogens: synthesis, *in vitro* biological activity, pharmacokinetics, and antitumor activity in the LAPC4 human prostate cancer xenograft model. *J. Med. Chem.*, 48(8): 2972-2984.
- Haran, G., Cohen, R., Bar, L. K., Barenholz, Y. 1993. Transmembrane ammonium sulfate gradients in liposomes produce efficient and stable entrapment of amphipathic weak bases. *BBA - Biomembranes*, 1151(2): 201-215.
- Harbone, J. R. 1976. *Phytochemical methods. a guide to modern techniques of plant analysis*. Charpan & Hall, London.
- Harborne, J. B. 2005. *Phytochemical methods. analysis. a guide to modern techniques of plant*. Springer International, India.
- Hashim, F., El-Ridy, M., Nasr, M., Abdallah, Y. 2010. Preparation and characterization of niosomes containing ribavirin for liver targeting. *Drug Deliv.*, 17(5): 282-287.
- Hashimoto, K., Ito, M., Suzuki, Y. 1990. Innervation and vasculature of the hair follicle. In: C. E. O. a. R. Happle, *Hair and hair diseases*. Springer-Verlag, New York.

Hashimoto, K., Shibazaki, S. 1976. Ultrastructural study on differentiation and function of hair. In: Kobori, T., Montagna, W., Toda, K., Ishibashi, Y., Hori, Y. and Morikawa, F. Biology and disease of the hair. University of Tokyo Press, Tokyo.

Hattori, M., Ogawa, H. 1983. Biochemical analysis of hair growth from the aspects of aging and enzyme activities. *J. Dermatol.*, 10: 45-54.

Hegde, M. N., C.Rodrigues, J., Kumari, S., Hegde, N. D. 2006. Toxicity evaluation of root canal sealers on human gingival fibroblasts. *Endod.*, 18(1): 42-48.

Hibberts, N. A., Howell, A. E., Randall, V. A. 1998. Balding hair follicle dermal papilla cells contain higher levels of androgen receptors than those from non-balding scalp. *J. Endocrinol.*, 156(1): 59-65.

Hirokawa, T., Hirokawa, S., Yamauchi, N., Kataoka, T., Woo, J.-T., Nagai, K. 1997. Immunomodulating activities of polysaccharide fractions from dried safflower petals. *Cytotech.*, 25 205-211.

Hoffman, R. M. 1998. Topical liposome targeting of dyes, melanins, genes, and proteins selectively to hair follicles. *J. Drug Targeting*, 5: 67-74.

Hofland, H. E. J., Van der Geest, R., Bodde, H. E., Junginger, H. E., Bouwstra, J. A. 1994. Estradiol permeation from nonionic surfactant vesicles through human stratum corneum *in vitro*. *Pharm. Res.*, 11(5): 659-664.

Hu, C., Rhodes, D. G. 1999. Proniosomes: A novel drug carrier preparation. *Int. J. Pharm.*, 185(1): 23-35.

Hueber, F., Wepierre, J., Schaefer, H. 1992. Role of Transepidermal and transfollicular routes in percutaneous absorption of hydrocortisone and testosterone: *In vivo* study in the hairless rat. *Skin Pharmacol. Physiol.*, 5(2): 99-107.

Imura, T., Gotoh, T., Otake, K., Yoda, S., Takebayashi, Y., Yokoyama, S., Takebayashi, H., Sakai, H., Yuasa, M., Abe, M. 2003. Control of physicochemical properties of liposomes using a supercritical reverse phase evaporation method. *Langmuir*, 19(6): 2021-2025.

Iqbal, S., Bhangar, M. I., Anwar, F. 2005. Antioxidant properties and components of some commercially available varieties of rice bran in Pakistan. *Food Chem.*, 93(2): 265-272.

Irving, R. 2007. Baldness cure firm heads for AIM. *The Times*. Times Newspapers Ltd., England.

Izzo, A. A., Ernst, E. 1992. Interaction between herbal medicines and prescribed drugs: a systematic review. *Drugs*, 6: 2163-2175.

Jadon, P., Gajbhiye, V., Jadon, R., Gajbhiye, K., Ganesh, N. 2009. Enhanced oral bioavailability of griseofulvin via niosomes. *AAPS Pharm. Sci.*, 10(4): 1186-1192.

Jain, C. P., Vyas, S. P. 1995. Preparation and characterization of niosomes containing rifampicin for lung targeting. *J. Microencaps.*, 12(4): 401-407.

Jain, C. P., Vyas, S. P., Dixit, V. K. 2006. Niosomal system for delivery of rifampicin to lymphatics. *Indian J. Pharm. Sci.*, 68: 575-578.

- Jain, S., Singh, P., Mishra, V., Vyas, S. P. 2005. Mannosylated niosomes as adjuvant carrier system for oral genetic immunization against Hepatitis B. *Immunol Lett*, 101(1): 41-49.
- Jain, S., Umamaheshwari, R. B., Bhadra, D., Jain, N. K. 2004. Ethosomes: A novel vesicular carrier for enhanced transdermal delivery of an anti HIV agent. *Indian J. Pharm. Sci.*, 66: 72-81.
- Jenkins, E. P., Andersson, S., Imperato-McGinley, J., Wilson, J. D., Russell, D. W. 1992. Genetic and pharmacological evidence for more than one human steroid 5 alpha-reductase. *J. Clin. Invest.*, 89(1): 293-300.
- Jennings, B. H., Akoh, C. C. 2009. Characterization of a rice bran oil structured lipid. *J Agric Food Chem*, 57(8): 3346-3350.
- Jin, Y., Xiao, Y.-s., Zhang, F.-f., Xue, X.-y., Xu, Q., Liang, X.-m. 2008. Systematic screening and characterization of flavonoid glycosides in *Carthamus tinctorius* L. by liquid chromatography/UV diode-array detection/electrospray ionization tandem mass spectrometry. *J. Pharmaceut. Biomed.*, 46(3): 418-430.
- Jung, S., Otberg, N., Thiede, G., Richter, H., Sterry, W., Panzner, S., Lademann, J. 2006. Innovative Liposomes as a transfollicular drug delivery system: penetration into porcine hair follicles. *J. Invest. Dermatol.*, 126(8): 1728-1732.
- Junyaprasert, V., Teeranachaideekul, V., Supaperm, T. 2008. Effect of charged and non-ionic membrane additives on physicochemical properties and stability of niosomes. *AAPS Pharm. Sci.*, 9(3): 851-859.
- Kalyanasundaram, K. 1987. Photochemistry in microheterogeneous systems. Academic Press, Inc., Orlando, FL.

- Kandimalla, K., Kanikkannan, N., Andega, S., Singh, M. 1999. Effect of fatty acids on the permeation of melatonin across rat and pig skin *in vitro* and on the transepidermal water loss in rats *in-vivo*. *J. Pharm. Pharmacol.*, 51(7): 783-790.
- Kang, B. M., Shin, S. H., Kwack, M. H., Shin, H., Oh, J. W., Kim, J., Moon, C., Moon, C., Kim, J. C., Kim, M. K., Sung, Y. K. 2010. Erythropoietin promotes hair shaft growth in cultured human hair follicles and modulates hair growth in mice. *J. Dermatol. Sci.*, 59(2): 86-90.
- Kato, K., Walde, P., Mitsui, H., Higashi, N. 2003. Enzymatic activity and stability of d-fructose dehydrogenase and sarcosine dehydrogenase immobilized onto giant vesicles. *Biotechnol. Bioeng.*, 84(4): 415-423.
- Kazuma, K., Takahashi, T., Sato, K., Takeuchi, H., Matsumoto, T. T. O., Okumo, T. 2000. Quinochalcones and flavonoids from fresh florets in different cultivars of *Carthamus tinctorius* L. *Biosci. Biotech. Biochem.*, 64: 1588-1599.
- Kertesz, Z., Szikszai, Z., Pelicon, P., Simcic, J., Telek, A., Biro, T. 2007. Ion beam microanalysis of human hair follicles. *Nucl. Instrum. Meth. B.*, 260(1): 218-221.
- Khandare, J. N., Madhavi, G., Tamhankar, B. M. 1994. Niosomes novel drug delivery system. *East. Pharm.*, 37: 61-64.
- Khazaeli, P., Pardakhty, A., Shoorabi, H. 2007. Caffeine-loaded niosomes: characterization and *in vitro* release studies. *Drug Deliv.*, 14(7): 447-452.
- Kitson, N., van-Lennep, E. W. 1984. Intercellular junctions and the permeability barrier in human sebaceous glands. *Br. J. Dermatol.*, 110: 601-606.
- Kligman, A. 1959. The human hair cycle. *J. Invest. Dermatol.*, 33: 307-316.

- Knorr, F., Lademann, J. r., Patzelt, A., Sterry, W., Blume-Peytavi, U., Vogt, A. 2009. Follicular transport route - Research progress and future perspectives. *Eur. J. Pharm. Biopharm.*, 71(2): 173-180.
- Krause, K., Foitzik, K. 2006. Biology of the hair follicle: The basics. *Semin. Cutan. Med. Surg.*, 25(1): 2-10.
- Kuk, M. S., Dowd, M. K. 1998. Supercritical CO₂ extraction of rice bran. *J. Am. Oil Chem. Soc.*, 75(5): 623-628.
- Kumar, G. P., Rajeshwarrao, P. 2011. Nonionic surfactant vesicular systems for effective drug delivery-an overview. *Acta Pharmaceut. B*, 1(4): 208-219.
- Kumarn, G. P., Rajeshwarrao, P. 2011. Nonionic surfactant vesicular systems for effective drug delivery-an overview. *Acta Pharmaceut. Sinica B*, 1(4): 208-219.
- Kumhar, S. K., Jain, S. K., Pancholi, S. S., Agrawal, S., Saraf, D. K., Agrawal, G. P. 2003. Provesicular transdermal drug delivery system of ethinylestradiol and levonorgestrel for contraception and hormone replacement therapy. *Indian J. Pharm. Sci.*, 65(6): 620-627.
- Kwon, O. S., Pyo, H. K., Oh, Y. J., Han, J. H., Lee, S. R., Chung, J. H., Eun, H. C., Kim, K. H. 2007. Promotive effect of minoxidil combined with all-trans retinoic acid (tretinoin) on human hair growth *in vitro*. *J Korean Med Sci*, 22(2): 283-289.
- Lachgar, S., Charveron, M., Gall, Y., Bonafe, J. 1998. Minoxidil upregulates the expression of vascular endothelial growth factor in human hair dermal papilla cells. *Br. J. Dermatol.*, 138: 407-411.

Lademann, J., Knorr, F., Richter, H., Blume-Peytavi, U., Vogt, A., Antoniou, C., Sterry, W., Patzelt, A. 2008. Hair follicles - an efficient storage and penetration pathway for topically applied substances. *Skin Pharmacol. Physiol.*, 21(3): 150-155.

Lademann, J., Richt, H., Teichmann, A., Otberg, N., Blume-Peytavi, U., Luengo, J., Weiß, B. F., Schaefer, U., Lehr, C., Wepf, R., Sterry, W. 2007. Nanoparticles - An efficient carrier for drug delivery into the hair follicles. *Eur. J. Pharm. Biopharm.*, 66: 159-164.

Lademann, J., Richter, H., Schaefer, U., Blume-Peytavi, U., Teichmann, A., Otberg, N., Sterry, W. 2006. Hair follicles - A long-term reservoir for drug delivery. *Skin Pharmacol. Physiol.*, 19: 232-236.

Lakshmi, P. K., Bhaskaran, S. 2011. Phase II study of topical niosomal urea gel-an adjuvant in the treatment of psoriasis. *International Journal of Pharmaceutical Sciences Review and Research*, 7: 1-7.

Lakshmi, P. K., Devi Gayathri, S., Shyamala, B., Sacchidanand, S. 2007. Niosomal methotrexate gel in the treatment of localized psoriasis: Phase I and phase II studies. *Indian J. Dermatol. Venereol. Leprol.*, 73(3): 157-161.

Lala, S., Nandy, A. K., Mahato, S. B., Basu, M. K. 2003. Delivery *in vivo* of 14-deoxy-11-oxoandrographolide, an antileishmanial agent, by different drug carriers. *Indian J Biochem Biophys*, 40(3): 169-174.

Lautenschläger, H. 2003. Essential fatty acids - cosmetic from inside and outside. *Beauty Forum*, 4: 54-56.

- Leaf, A., Kang, J. X., Xiao, Y.-F., Billman, G. E. 2003. Clinical prevention of sudden cardiac death by n-3 polyunsaturated fatty acids and mechanism of prevention of arrhythmias by n-3 fish oils. *Circulation*, 107(21): 2646-2652.
- Lee, G.-S., Hong, E.-J., Gwak, K.-S., Park, M.-J., Choi, K.-C., Choi, I.-G., Jang, J.-W., Jeung, E.-B. 2010. The essential oils of *Chamaecyparis obtusa* promote hair growth through the induction of vascular endothelial growth factor gene. *Fitoterapia*, 81(1): 17-24.
- Lee, Z., Hong, G., Cho, M. 2008. Antioxidant, antibacterial and tyrosinase inhibition activities of compounds from rice bran and fermented rice bran. *J. Biotech.*, 136: S442-S442.
- Lekki, J., Stachura, Z., Dabros, W., Stachura, J., Menzel, F., Reinert, T., Butz, T., Pallon, J., Gontier, E., Ynsa, M. D., Moretto, P., Kertesz, Z., Szikszai, Z., Kiss, A. Z. 2007. On the follicular pathway of percutaneous uptake of nanoparticles: Ion microscopy and autoradiography studies. *Nucl. Instrum. Meth. B.*, 260(1): 174-177.
- Li, Y., Mitra, A. K. 1996. Effects of phospholipid chain length, concentration, charge, and vesicle size on pulmonary insulin absorption. *Pharm. Res.*, 13(1): 76-79.
- Liang, T., Liao, S. 1992. Inhibition of steroid 5 alpha-reductase by specific aliphatic unsaturated fatty acids. *Biochem. J.*, 285(2): 557-562.
- Limat, A., Suormala, T., Hunziker, T., Waelti, E., Braathen, L., Baumgartner, R. 1996. Proliferation and differentiation of cultured human follicular keratinocytes are not influenced by biotin. *Arch. Dermatol. Res.*, 288: 31-38.
- Liyana-Pathirana, C. M., Shahidi, F. 2007. Antioxidant and free radical scavenging activities of whole wheat and milling fractions. *Food Chem.*, 101: 1151-1157.

- Löntz, W., Sirsjö, A., Liu, W., Lindberg, M., Rollman, O., Törmä, H. 1995. Increased mRNA expression of manganese superoxide dismutase in psoriasis skin lesions and in cultured human keratinocytes exposed to IL-1[beta] and TNF-[alpha]. *Free Radical Bio. Med.*, 18(2): 349-355.
- Lou, X., Janssen, H. G., Cramers, C. A. 1996. Effects of modifier addition and temperature variation in SFE of polymeric materials. *J. Chromatogr. Sci.*, 34: 282-290.
- Lu, X., Yu, H., Ma, Q., Shen, S., Das, U. 2010. Linoleic acid suppresses colorectal cancer cell growth by inducing oxidant stress and mitochondrial dysfunction. *Lipids Health and Dis.*, 9(1): 106.
- Luciani, A., Olivier, J.-C., Clement, O., Siauve, N., Brillet, P.-Y., Bessoud, B., Gazeau, F., Uchegbu, I. F., Kahn, E., Frija, G., Cuenod, C. A. 2004. Glucose-receptor MR imaging of tumors: study in mice with PEGylated paramagnetic niosomes. *Radiology*, 231(1): 135-142.
- Lucky, A., Picquadio, D., Ditre, C., Dunlap, F., Kantor, I., Pandya, A., AG, P., Savin, R., Tharp, M. 2004. A randomized, placebo-controlled trial of 5% and 2% topical minoxidil solutions in the treatment of female pattern hair loss. *J. Am. Acad. Dermatol.*, 50: 541-553.
- Luque de Castro, M. D., Garcia-Ayuso, L. E. 1998. Soxhlet extraction of solid materials: an outdated technique with a promising innovative future. *Anal. Chim. Acta*, 369: 1-10.
- MacDonald, H. B. 2000. Conjugated linoleic acid and disease prevention: a review of current knowledge. *J. Am. Coll. Nutri.*, 19: 111s-118s.

- Madhusudhan, B., Rambhau, D., Apte, S. S., Gopinath, D. 2007. Oral bioavailability of flutamide from alkylglycerol stabilized o/w nanoemulsions. *J. Disper. Sci. Technol.*, 28(8): 1254-1261.
- Maisuthisakul, P., Gordon, M. H. 2009. Antioxidant and tyrosinase inhibitory activity of mango seed kernel by product. *Food Chem.*, 117(2): 332-341.
- Malenčić, D., Popović, M., Miladinović, J. 2007. Phenolic Content and antioxidant properties of soybean (*Glycine max* (L.) Merr.) seeds. *Molecules*, 12: 576-581.
- Manconi, M., Valenti, D., Sinico, C., Lai, F., Loy, G., Fadda, A. M. 2003. Niosomes as carriers for tretinoin: II. Influence of vesicular incorporation on tretinoin photostability. *Int. J. Pharm.*, 260(2): 261-272.
- Manosroi, A., Chutoprapat, R., Abe, M., Manosroi, J. 2008a. Characteristics of niosomes prepared by supercritical carbon dioxide (scCO₂) fluid. *Int. J. Pharm.*, 352(1-2): 248-255.
- Manosroi, A., Jantrawut, P., Manosroi, J. 2008b. Anti-inflammatory activity of gel containing novel elastic niosomes entrapped with diclofenac diethylammonium. *Int. J. Pharm.*, 360: 156-163.
- Manosroi, A., Jantrawut, P., Khositsuntiwong, N., Manosroi, W., Manosroi, J. 2009. Novel elastic nanovesicles for cosmeceutical and pharmaceutical applications. *Chiang Mai J. Sci.*, 36(2): 168-178.
- Manosroi, A., Jantrawut, P., Akihisa, T., Manosroi, W., Manosroi, J. 2011. *In vitro* and *in vivo* skin anti-aging evaluation of gel containing niosomes loaded with a semi-purified fraction containing gallic acid from *Terminalia chebula* galls. *Pharm. Biol.*, 49(11): 1190-1203.

Manosroi, A., Ruksiriwanich, W., Abe, M., Sakai, H., Manosroi, W., Manosroi, J. 2010. Biological activities of the rice bran extract and physical characteristics of its entrapment in niosomes by supercritical carbon dioxide fluid. *J. Supercrit. Fluids*, 54(2): 137-144.

Manosroi, A., Ruksiriwanich, W., Abe, M., Manosroi, W., Manosroi, J. 2012a. Transfollicular enhancement of gel containing cationic niosomes loaded with unsaturated fatty acids in rice (*Oryza sativa*) bran semi-purified fraction. *Eur. J. Pharm. Biopharm.*, Accepted.

Manosroi, A., Ruksiriwanich, W., Manosroi, W., Abe, M., Manosroi, J. 2012b. *In vivo* hair growth promotion activity of gel containing niosomes loaded with the *Oryza sativa* bran fraction (OSF3). *Advanced Science Letters*, Accepted.

Manosroi, A., Wongtrakul, P., Manosroi, J., Midorikawa, U., Hanyu, Y., Yuasa, M., Sugawara, F., Sakai, H., Abe, M. 2005. The entrapment of kojic oleate in bilayer vesicles. *Int. J. Pharm.*, 298(1): 13-25.

Manosroi, A., Wongtrakul, P., Manosroi, J., Sakai, H., Sugawara, F., Yuasa, M., Abe, M. 2003. Characterization of vesicles prepared with various non-ionic surfactants mixed with cholesterol. *Colloid. Surface. B*, 30: 129-138.

Maria, H. 2008. Advances in hair diseases. *Adv. Dermatol.*, 24(0): 245-259.

Marrone, C., Poletto, M., Reverchon, E., Stassi, A. 1998. Almond oil extraction by supercritical CO₂: experiments and modelling. *Chem. Eng. Sci.*, 53(21): 3711-3718.

- Matsuda, H., Yamazaki, M., Naruto, S., Asanuma, Y., Kubo, M. 2002. Anti-androgenic and hair growth promoting activities of *Lygodii Spora* (spore of *Lygodium japonicum*) I. active constituents inhibiting testosterone 5 α -reductase. *Biol. Pharm. Bull.*, 25: 622-626.
- Maurer, N., Wong, K. F., Stark, H., Louie, L., McIntosh, D., Wong, T., Scherrer, P., Semple, S. C., Cullis, P. R. 2001. Spontaneous entrapment of polynucleotides upon electrostatic interaction with ethanol-destabilized cationic liposomes. *Biophys J*, 80(5): 2310-2326.
- Meidan, V. M., Bonner, M. C., Michniak, B. B. 2005. Transfollicular drug delivery - Is it a reality? *Int. J. Pharm.*, 306(1-2): 1-14.
- Meisheri, K., Cipkus, L., Taylor, C. 1988. Mechanism of action of minoxidil sulfate-induced vasodilation: a role for increased K⁺ permeability. *J. Pharmacol. Exp. Ther.*, 245: 751-760.
- Meng, M., Liu, Y., Wang, Y.-B., Wang, J.-C., Zhang, H., Wang, X.-Q., Zhang, X., Lu, W.-L., Zhang, Q. 2008. Increase of the pharmacological and pharmacokinetic efficacy of negatively charged polypeptide recombinant hirudin in rats via parenteral route by association with cationic liposomes. *J. Controlled Release*, 128(2): 113-119.
- Merrill, L. I., Pike, O. A., Ogden, L. V., Dunn, M. L. 2008. Oxidative stability of conventional and high-oleic vegetable oils with added antioxidants. *J. Am. Oil Chem. Soc.*, 85: 771-776.
- Mezei, M. 1994. Liposomes as penetration promoters and localizers of topically applied drugs. *J. Pharm. Sci.*, 62: 190-195.

- Michael, J. 2004. Quinoline, quinazoline and acridone alkaloids. *Nat. Prod. Rep.*, 21: 650-668.
- Miyachi, Y. 1995. Photoaging from an oxidative standpoint. *J. Dermatol. Sci.*, 9(2): 79-86.
- Most, M. M., Tulley, R., Morales, S., Lefevre, M. 2005. Rice bran oil, not fiber, lowers cholesterol in humans. *American Journal of Clinical Nutrition*, 81(1): 64-68.
- Mukherjee, B., Patra, B., Layek, B., Mukherjee, A. 2007. Sustained release of acyclovir from nano-liposomes and nano-niosomes: An *in vitro* study. *Int. J. Nanomed.*, 2: 213-225.
- Muller, J. M., Lelievre, V., Becq-Giraudon, L., Meunier, A. C. 1995. VIP as a cell-growth and differentiation neuromodulator role in neurodevelopment. *Mol. Neurobiol.*, 10(2-3): 115-134.
- Muller-Rover, S., Handjiski, B., van der Veen, C., Eichmuller, S., Foitzik, K., McKay, I. A., Stenn, K. S., Paus, R. 2001. A comprehensive guide for the accurate classification of murine hair follicles in distinct hair cycle stages. *J. Invest. Dermatol.*, 117(1): 3-15.
- Munster, U., Nakamura, C., Haberland, A., Jores, K., Mehnert, W., Rummel, S., Schaller, M., Korting, H. C., Zouboulis, C. C., Blume-Peytavi, U., Schafer-Korting, M. 2005. RU 58841-myristate - Prodrug development for topical treatment of acne and androgenetic alopecia. *Pharmazie*, 60(1): 8-12.
- Mura, S., Pirot, F., Manconi, M., Falson, F., Fadda, A. M. 2007. Liposomes and niosomes as potential carriers for dermal delivery of minoxidil. *J. Drug Targeting*, 15: 101-108.

- Nagatsu, A., Zhang, H., Mizukami, H., Okuyama, H., Sakakibara, J., Tokuda, H., Nishino, H. 2000. Tyrosinase inhibitory and anti-tumor promoting activities of compounds isolated from safflower (*Carthamus tinctorius* L.) and cotton (*Gossypium hirsutum* L.) oil cakes. *Nat. Prod. Lett.*, 14(3): 153-158.
- Nasr, M. 2010. *In vitro* and *in vivo* evaluation of proniosomes containing celecoxib for oral administration. *AAPS Pharm Sci.*, 11(1): 85-89.
- Naudé, Y., de Beer, W. H. J., Jooste, S., van der Merwe, L., van Rensburg, S. J. 1998. Comparison of supercritical fluid extraction and Soxhlet extraction for the determination of DDT, DDD and DDE in sediment. *Water SA*, 24(3): 205-214.
- Naziroglu, M., Kokcam, I. 2000. Antioxidants and lipid peroxidation status in the blood of patients with alopecia. *Cell Biochem. Funct.*, 18: 169-173.
- Nicoletti, G., Boghossian, V., Gurevitch, F., Borland, R., Morgenroth, P. 1993. The antimicrobial activity *in vitro* of chlorhexidine, a mixture of isothiazolinones (~Kathon) and cetyl trimethyl ammonium bromide (CTAB). *J. Hosp. Infect.*, 23(2): 87-111.
- Ning, M., Guo, Y., Pan, H., Chen, X., Gu, Z. 2005. Preparation, *in vitro* and *in vivo* evaluation of liposomal/niosomal gel delivery systems for clotrimazole. *Drug Dev. Ind. Pharm.*, 31(4-5): 375-383.
- Norman, W. 1998. Targeted follicular delivery of macromolecules via liposomes. *Int. J. Pharm.*, 162: 29-38.
- Nutan, M. T. H., Reddy, I. K. 2009. General principles of suspension. In: A. K. Kulshreshtha, O. N. Singh, & G. M. Wall, *Pharmaceutical suspensions*. Springer, London.

OECD 2002. OECD Guidelines for testing of chemicals. Acute dermal irritation corrosion Method 404. OECD publishing.

Oliver, R., Jahoda, C. 1989. The dermal papilla and the maintenance of hair growth. Chapman and Hall, London.

Olsen, E. A., Hordinsky, M., Whiting, D., Stough, D., Hobbs, S., Ellis, M. L., Wilson, T., Rittmaster, R. S. 2006. The importance of dual 5-alpha reductase inhibition in the treatment of male pattern hair loss: results of a randomized placebo-controlled study of dutasteride vs. finasteride. *J. Am. Acad. Dermatol.*, 55: 1014-1023.

Olsen, E., Weiner, M., Amara, I., DeLong, E. 1990. Five-year follow up of men with androgenetic alopecia treated with topical minoxidil. *J. Am. Acad. Dermatol.*, 22: 643-646.

Onwukaeme, D. N., Ikuegbvweha, T. B., Asonye, C. C. 2007. Evaluation of phytochemical constituents, antibacterial activities and effect of exudates of *Pycanthus angolensis* Weld Warb (Myristicaceae) on corneal ulcers in rabbits. *Trop. J. Pharm. Res.*, 6: 725-730.

Osawa, T., Namiki, M. 1981. A novel type of antioxidant isolated from leaf wax of *Eucalyptus* leaves. *Agr. Biol. Chem.*, 45(3): 735-739.

Ossadnik, M., Czaika, V., Teichmann, A., Sterry, W., Tietz, H. J., Lademann, J., Koch, S. 2007. Differential stripping: introduction of a method to show the penetration of topically applied antifungal substances into the hair follicles. *Mycoses*, 50(6): 457-462.

- Ossadnik, M., Richter, H., Teichmann, A., Koch, S., Schaefer, U., Wepf, R., Sterry, W., Lademann, J. 2006. Investigation of differences in follicular penetration of particle-and nonparticle-containing emulsions by laser scanning microscopy. *Laser Phys*, 16(5): 747-750.
- Otake, K., Imura, T., Sakai, H., Abe, M. 2001. Development of a new preparation method of liposomes using supercritical carbon dioxide. *Langmuir*, 17(13): 3898-3901.
- Otberg, N., Patzelt, A., Rasulev, U., Hagemester, T., Linscheid, M., Sinkgraven, R., Sterry, W., J. L. 2008. The role of hair follicles in the percutaneous absorption of caffeine. *Br. J. Clin. Pharmacol.*, 65: 488-492.
- Otberg, N., Richter, H., Schaefer, H., Blume-Peytavi, U., Sterry, W., Lademann, J. 2004. Variations of hair follicle size and distribution in different body sites. *J. Invest. Dermatol.*, 122: 14-19.
- Otberg, N., Teichmann, A., Rasuljev, U., Sinkgraven, R., Sterry, W., Lademann, J. 2007. Follicular penetration of topically applied caffeine via a shampoo formulation. *Skin Pharmacol. Physiol.*, 20(4): 195-198.
- Pérez-Ornelas, V., Cabeza, M., Bratoeff, E., Heuze, I., Sánchez, M., Ramirez, E., Naranjo-Rodriguez, E. 2005. New 5 alpha-reductase inhibitors: *In vitro* and *in vivo* effects. *Steroids Lipids Res.*, 70: 217-224.
- Padula, C., Fulgoni, A., Santi, P. 2010. *In vivo* stratum corneum distribution of lidocaine, assessed by tape stripping, from a new bioadhesive film. *Skin Res. Technol.*, 16: 125-130.

- Paladini, R. D., Saleh, J., Qian, C., Xu, G.-X., Rubin, L. L. 2005. Modulation of hair growth with small molecule agonists of the hedgehog signaling pathway. *J. Invest. Dermatol.*, 125(4): 638-646.
- Palter, R., Lundin, R. E., Haddon, W. F. 1972. A cathartic lignan glycoside isolated from *Carthamus tinctorius*. *Phytochem.*, 11: 2871-2874.
- Paolino, D., Muzzalupo, R., Ricciardi, A., Celia, C., Picci, N., Fresta, M. 2007. *In vitro* and *in vivo* evaluation of Bola-surfactant containing niosomes for transdermal delivery. *Bio. Mi.*, 9(4): 421-433.
- Papazisis, K. T., Geromichalos, G. D., Dimitriadis, K. A., Kortsaris, A. H. 1997. Optimization of the sulforhodamine B colorimetric assay. *J. Immunol. Methods*, 208(2): 151-158.
- Pardakhty, A., Varshosaz, J., Rouholamini, A. 2007. *In vitro* study of polyoxyethylene alkyl ether niosomes for delivery of insulin. *Int. J. Pharm.*, 328(2): 130-141.
- Park, S., Shin, W.-S., Ho, J. 2011. *Fructus panax ginseng* extract promotes hair regeneration in C57BL/6 mice. *J. Ethnopharmacol.*, 138(2): 340-344.
- Parrado, J., Miramontes, E., Jover, M., Márquez, J. C., Angeles Mejias, M., Collantes de Teran, L., Absi, E., Bautista, J. 2003. Prevention of brain protein and lipid oxidation elicited by a water-soluble oryzanol enzymatic extract derived from rice bran. *Eur. J. Nutr.*, 42(6): 307-314.
- Paus, R., Stenn, K. S., Link, R. E. 1990. Telogen skin contains an inhibitor of hair growth. *Br. J. Dermatol.*, 122: 777-784.
- Perez, C. A., Wei, Y., Guo, M. 2009. Iron-binding and anti-Fenton properties of baicalein and baicalin. *J. Inorg. Biochem.*, 103(3): 326-332.

- Perez-Marcos, B., Ford, J. L., Armstrong, D. J., Elliott, P. N. C., Rostron, C., Hogan, J. E. 1994. Release of propranolol hydrochloride from matrix tablets containing hydroxypropylmethylcellulose K4M and carbopol 974. *Int. J. Pharm.*, 111(3): 251-259.
- Pillai, G. K., Salim, M. L. D. 1999. Enhanced inhibition of platelet aggregation *in vitro* by niosome-encapsulated indomethacin. *Int. J. Pharm.*, 193(1): 123-127.
- Plessis, J., Egbaria, K., Weiner, N. 1992. Influence of formulation factors on the deposition of liposomal components into the different strata of the skin. *J. Soc. Cosmet. Chem.*, 43: 93-100.
- Prager, N., Bickett, K., French, N., Marcovici, G. 2002. A randomized, double-blind, placebo-controlled trial to determine the effectiveness of botanically derived inhibitors of 5-alpha reductase in the treatment of androgenetic alopecia. *J. Altern. Complement. Med.*, 8: 143-152.
- Prakash, A. 2001. The BP and EP microbial limit test. In: L. Clontz, Microbial limit and bioburden tests: Validation approaches and global requirements. vol.19 CRC press, Boston.
- Prendergast, F. G., Haugland, R. P., Callahan, P. J. 2002. 1-[4-(Trimethylamino)phenyl]-6-phenylhexa-1,3,5-triene: synthesis, fluorescence properties and use as a fluorescence probe of lipid bilayers. *Biochem.*, 20(26): 7333-7338.
- Pyo, H., Yoo, H., Won, C., Lee, S., Kang, Y., Eun, H., Cho, K., KH, K. 2007. The effect of tripeptide-copper complex on human hair growth *in vitro*. *Arch. Pharmacol Res.*, 30: 834-839.

- Raja Naresh, R. A., Chandrashekhar, G., Pillai, G. K., Udupa, N. 1994. Anti-inflammatory activity of niosome encapsulated diclofenac sodium in arthritic rats *Indian J Pharmacol*, 26: 46-48.
- Rajera, R., Nagpal, K., Singh, S. K., Mishra, D. N. 2011. Niosomes: a controlled and novel drug delivery system. *Biol. Pharm. Bull.*, 34: 945-953.
- Rao, A. S., Reddy, S. G., Babu, P. P., Reddy, A. R. 2010. The antioxidant and antiproliferative activities of methanolic extracts from Njavara rice bran. *BMC Comp. Alter. Med.*, 10: 1-9.
- Rho, S.-S., Park, S.-J., Hwang, S.-L., Lee, M.-H., Kim, C. D., Lee, I.-H., Chang, S.-Y., Rang, M.-J. 2005. The hair growth promoting effect of *Asiasari radix* extract and its molecular regulation. *J. Dermatol. Sci.*, 38(2): 89-97.
- Roehrborn, C. G., Marks, L. S., Fenter, T., Freedman, S., Tuttle, J., Gittleman, M., Morrill, B., Wolford, E. T. 2004. Efficacy and safety of dutasteride in the four-year treatment of men with benign prostatic hyperplasia. *Urology*, 63: 709-715.
- Rogers, N. E., Avram, M. R. 2008. Medical treatments for male and female pattern hair loss. *J. Am. Acad. Dermatol.*, 59(4): 547-566.
- Roh, J. S., Han, J. Y., Kim, J. H., Hwang, J. K. 2004. Inhibitory effects of active compounds isolated from safflower (*Carthamus tinctorius* L.) seeds for melanogenesis. *Biol. Pharm. Bull.*, 27(12): 1976-1978.
- Roh, S.-S., Kim, C. D., Lee, M.-H., Hwang, S.-L., Rang, M.-J., Yoon, Y.-K. 2002. The hair growth promoting effect of *Sophora flavescens* extract and its molecular regulation. *J. Dermatol. Sci.*, 30(1): 43-49.

- Rose, D. J., Inglett, G. E., Liu, S. X. 2010. Utilisation of corn (*Zea mays*) bran and corn fiber in the production of food components. *J. Sci. Food Agric.*, 90(6): 915-924.
- Roy, S., Mohanty, A., Dey, J. 2005. Microviscosity of bilayer membranes of some N-acylamino acid surfactants determined by fluorescence probe method. *Chem. Phys. Lett.*, 414(1-3): 23-27.
- Ruksiriwanich, W., Manosroi, J., Abe, M., Manosroi, W., Manosroi, A. 2011. 5 α -Reductase type 1 inhibition of *Oryza sativa* bran extract prepared by supercritical carbon dioxide fluid. *J. Supercrit. Fluids*, 59(1): 61-71.
- Ryan, T. J. 1991. Cutaneous circulation. In: Goldsmith, L. A. Physiology, biochemistry and molecular biology of the skin. Oxford Press, Oxford.
- Sahena, F., Zaidul, I. S. M., Jinap, S., Karim, A. A., Abbas, K. A., Norulaini, N. A. N., Omar, A. K. M. 2009. Application of supercritical CO₂ in lipid extraction - A review. *J. Food Eng.*, 95(2): 240-253.
- Sankar, V., Ruckmani, K., Durga, S., Jailani, S. 2010. Proniosomes as drug carriers. *Pakistan J. Pharm. Sci.*, 23: 103-107.
- Sawada, M., Terada, N., Taniguchi, H., Tateishi, R., Mori, Y. 1987. Cyclosporin A stimulates hair growth in nude mice. *Lab. Invest.*, 56: 684-686.
- Sawaya, M. E., Price, V. H. 1997. Different levels of 5 alpha-reductase type I and II, aromatase, and androgen receptor in hair follicles of women and men with androgenetic alopecia. *J. Invest. Dermatol.*, 109(3): 296-300.
- Sayre, R. N., Saunders, R. M. 1990. Lipid technologies and applications: rice bran and rice bran oil. Marcel Dekker, New York.

Schaefer, H., Lademann, J. 2001. The role of follicular penetration. A differential view. *Skin Pharmacol. Appl. Skin Physiol.*, 14: 23-27.

Schaefer, H., Redelmeier, T. 1996. Skin barrier: principles of percutaneous absorption. Karger, New York.

Scheuplein, R. J. 1967. Mechanism of percutaneous absorption. II. Transient diffusion and the relative importance of various routes of skin penetration. *J. Invest. Dermatol.*, 48: 79-88.

Shanmugam, S., Song, C., Nagayya-Sriraman, S., Baskaran, R., Yong, C., Choi, H., Kim, D., Woo, J., Yoo, B. 2009. Physicochemical characterization and skin permeation of liposome formulations containing clindamycin phosphate. *Arch. Pharmacol. Res.*, 32: 1067-1075.

Shen, Z., Palmer, M. V., Ting, S. S. T., Fairclough, R. J. 1996. Pilot scale extraction of rice bran oil with dense carbon dioxide. *J. Agr. Food Chem.*, 44(10): 3033-3039.

Sherwin, E. R. 1978. Oxidation and antioxidants in fat and oil processing. *J. Am. Oil Chem. Soc.*, 55 809-841.

Shim, J., Seok Kang, H., Park, W.-S., Han, S.-H., Kim, J., Chang, I.-S. 2004. Transdermal delivery of minoxidil with block copolymer nanoparticles. *J. Controlled Release*, 97(3): 477-484.

Shimizu, K., Kondo, R., Sakai, K., Lee, S., Sato, H. 1998. The inhibitory components from *Artocarpus incisus* on melanin biosynthesis. *Planta Medica*, 64: 408-412.

- Shin, H., Won, C., Lee, S., Kwon, O., Kim, K., Eun, H. 2007. Efficacy of 5% minoxidil versus combined 5% minoxidil and 0.01% tretinoin for male pattern hair loss: a randomized, doubleblind, comparative trial. *Am. J. Clin. Dermatol.*, 8: 285-290.
- Sinclair, R. 1998. Male pattern androgenetic alopecia. *Br Med J*, 317: 865-869.
- Singh, R., Arora, S. 2007. Attenuation of free radicals by acetone extract/fractions of *Acacia nilotica* (L.) Willd.Ex del. *J. Chinese Clin. Med.*, 2: 196-204.
- Skalnaya, M. G., Tkachev, V. P. 2011. Trace elements content and hormonal profiles in women with androgenetic alopecia. *J. Trace Elem. Med. Bio.*, 25: S50-S53.
- Solanki, A. B., Parikh, J. R., Parikhb, R. H., Patel, M. R. 2010. Evaluation of different compositions of niosomes to optimize aceclofenac transdermal delivery. *Asian J. Pharm. Sci.*, 5(3): 87-95.
- Suja, K. P., Jayalekshmy, A., Arumughan, C. 2005. Antioxidant activity of sesame cake extract. *Food Chem.*, 91: 213-219.
- Sung-Yum, S., Vinay, K. S., Niti, S. 2003. Mushroom tyrosinase: Recent prospects. *J. Agric. Food Chem.*, 51: 2837-2853.
- Tabbakhian, M., Tavakoli, N., Jaafari, M. R., Daneshamouz, S. 2006. Enhancement of follicular delivery of finasteride by liposomes and niosomes: 1. *In vitro* permeation and *in vivo* deposition studies using hamster flank and ear models. *Int. J. Pharm.*, 323(1-2): 1-10.
- Tachibana, Y., Kikuzaki, H., Hj-Lajis, N., Nakatani, N. 2001. Antioxidant activity of carbazoles from *Murraya koenigii* leaves. *J. Agr. Food Chem.*, 49: 5589-5594.

Tahira, R., Ata-ur, R., Muhammad, A. B. 2007. Characterization of rice bran oil. *J. Agr. Res.*, 45: 225-230.

Takamatsu, Y., Iwata, N., Tsubone, K., Torigoe, K., Endo, T., Sakai, K., Sakai, H., Abe, M. 2009. Synthesis and aqueous solution properties of novel anionic heterogemini surfactants containing a phosphate headgroup. *J. Colloid Interf. Sci.*, 338(1): 229-235.

Teichmann, A., Heuschkel, S., Jacobi, U., Presse, G., Neubert, R. H. H., Sterry, W., Lademann, J. 2007. Comparison of stratum corneum penetration and localization of a lipophilic model drug applied in an o/w microemulsion and an amphiphilic cream. *Eur. J. Pharm. Biopharm.*, 67(3): 699-706.

Teichmann, A., Otberg, N., Jacobi, U., Sterry, W., Lademann, J. 2006. Follicular penetration: development of a method to block the follicles selectively against the penetration of topically applied substances. *Skin Pharmacol. Physiol.*, 19: 216-223.

Teutsch, G., Goubet, F., Battmann, T., Bonfils, A., Bouchoux, F., Cerede, E., Gofflo, D., Gaillard-Kelly, M., Philibert, D. 1994. Non-steroidal antiandrogens: Synthesis and biological profile of high-affinity ligands for the androgen receptor. *J. Steroid Biochem. Mol. Biol.*, 48(1): 111-119.

Thiboutot, D., Harris, G., Iles, V., Cimisi, G., Gilliland, K., Hagari, S. 1995. Activity of the type 1 5 alpha reductase exhibits regional differences in isolated sebaceous glands and whole skin. *J. Invest. Dermatol.*, 105(2): 209-214.

Thielitz, A., Helmdach, M., Röpke, E. M., Gollnick, H. 2001. Lipid analysis of follicular casts from cyanoacrylate strips as a new method for studying therapeutic effects of antiacne agents. *Br. J. Dermatol.*, 145(1): 19-27.

- Titus, M. A., Schell, M. J., Lih, F. B., Tomer, K. B., Mohler, J. L. 2005. Testosterone and dihydrotestosterone tissue levels in recurrent prostate cancer. *Clin. Cancer Res.*, 11(13): 4653-4657.
- Toll, R., Jacobi, U., Richter, H., Lademann, J., Schaefer, H., Blume-Peytavi, U. 2004. Penetration profile of microspheres in follicular targeting of terminal hair follicles. *J. Invest. Dermatol.*, 123: 168-176.
- Torres, J. M., Ortega, E. 2004. Quantitation of mRNA levels of steroid 5[alpha]-reductase isozymes: a novel method that combines quantitative RT-PCR and capillary electrophoresis. *Int. J. Biochem. Cell Biol.*, 36(1): 78-88.
- Touitou, E., Dayan, N., Bergelson, L., Godin, B., Eliaz, M. 2000. Ethosomes-novel vesicular carriers for enhanced delivery: characterization and skin penetration properties. *J. Controlled Release*, 65(3): 403-418.
- Touitou, E., Godin, B., Dayan, N., Weiss, C., Piliponsky, A., Levi-Schaffer, F. 2001. Intracellular delivery mediated by an ethosomal carrier. *Biomaterials*, 22(22): 3053-3059.
- Trüeb, R. M. 2002. Molecular mechanisms of androgenetic alopecia. *Exp. Gerontol.*, 37(8-9): 981-990.
- Tuberoso, C. I. G., Kowalczyk, A., Sarritzu, E., Cabras, P. 2007. Determination of antioxidant compounds and antioxidant activity in commercial oilseeds for food use. *Food Chem.*, 103(4): 1494-1501.
- Uchegbu, I. F., Bouwstra, J. A., Florence, A. T. 1992. Large disk-shaped structures (discomes) in nonionic surfactant vesicle to micelle transitions. *J. Phys. Chem.*, 96(25): 10548-10553.

Uchegbu, I. F., Vyas, S. P. 1998. Non-ionic surfactant based vesicles (niosomes) in drug delivery. *Int. J. Pharm.*, 172(1-2): 33-70.

Van der Molen, R., Spies, F., Van 't Noordende, J., Boelsma, E., Mommaas, A., Koerten, H. 1997. Tape stripping of human stratum corneum yields cell layers that originate from various depths because of furrows in the skin. *Arch. Dermatol. Res.*, 289: 514-518.

VanMiddlesworth, F., Cannell, R. J. P. 1998. Dereplication and partial identification of natural products, *Methods in biotechnology: Natural product isolation*. Humana Press, New Jersey.

Velioglu, Y. S., Mazza, G., Gao, Y. L., Oomah, B. D. 1998. Antioxidant activity and total phenolics in selected fruits, vegetables and grain products. *J. Agr. Food Chem.*, 46: 4113-4117.

Vogt, A., Combadiere, B., Hadam, S., Stieler, K. M., Lademann, J., Schaefer, H., Autran, B., Sterry, W., Blume-Peytavi, U. 2006. 40 nm, but not 750 or 1,500 nm, Nanoparticles enter epidermal CD1a⁺ cells after transcutaneous application on human skin. *J. Invest. Dermatol.*, 126(6): 1316-1322.

Vora, B., Khopade, A. J., Jain, N. K. 1998. Proniosome based transdermal delivery of levonorgestrel for effective contraception. *J. Controlled Release*, 54(2): 149-165.

Vyas, S. P., Khar, R. K. 2002. Targeted and control drug delivery. CBS Publishers and Distributors, New Delhi.

Vyas, S. P., Mysore, N., Jaitely, V., Venkatesan, N. 1998. Discoidal niosome based controlled ocular delivery of timolol maleate. *Pharmazie*, 53(7): 466-469.

- Wadhe, K., Kalsait, R., Umekar, M. 2009. Alternate drug delivery system: recent advancement and future challenges. *Arch. Pharmacol Res.*, 1: 97-105.
- Waters, R. E., White, L. L., May, J. M. 1997. Liposomes containing alpha-tocopherol and ascorbate are protected from external oxidant stress. *Free Radical Res.*, 26(4): 373-379.
- Weisz, G. M., Kammerer, D. R., Carle, R. 2009. Identification and quantification of phenolic compounds from sunflower (*Helianthus annuus* L.) kernels and shells by HPLC-DAD/ESI-MSn. *Food Chem.*, 115(2): 758-765.
- Whiting, D. A. 2001. Advances in the treatment of male androgenetic alopecia: a brief review of finasteride studies. *Eur. J. Dermatol.*, 11: 332-334.
- Williams, C. A., Grayer, R. J. 2004. Anthocyanins and other flavonoids. *Nat. Prod. Rep.*, 21: 539-573.
- Wilton, L., Pearce, G., Edet, E., Freemantle, S., Stephens, M., Mann, R. 1996. The safety of finasteride used in benign prostatic hypertrophy: a non-interventional observational cohort study in 14,772 patients. *Br. J. Urol.*, 78: 379-384.
- Winitchai, S., Manosroi, J., Abe, M., Boonpisuttinant, K., Manosroi, A. 2011. Free radical scavenging activity, tyrosinase inhibition activity and fatty acids composition of oils from pupae of native Thai silkworm (*Bombyx mori* L.). *Kasetsart J Nat Sci*, 45: 404 - 412.
- Wu, H., Ramachandran, C., Weiner, N. D., Roessler, B. J. 2001. Topical transport of hydrophilic compounds using water-in-oil nanoemulsions. *Int. J. Pharm.*, 220: 63-75.
- Yen, G.-C., Duh, P.-D., Su, H.-J. 2005. Antioxidant properties of lotus seed and its effect on DNA damage in human lymphocytes. *Food Chem.*, 89(3): 379-385.

- Yoon, J. I., Al-Reza, S. M., Kang, S. C. 2010. Hair growth promoting effect of *Zizyphus jujuba* essential oil. *Food Chem. Toxicol.*, 48(5): 1350-1354.
- Zhang, W., Sun, Y., Li, Y., Shen, R., Ni, H., Hu, D. 2012. Preparation and influencing factors of sirolimus liposome by supercritical fluid. *Artif. Cells Blood Substit. Biotechnol.*, 40(1-2): 62-65.
- Zhao, W., Shishikura, A., Fujimoto, K., Arai, K., Saito, S. 1987. Fractional extraction of rice bran oil with supercritical carbon dioxide. *Agr. Biol. Chem.*, 51: 1773-1777.
- Zhou, K., Su, L., Yu, L. 2004. Phytochemicals and Antioxidant Properties in Wheat Bran. *J. Agr. Food Chem.*, 52(20): 6108-6114.
- Zidan, A. S., Rahman, Z., Khan, M. A. 2011. Product and process understanding of a novel pediatric anti-HIV tenofovir niosomes with a high-pressure homogenizer. *Eur. J. Pharm. Sci.*, 44: 93-102.