

TABLE OF CONTENTS

	Page
ACKNOWLEDGEMENT	iii
ENGLISH ABSTRACT	iv
THAI ABSTRACT	vi
LIST OF TABLES	xiii
LIST OF FIGURES	xiv
ABBREVIATIONS AND SYMBOLS	xvi
CHAPTER 1 INTRODUCTION	
1.1 Statement and Significance of the problem	1
1.2 Objectives	3
CHAPTER 2 LITERATURE REVIEWS	
2.1 Skin structure and function	
2.1.1 Structure of the skin	5
2.1.2 Functions of the skin	7
2.2 Skin aging	8
2.3 Free radical	10
2.4 Antioxidant	15
2.5 Nanostructured Lipid Carriers	18

2.6 Marigold (<i>Tegetes ereta</i> Linn.)	22
--	----

CHAPTER 3 MATERIALS AND METHODS

3.1 Chemicals	27
3.2 Plants	28
3.3 Instruments	28
3.4 Extraction of marigold flower	29
3.5 Determination of antioxidant activity of marigold flower extract	
3.5.1 Determination of antioxidant activity with 1, 1-diphenyl-2-2picrylhydrazyl (DPPH) radical scavenging method	30
3.5.2 Determination of antioxidant activity with thiobarbituric acid-reactive substance (TBARS) method	31
3.6 Determination of Total Phenolic content in marigold flower extracts	31
3.7 Skin irritation testing in animal	32
3.8 Formulation, characterization, stability test of nanostructured lipid carriers (NLC)	
3.8.1 Formulation of NLC (unloaded NLC)	34
3.8.2 Characterization method for unloaded NLC	35
3.8.3 Stability test of unloaded NLC	36

3.9 Formulation, characterization, stability test and antioxidant activity of marigold flower extract loaded nanostructured lipid carriers (ME-NLC)	
3.9.1 Formulation of marigold flower extract loaded NLC (ME-NLC)	36
3.9.2 Characterization method for marigold flower extract loaded nanostructured lipid carriers (ME-NLC)	37
3.9.3 Stability test of marigold flower extract loaded nanostructured lipid carriers (ME-NLC)	37
3.9.4 Antioxidant activity test of marigold flower extract loaded nanostructured lipid carriers (ME-NLC)	38
3.10 Formulation and stability test of cream base	38
3.11 Formulation, characterization, stability test and antioxidant activity of marigold flower extract loaded nanostructured lipid carriers cream (ME-NLC cream)	
3.11.1 Formulation of marigold flower extract loaded nanostructured lipid carriers cream (ME-NLC cream)	40
3.11.2 Characterization of marigold flower extract loaded nanostructured lipid carriers cream (ME-NLC cream)	40
3.11.3 Stability test of marigold flower extract loaded nanostructured lipid carriers cream (ME-NLC cream)	40
3.11.4 Antioxidant activity test of marigold flower extract loaded nanostructured lipid carriers cream (ME-NLC cream)	41

3.12	Skin irritation testing in human volunteers	41
3.13	Wrinkle reducing capacity test of marigold flower extract loaded nanostructured lipid carriers cream (ME-NLC cream)	43

CHAPTER 4 RESULTS AND DISCUSSION

4.1	Determination of antioxidant activity of marigold flower extracts	
4.1.1	Scavenging effect on 1, 1-diphenyl-2-2picrylhydrazyl (DPPH) radical	46
4.1.2	Thiobarbituric acid reactive substances species (TBARS) assay	47
4.2	Determination of Total Phenolic content in marigold flower extracts	50
4.3	Formulation, characterization and stability of unloaded nanostructured lipid carriers (unloaded NLC)	53
4.4	Formulation, characterization and stability of marigold flower extract loaded nanostructured lipid carriers (ME-NLC)	61
4.5	Selection of good marigold flower extract loaded nanostructured lipid carriers (ME-NLC)	67
4.6	Formulation, stability and selection of good cream base	68
4.7	Formulation and antioxidant activity of marigold flower extract loaded nanostructured lipid carriers cream (ME-NLC cream)	69
4.8	Selection of good marigold flower extract loaded nanostructured lipid carriers cream (ME-NLC cream)	77

4.9	Skin irritation testing in animal	77
4.10	Skin irritation testing in human volunteers	77
4.11	Wrinkle reducing capacity of marigold flower extract loaded nanostructured lipid carriers cream (ME-NLC cream)	81
CHAPTER 5 CONCLUSION		95
REFERENCES		98
APPENDICES		
	APPENDIX A	106
	APPENDIX B	111
	APPENDIX C	113
	APPENDIX D	116
	APPENDIX E	121
CURRICULUM VITAE		129

LIST OF TABLES

Table	Page
2.1 Reactive oxygen species found in vivo	11
3.1 Modified Draize-FHSA Model used in this research	32
3.2 Draize-FHSA Scoring System	33
3.3 Classification of skin irritation	33
4.1 Total phenolic and antioxidant activities of marigold flower extracts and fractions from EA	52
4.2 The mean particle size, polydispersity index (PDI) and zeta potential value of unloaded NLC dispersion after storage for 30, 60 and 90 days	57
4.3 The mean particle size, polydispersity index (PDI) and zeta potential value of marigold flower extract loaded NLC dispersion (ME-NLC) after storage for 30, 60 and 90 days	64
4.4 Primary irritation index (PII) and skin irritation reaction in rabbits	78
4.5 Primary irritation index (PII) and skin irritation reaction in 25 volunteers	80
4.6 The wrinkle reducing capacity on the volume, surface, R_a and R_z after 4 and 8 weeks of treatment	83
4.7 The percentage of satisfaction on test cream before and after use	89


LIST OF FIGURES

Figure	Page
2.1 Diagrams of Skin Layers	4
2.2 Layers of the epidermis	6
2.3 Production of free radicals in human	13
2.4 Antioxidants defenses against free radicals attack	16
2.5 Schematic procedure of hot and cold homogenization techniques	20
2.6 The three types of NLC	22
2.7 <i>Tagetes erecta</i> Linn.	22
3.1 The appearance of three marigold flower extracts	29
3.2 Application sites of wrinkle reducing capacity test	44
4.1 The IC ₅₀ values (µg/ml) of marigold flower extracts and fractions from EA extract by DPPH assay	48
4.2 The IC ₅₀ values (µg/ml) of marigold flower crude extracts and EA fractions by TBARS assay.	50
4.3 The total phenolic contents of marigold flower crude extracts and EA fractions	51
4.4 The physical appearance of selected unloaded NLC; A1, B2 and B6	61
4.5 The physical appearance of selected EA-NLC and F9-NLC	68
4.6 The 8 formulations of marigold cream; 15% EA-NLC cream,	70

30% EA-NLC cream, 15% F9-NLC cream, 30% F9-NLC cream, 0.0225% EA cream, 0.045% EA cream, 0.009%F9 cream and 0.018%F9 creams	
4.7 The antioxidant activity (percent inhibition) of F9 cream at 0, 30, 60 and 90 days after storage at different conditions by DPPH and TBARS assay	72
4.8 The antioxidant activity (percent inhibition) of EA creams at 0, 30, 60 and 90 days after storage at different conditions by DPPH and TBARS assay	75
4.9 Example of skin irritation test in rabbit of 10 substances	79
4.10 Sequence of application on skin irritation test in human	80
4.11 The example of skin irritation test in human with Finn Chamber [®]	81
4.12 The percent Efficiency value on volume, surface, <i>Ra</i> and <i>Rz</i> parameter after 8 weeks of treatment	84
4.13 Skin surface of forearms using Skin-Visiometer [®] SV 600 FW before and after 8 weeks of treatment	88
4.14 The percentage of satisfaction on creams of volunteers	92

ABBREVIATIONS AND SYMBOLS

=	Aequalis, equal to
BA	Behenyl alcohol
CCM	Cyclomethicone
EA	Ethyl acetate
Et	Ethanol
GMS	Glyceryl monostearate
GB	Glyceryl behenate
GC	Glyceryl cocoate
H	Hexane
OD	Octyldodecanol
SA	Stearic acid
S	Surfactant
ME	Marigold flower extract
mV	Millivolt
NLC	Nanostructured lipid carriers
nm	Nanometer
PDI	Polydispersity index
RT	Room temperature
Ra	Arithmetic roughness



Rz	Average roughness
mm ³	Cubic millimeter
°C	degree Celcius
et al	et alibi, and others
e.g.	exempli gratia, for example
IC ₅₀	half maximal inhibitory concentration
H/C	heating-cooling
hr	hour
i.e.	id est, that is to say
<	less than
>	more than
μM	micromolar
mg	milligram
μg	microgram
ml	milliliter
min	minute

ลิขสิทธิ์มหาวิทยาลัยเชียงใหม่

Copyright© by Chiang Mai University
All rights reserved