

REFERENCES

- Alster, T.S. (1999). Cutaneous resurfacing with CO₂ and erbium:YAG lasers—preoperative, intraoperative, and postoperative considerations. *Plast Reconstr Surg*, 103, 619–632.
- Abba, D., Ibabi, H.I., Yakubu, S.E. and Olonitola, O.S. (2009). Phytochemical analysis and antibacterial activity of some powdered herbal preparations marketed in Kaduna metropolis. *Science World Journal*, 4 (1), 23–26.
- Ajayi, I.A. and Oderinde, R.A. (2011). Preliminary Phytochemical Analysis of some Plant Seeds. *Res.J.Chem.Sci.*, 1 (3), 58–62.
- Alam, N., Yoon, K.N., Cha, Y.J., Kim, J.H., Lee, K.R. and Lee, T.S. (2011). Appraisal of the antioxidant, phenolic compounds concentration, xanthine oxidase and tyrosinase inhibitory activities of *Pleurotus salmoneostramineus*. *African Journal of Agricultural Research*, 6 (6), .
- Alster, T.S., Doshi, N.S. and Hopping, S.B. (2004). Combination surgical lifting with ablative laser skin resurfacing of facial skin: a retrospective analysis. *Dermatol Surg*, 30 (9), 1191–1195.
- Ardestani, A. and Yazdanparast, R. (2007). Antioxidant and free radical scavenging potential of *Achillea santolina* extracts. *Food Chem*, 104, 21–29.
- Arican, M. and Ceylan, C. (1999). Metalloproteinases in canine experimental traumatic keratoconjunctivitis. *J Vet Med*, 46, 527–532.

- Arung, E.T., Kusuma, I.W., Christy, E.O., Shimizu, K. and Kondo, R. (2009). Evaluation of medicinal plants from Central Kalimantan for antimelanogenesis. *J Nat Med*, 63 (4), 473–480.
- Aslan, M., Orhan, D.D. and Orhan, N. (2011). Effect of *Gentiana olivieri* on experimental epilepsy models. *Phcog Mag* 7(28), 344–349.
- Astner, S., Wu, A. and Chen, J. (2007). Dietary lutein/zeaxanthin partially reduces photoaging and photocarcinogenesis in chronically UVB-irradiated Skh-1 hairless mice. *Skin Pharmacology and Physiology*, 20 (6), 283–291.
- Aucamp, J.P., Hara, Y. and Apostolides, Z. (2000). Simultaneous analysis of tea catechins, caffeine, gallic acid, theanine and ascorbic acid by micellar electrokinetic capillary chromatography. *J. of Chromatogr. A*, 876, 235–242.
- Bagrov, A.Y. and Shapiro, J.I. (2008). Endogenous digitalis: pathophysiologic roles and therapeutic applications. *Nature Clinical Practice Nephrology* (2008) 4, 378-392, 4, 378–392.
- Barel, A.O. and Clarys, P. (1995). Study of the stratum corneum barrier function by TEWL measurements. *Skin Pharmacol*, 8, 186–195.
- Barry, B.W. and Meyer, M.C. (1979). The rheological properties of carbopol gels I. Continuous shear and creep properties of carbopol gels. *Int J Pharm*, 2, 1-25.
- Beek, T.A.V. (2002). Chemical analysis of Ginkgo biloba leaves and extracts. *Journal of Chromatography A*, 967, 21–55.
- Beurden, P.A.M.S.-V. and Hoff, J.W.V.D. (2005). Zymographic techniques for the analysis of matrix metalloproteinases and their inhibitors. *Biotechniques*, 38 (1), 73–83.

- Blichmann, C.W. and Serup, J. (1988). Assessment of skin moisture. Measurement of electrical conductance, capacitance and transepidermal water loss. *Acta Derm. Venereol.*, 68, 284–290.
- Bonte, F., Dumas, M., Chaudagne, C. and Meybeck, A. (1995). Asiaticoside and madecassoside activity on human fibroblast type I and III collagen secretion. *Ann. Pharm. Fr.*, 53, 38–42.
- Borel, P., Grolier, P., Armand, M., Partier, A., Lafont, H., Lairon, D. and Azais-Braesco, V. (1996). Carotenoids in biological emulsions: solubility, surface-to-core distribution, and release from lipid droplets. *Journal of Lipid Research* 37, 250–261.
- Boron, W.F. and Emile, B.L. (2003). Medical physiology: a cellular and molecular approach. Philadelphia, PA, W.B. Saunders.
- Bradley, D.G. and Min, D.B. (1992). Singlet oxygen oxidation of food. *Crit. Rev. Food Sci. Nutr.*, 31, 211–236.
- Cannell, R.J. (1990). Natural products isolation. Methods in Biotechnology. Totowa, NJ, USA, Humana Press Inc. .
- Carmeliet, P., Moons, L., Herbert, J., Crawley, J., Lupu, F., Lijnen, R. and Collen, D. (1997). Urokinase but not tissue plasminogen activator mediates arterial neointima formation in mice. *Circ Res*, 81, 829–839.
- Castañeda-Ovando, A., Pacheco-Hernandez, M.D.L., P´Aez-Hern´andez, M.E., Rodr´ıguez, J.A. and Gal´an-Vidal, C.A. (2009). Chemical studies of anthocyanins: a review. *Food Chemistry*, 113 (4), 859–871.

- Chairgulprasert, V., Krisornpornsan, B. and Hamad, A. (2006). Chemical constituents of the essential oil and organic acids from longkong (*Aglaia dookkoo* Grif.) fruits. *Songklanakarin J. Sci. Technol.*, 28 (2), 321–326.
- Cho, E.J., Yokozawa, T., Rhyu, D.Y., Kim, H.Y. and Shibahara, N. (2003). The inhibitory effects of 12 medicinal plants and their component compounds on lipid peroxidation. *The American Journal of Chinese Medicine*, 31 (6), 907–917.
- Clarys, P., Alewaeters, K., Lambrecht, R. and Barel, A.O. (2000). Skin color measurements: comparison between three instruments: the Chromameter, the Derma Spectrometer and the Mexameter. *Skin Res Technol*, 6, 230–238.
- Connolly, J.D., Mccrindle, R. and Overton, K.H. (1968). Tetranortriterpenoids V (bicyclonanolides III) the PMR spectrum of Mexicanolide. *Tetrahedron Lett*, 24, 1497–1502.
- Cosgrove, M. (2007). Dietary nutrients intakes and skin-aging appearance among middle-aged American women. *Am.J.Clin.Nutr*, 86, 1225–1231.
- Dahms, G.H. and Zombeck, A. (1995). New formulation possibilities offered by silicone copolyols. *Cosmetics and Toiletries*, 110 (3), 91–96.
- Decker, E.A. and Welch, B. (1990). Role of ferritin as a lipid oxidation catalyst in muscle food. *J Agric Food Chem*, 38, 674–677.
- Demirezer, L.O., Karahan, N., Ucakturk, E., Kuruuzum-Uz, A., Guvenalp, Z. and Kazaz, C. (2011). HPLC fingerprinting of sennosides in laxative drugs with isolation of standard substances from some senna leaves. *Rec. Nat. Prod.*, 5 (4), 261–270.

- Dobarganes, M.C. and Velasco, J. (2002). Analysis of lipid hydroperoxides. *Eur. J. Lipid Sci. Technol.*, 104, 420–428.
- Dobrev, H. (2005). Application of Cutometer area parameters for the study of human skin fatigue. *Skin Res Technol*, 11 (2), 120–122.
- Dolotov, L.E., Sinichkin, Y.P., Tuchin, V.V., Utz, S.R., Altshuler, G.B. and Yaroslavsky, I.V. (2004). Design and evaluation of a novel portable erythema-melanin-meter. *Lasers in Surgery and Medicine*, 34, 127–135.
- Dong, S.-H., Zhang, C.-R., Dong, L., Wu, Y. and Yue, J.-M. (2011). Onoceranoid-Type Triterpenoids from *Lansium domesticum*. *J. Nat. Prod.*, 74, 1042–1048.
- Donghyun, K., Jiyeoun, P., Jinhee, K., Cheolkyu, H., Jeonghyeok, Y., Namdoo, K., Jinho, S. and Choongwan, L. (2006). Flavonoids as mushroom tyrosinase inhibitors: A fluorescence quenching study. *J. Agric. Food Chem*, 54, 935–941.
- Draelos, Z.D. (2001). Botanicals as topical agents. *Clinics Dermatol*, 19(4), 474–477.
- Draize, J., Woodard, G. and Calvery, H. (1944). Methods for the study of irritation and toxicity of substances applied topically to the skin and mucous membranes. *J Pharmacol Exp Ther*, 82, 377–390.
- Dutia, P. (2004). Ethyl Acetate: A Techno-Commercial Profile. *Chemical Weekly*, 184.
- Eder, M. and Mehnert, W. (1998). The importance of concomitant compounds in plant extracts. *Pharmazie*, 53 (5), 285–293.
- Egwaikhide, P.A. and Gimba, C.E. (2007). Analysis of the phytochemical content and anti-microbial activity of *Plectranthus glandulosus* whole plant. *Middle-East Journal of Scientific Research*, 2 (3-4), 135–138.

El-Kamali, H.H. and El-Amir, M.Y. (2010). Antibacterial activity and phytochemical screening of ethanolic extracts obtained from selected sudanese medicinal plants. *Curr. Res. J. Biol. Sci.*, 2 (2), 143-146.

Elsner, P., Wilhelm, D. and Maibach, H. (1990). Mechanical properties of human forearm and vulvar skin. *Br J Dermatol*, 122, 607–614.

Eymard, S., Genot, C. (2003) A modified xylenol orange method to evaluate formation of lipid hydroperoxides during storage and processing of small pelagic fish. *Eur. J. Lipid Sci. Technol*, 105, 497–501.

Fadeel, B. (2005). Raji revisited: cytogenetics of the original Burkitt's lymphoma cell line. *Leukemia*, 19, 159–161.

Fais, A., Corda, M., Era, B., Fadda, M.B., Matos, M.J., Quezada, E., Santana, L., Picciau, C., Podda, G. and Delogu, G. (2009). Tyrosinase Inhibitor Activity of Coumarin-Resveratrol Hybrids. *Molecules*, 14, 2514–2520.

Fisher, G.J., Varani, J. and Voorhees, J.J. (2008). Looking older: fibroblast collapse and therapeutic implications. *Archives of Dermatology*, 144 (5), 666–672.

Florence, A.T. and Attwood, D. (2006). *Physicochemical Principles of Pharmacy* Fourth ed. UK, Pharmaceutical Press.

Fricker S.P., Buckley R.G. (1996). Comparison of two colorimetric assays as cytotoxicity endpoints for an in vitro screen for antitumour agents. *Anticancer Res.*, 16, 3755–3760.

Galloway, J.A., Koester, K.J., Paasch, B.J. and Macosko, C.W. (2004). Effect of sample size on solvent extraction for detecting cocontinuity in polymer blends.

Polymer, 45, 423–428.

- Garrett, E.R. (1962). Prediction of stability of drugs and pharmaceutical preparations. *Journal of Pharmaceutical Sciences*, 51, 811–833.
- Gaspar, L.R., Camargo Jr, F.B., Gianeti, M.D. and Maia Campos, P.M.B.G. (2008). Evaluation of dermatological effects of cosmetic formulations containing *Saccharomyces cerevisiae* extract and vitamins. *Food and Chemical Toxicology*, 46 (11), 3493–3500.
- Giral, F. (1947). The stability of vitamin C in aqueous solutions and in pharmaceutical preparations. *Journal of the American Pharmaceutical Association (Scientific Edition)*, 36, 82–84.
- Githinji, C.G., Mbugua, P.M., Kanui, T.I. and Kariuki, D.K. (2012). Phytochemical and analgesic evaluation of *Mondia whytei* (hook.f) root. *Journal of Pharmacognosy and Phytotherapy*, 4 (3), 26–32.
- Giusti, M.M. and Wrolstad, R.E. (2003). Acylated anthocyanins from edible sources and their applications in food systems. *Biochemical Engineering Journal*, 14 (3), 217–225.
- Glaser, D. (2004). Anti-aging products and cosmeceuticals. *Facial Plast Surg Clin North Am*, 12, 363–372.
- Goin, J.M. and Goin, M.K. (1989). Changing the body: psychological effects of plastic surgery. Baltimore, Williams & Wilkins.
- Gram, F., Brandstetter, H., D'alò, S., Geppert, D., Krell, H.W., Leinert, H., Livi, V., Menta, E., Oliva, A. and Zimmermann, G. (2001). Pyrimidine-2,4,6-Triones: a new effective and selective class of matrix metalloproteinase inhibitors. *Biol. Chem.*, 382, 1277–1285.

Grossbart, T. and Sarwer, D. (1999). Cosmetic surgery: surgical tools – psychosocial goals. *Cutan Med Surg*, 18 (101)

Guangyi, W., Weiping, T. and Robert, R.B. (2005). Bioactive terpenoids as pharmaceutical agents. Natural products drug discovery and therapeutic medicine. L. Zhang. Totowa, New Jersey, Humana Press: 197–227.

Gulcin, I. (2006). Antioxidant and antiradical activities of L-carnitine. *Life Sci*, 78, 803–811.

Habaguchi, K., Watanabe, M., Nakadaira, Y., Nakanishi, K., Kaing, A.K. and Lim, F.L. (1968). The full structures of lansic acid and its minor congener, an unsymmetric onoceradienedione. *Tetrahedron Lett*, 34, 3731–3734.

Hagerman, A.E. and Butler, L.G. (1989). Choosing appropriate methods and standards for assaying tannin. *J. Chem. Ecol.*, 15, 1795–1810.

Handa, S.S., Khanuja, S.P.S., Longo, G. and Rakesh, D.D. (2008). Extraction technologies for medicinal and aromatic plants. Italy, United Nations Industrial Development Organization and the International Centre for Science and High Technology.

Harborne, J.B. (2005). Phytochemical methods. analysis. a guide to modern techniques of plant., Springer International, India.

Harborne, J.B. and Williams, C.A. (2000). Advances in flavonoid research since 1992. *Phytochemistry*, 55, 481-504.

Heinrich, U., Neukam, K., Tronnier, H., Sies, H. and Stahl, W. (2011). Long-Term Ingestion of High Flavanol Cocoa Provides Photoprotection against UV-Induced Erythema and Improves Skin Condition in Women¹. *J. Nutr.*, 141 (6), 1202-1208.

Herrero, M., Ez, E.I.N., Sen~Ora'Ns, J. and Cifuentes, A. (2004). Pressurized liquid extracts from *Spirulina platensis* microalga: Determination of their antioxidant activity and preliminary analysis by micellar electrokinetic chromatography. *J. Chromatogr. A*, 1047, 195–203.

Heyne, K. (1987). *Tumbuhan Berguna Indonesia II*. Jakarta, Badan Litbang Departement Kehutanan.

Hideaki, O. (2005). Purification by solvent extraction using partition coefficient. *Methods in Biotechnology*, 20, 269–273.

Hirose, M., Takesada, Y., Tanaka, H., Tamano, S., Kato, T. and Shirai, T. (1997). *Carcinogenesis*, 19 (1), 27.

Hostettman, K. and Marston, A. (1995). *Saponins*, Cambridge University Press, United Kingdom.

Ito, N., Hirose, M., Fukushima, S., Shirai, T. and Tatematsu, M. (1986). *Food Chem. Toxicol.*, 24 (10-11), 1071.

Jennifer, S., David, A., Grady, G. and Tonalee, C. (1988). Aqueous thermal degradation of gallic acid. *Geochim Cosmochim Acta*, 52, 341–344.

Jiang, Z.Y., Woollard, A.C.S. and Wolff, S.P. (1991). *Lipids*, 26, 853–856.

Jing, P. and Giusti, M.M. (2005). Characterization of Anthocyanin-Rich Waste from Purple Corncobs (*Zea mays* L.) and Its Application to Color Milk. *J Agric Food Chem*, 53, 8775–8781.

Judis, M.A., Sturla, M.A., Doval, M.M., Romero, A.M. and Fernández, C.L. (2010). Influence of meat components on antioxidant activity of beef sarcoplasmic proteins/malondialdehyde reaction products in model emulsions. *Electronic Journal of Environmental, Agricultural and Food Chemistry*, 9 (3), 636–645.

Kasolo, J.N., Bimenya, G.S., Ojok, L., Ochieng, J. and Ogwal-Okeng, J.W. (2010). Phytochemicals and uses of *Moringa oleifera* leaves in Ugandan rural communities. *Journal of Medicinal Plants Research* 4(9), 753–757.

Keepers, Y.P., Pizao, P.E., Peters, G.J., Arke-Otte, J., Winograd, B. and Pinedo, H.M. (1991). *Eur. J. Cancer*, 27, 897–900.

Khan, B.A. and Akhtar, N. (2012). Phytochemical analysis and acute toxicity tests of two medicinal plant extracts. *Journal of Medicinal Plants Research*, 6 (19), 3545–3548.

Khongsuwan, P. (2005). Detection of Apomixis in *Lansium domesticum* Corr. Department of Plant Science, Faculty of Natural Resources University. Songkla, Prince of Songkla University.

Kiang, A.K., Tan, E.L., Lim, F.Y., Habaguchi, K., Nakanishi, K., Fachan, L. and Ourisson, G. (1967). Lansic acid, a bicyclic triterpene. *Tetrahedron Lett.*, 37, 3571–3574.

Kim, I., Jeong, S.-J., Kim, E.-S., Kim, S.H. and Moon, A. (2007). Type I Collagen-induced Pro-MMP-2 Activation is Differentially Regulated by H-Ras and N-Ras in Human Breast Epithelial Cells. *Journal of Biochemistry and Molecular Biology*, 40 (5), 825–831.

Kim, S., Kim, Y., Kim, J., Cho, K. and Chung, J. (2007). Berberine inhibits TPA-induced MMP-9 and IL-6 expression in normal human keratinocytes. *Phytomedicine*, 15, 340–347.

Krasnovsky, A.A. (1953). The primary processes of photosynthesis in plants. *Plant Biotechnology*, 11, 363–410.

Krishnamurthy, K.H. (2001). Ayurvedic technical studies and herbal cosmetics of ancient India. New Delhi, Vedams Books Ltd.

Kujur, R., Singh, V., Ram, M., Yadava, H., Singh, K., Kumari, S. and Roy, B. (2010). Antidiabetic activity and phytochemical screening of crude extract of *Stevia rebaudiana* in alloxan-induced diabetic rats. *Phcog Res*, 2, 258–263.

Kumar, A., Rajput, G., Dhatwalia, V.K. and Srivastav, G. (2009). Phytocontent screening of mucuna seeds and exploit in opposition to pathogenic microbes. *J. Biol. Environ. Sci.*, 3 (9), 71–76.

Kumguan, K., Manosroi, J., Jantrawut, P., Pankasemsuk, T., Manosroi, W. and Manosroi, A. (2011). Antioxidant activities of aqueous extract from leaves of Longkong (*Lansium domesticum* Corr.). 3E IFSCC 2011 Conference, Bangkok, Thailand.

Lavker, R.M. (1995). Cutaneous aging:chronologic versus photoaging. In: *Photodamage* Cambridge, MA, Blackwell Science.

Lazarowych, N.J. and Pekos, P. (1998). Fingerprinting and marker compounds for identifying and standardizing botanicals. *Drug Information Journal*, 32, 497–512.

Leaman, D.J. (1995). Malaria remedies of the Kenyah of the Apo Kayan, East Kalimantan, Indonesian Borneo: A quantitative assessment of local consensus as an indicator of biological efficacy. *J Ethnopharmacol*, 49, 1–16.

Lee, S.M., Olsen, J.I., Schweizer, M.P. and Klocke, J.A. (1988). 7-Deacetyl-17b-hydroxyazadione, a new limonoid insect growth inhibitor from *Azadirachta indica*. *Phytochemistry*, 27, 2773–2775.

Leonhardt, H. (1990). Histologie. Zytologie and Mikroanatomie des Menschen, Thieme. 3.

Letelier, M., Berríos, A., Troncoso, J., Sandoval, J., Holst, M., Palma, K., Montoya, M., Miranda, D. and Lira, V. (2008). DPPH and oxygen free radicals as pro-oxidant of biomolecules. *Toxicol In Vitro*, 22, 279–286.

Li, Z., Wang, B., Zhang, Q., Qu, Y., Xu, H. and Li, G. (2012). Preparation and antioxidant property of extract and semipurified fractions of *Caulerpa racemosa*. *J Appl Phycol*, 1–10.

Lim, Y.Y., Lim, T.T. and Tee, J.J. (2007). Antioxidant properties of several tropical fruits: A comparative study. *Food Chem*, 103, 1003–1008.

Liyana-Pathirana, C. and Shahidi, F. (2007). Antioxidant and free radical scavenging activities of whole wheat and milling fractions. *Food Chem*, 101, 1151–1157.

Lopes, G.C., Longhini, R., Santos, P.V.P.D., Araujo, A.A.S., Bruschi, M.L. and Demello, J.A.C.P. (2011). Preliminary assessment of the chemical stability of dried extracts from *Guazuma ulmifolia* Lam. (Sterculiaceae). *International Journal of Analytical Chemistry*, 2012, 1–7.

Lopez-Avila, V. and Spencer, J.V. (2008). Methods for detection of matrix metalloproteinases as biomarkers in cardiovascular disease. *Clinical Medicine: Cardiology*, 2, 75–87.

Luque-Garciá, J.L. and Castro, M.D. (2003). Ultrasound: A powerful tool for leaching. *TrAC-Trend. Anal. Chem.*, 22, 41–47.

Ma, J., Lü, X. and Huang, Y. (2011). Genomic Analysis of Cytotoxicity Response to Nanosilver in Human Dermal Fibroblasts. *J. Biomed. Nanotechnol.*, 7, 263–275.

Maci'as-Sa'Nchez, M.D., Mantell, C. and Rodri'Guez, M. (2005). Supercritical fluid extraction of carotenoids and chlorophyll a from *Nannochloropsis gaditana*. *J. Food Eng.*, 66, 245–251.

Manosroi, A., Jantrawut, P., Akihisa, T., Manosroi, W. and Manosroi, J. (2010). *In vitro* anti-aging activities of *Terminalia chebula* gall extract. *Pharmaceutical Biology*, 48 (4), 469–481.

Manosroi, A., Jantrawut, P., Akihisa, T., Manosroi, W. and Manosroi, J. (2011). *In vitro* and *in vivo* skin anti-aging evaluation of gel containing niosomes loaded with a semi-purified fraction containing gallic acid from *Terminalia chebula* galls. *Pharmaceutical Biology*, 49 (11), 1190–1203.

Manosroi, A., Podjanasoonthon, K. and Manosroi, J. (2002). Stability and release of topical tranexamic acid liposome formulations. *J Soc Cosmet Chem*, 53, 375–386.

Martha, T., Wih, W.L., Ranti, A.S., Wasitaatamadja, S.M., Suryninsih, F.D. Junardy and Maily (2007). In Search of Naturally Derived Whitening agent-Pragmatic approach. Asian Societies of Cosmetic Scientists 8th Conference ASCS 2007. Delivering Science to the Depths of Asian Skin. Singapore: 116–117.

Martha, T., Wih, W.L., Ranti, A.S., Wasitaatamadja, S.M., Suryninsih, F.D. Junardy and Maily (2007). Review of *Lansium domesticum* Correa and its use in cosmetics. *BLACPMA*, 7 (4), 183-189.

Martin, P.D. (1993). Sonochemistry in industry. Progress and prospects. *Chem. Ind-London.*, 1993, 233–236.

- Martins, A.M., Santos, M.I., Azevedo, H.S., B., P., Malafaya and Reis, R.L. (2008). Natural origin scaffolds with in situ pore forming capability for bone tissue engineering applications. *Acta Biomaterialia*, 4, 1637–1645.
- Maw-Rong, L., Chueh-Yu, L., Zu-Guang, L. and Tzu-Feng, T. (2006). *Chromatographia*, 1120, 244–251.
- Mayanti, T., Soidah, S., Natawigena, W.D., Supratman, U. and Tjokronegoro, D.R.-A. (2007). Antibacterial Terpenoid from The Bark of *Lansium domesticum* Corr cv. Kokossan (Meliaceae). *Medicine and Medicinal Plants*. Surabaya.
- Mccloud, T.G. (2010). High throughput extraction of plant, marine and fungal specimens for preservation of biologically active molecules. *Molecules*, 15, 4526–4563.
- Mercer, P. and Armenta, R.E. (2011). Developments in oil extraction from microalga. *Eur. J. Lipid Sci. Technol.*
- Michael, J. (2004). Quinoline, quinazoline and acridone alkaloids. *Nat. Prod. Rep.*, 21, 650–668.
- Michel, V.D.R. and Garrone, R. (1991). Collagen family of proteins. *FASEB J*, 5, 2814-2823.
- Milner, J.M., Rowan, A.D., Cawston, T.E. and Young, D.A. (2006). Metalloproteinase and inhibitor expression profiling of resorbing cartilage reveals pro-collagenase activation as a critical step for collagenolysis. *Arthritis Research & Therapy*, 8 (5), 1–12.
- Minaker, K.L. (2007). *Common clinical sequelae of aging*. Philadelphia, Saunders Elsevier.

Mitra, S.K., Walters, B.T., Clouse, S.D. and Goshe, M.B. (2009). An efficient organic solvent based extraction method for the proteomic analysis of Arabidopsis plasma membranes. *Journal of Proteome Research*, 8, 2752–2767.

Mohanlall, V., Steenkamp, P. and Odhav, B. (2011). Isolation and characterization of anthraquinone derivatives from *Ceratopthea triloba* (Bernh.). *Journal of Medicinal Plants Research* 5(14), 3132–3141.

Monheit, G.D. and Chastain, M.A. (2001). Chemical peels. *Facial Plast Surg Clin North Am*, 9, 239–255.

Morton, J. (1987). In: *Fruits of Warm Climates*. Miami, Florida, Julia F. Marton.

Nantitanon, W., Yotsawimonwat, S. and Okonogi, S. (2010). Factors influencing antioxidant activities and total phenolic content of guava leaf extract. *LWT - Food Science and Technology*, 43, 1095–1103.

Newman, A.W. and Stahly, G.P. (2002). Form selection of pharmaceutical compounds. *Handbook of pharmaceutical analysis*. L. Ohannesian and A. J. Streeter. New York, Marcel Dekker Inc.: 18-74.

Nishizawa, M., Emura, M., Yamada, H., Shiro, M., Hayashi, Y. and Tokuda, H. (1989). Isolation of a new cycloartanoid triterpene from leaves of *Lansium domesticum* novel skin-tumor promotion inhibitor. *Tetrahedron Letters*, 30 (41), 5615–5618.

Nishizawa, M., Nademoto, Y., Sastrapradja, S., Shiro, M. and Hayashi, Y. (1985). The Structure of Dukunolide A: A Tetranortriterpenoid with a New Carbon Skeleton from *Lansium domesticum*. *J. Chem. Soc., Chem. Commun.*, 395–396.

Nishizawa, M., Nademoto, Y., Sastrapradja, S., Shiro, M. and Hayashi, Y. (1985).

Structure of dukunolides, bitter principles of *Lansium domesticum*. *J. Org Chem*, 50, 5487–5490.

Nishizawa, M., Nademoto, Y., Sastrapradja, S., Shiro, M. and Hayashii, Y. (1988).

Dukunolide D, E and F: New tetranortriterpenoids from the seed of *Lansium domesticum*. *Phytochemistry*, 27 (1), 237–239.

Nishizawa, M., Nishide, H. and Hayashi, Y. (1984). Total synthesis of (\pm)- α,γ -

onoceradienedione and lansic acid. *Tetrahedron Lett*, 25, 5071-5074.

Nishizawa, M., Nishide, H., Hayashi, Y. and Kosela, S. (1982). The structure of

lansioside A: a novel triterpene glycoside with amino-sugar from *Lansium domesticum*. *Tetrahedron Lett*, 23, 1349–1350.

Nishizawa, M., Nishide, H., Kosela, S. and Hayashi, Y. (1983). Structure of

lansiosides: biologically active new triterpene glycosides from *Lansium domesticum*. *J. Org Chem*, 48, 4462–4466.

Nishizawa, M., Nishide, H., Kuriyama, K. and Hayashi, Y. (1986). *Chem Pharm Bull*,

34, 4443–4446.

Nobakht, G.M., Kadir, M.A. and Stanslas, J. (2010). Analysis of preliminary

phytochemical screening of *Typhonium flagelliforme*. *African Journal of Biotechnology*, 9 (11), 1655-1657.

Okorie, D.A. and Taylor, D.A.H. (1968). Extractives from the seed of *Cedrela*

odorata L. *Phytochemistry*, 7, 1683–1686.

Omar, S., Marcotte, M., Fields, P., Sanchez, P.E., Poveda, L., Mata, R., Jimenez, A., Durst, T., Zhang, J., Mackinnon, S., Leaman, D., Arnason, J.T. and Philogene, B.J.R. (2007). Antifeedant activities of terpenoids isolated from tropical Rutales. *Journal of Stored Products Research* 43, 92–96.

Omura S, Enomoto Y, Shinose M, Takahashi Y, Iwai Y, Shiomi K. (1999). Isolation and structure of a new antibiotic viridomycin F produced by *Streptomyces* sp. K96-0188. *J Antibiot*, 52, 61–64.

Onwukaeme, D.N., Ikuegbvweha, T.B. and Asonye, C.C. (2007). Evaluation of phytochemical constituents, antibacterial activities and effect of exudates of *Pycnanthus angolensis* Weld Warb (Myristicaceae) on corneal ulcers in rabbits. *Trop J Pharm Res*, 6, 725–730.

Osawa, T. and Namiki, M. (1981). A novel type of antioxidant isolated from leaf wax of *Eucalyptus* leaves. *Agric Biol Chem*, 45, 735-739.

Özgen, U., Mavi, A., Terzi, Z., Kazaz, C., Asçı, L., Kaya, Y. and Seçen, H. (2011). Relationship Between Chemical Structure and Antioxidant Activity of Luteolin and Its Glycosides Isolated from *Thymus*. *Rec. Nat. Prod.*, 5 (1), 12-21.

Papazisis, K.T., Geromichalos, G.D., Dimitriadis, K.A. and Kortsaris, A.H. (1997). Optimization of the sulforhodamine B colorimetric assay. *J Immunol Methods*, 208, 151–158.

Patterson, M., Atkinson, S., Knuper, V. and Murphy, G. (2001). Specific collagenolysis by gelatinase A, MMP-2, is determined by the hemopexin domain and not the fibronectin-like domain. *FEBS Lett*, 503, 158-162.

Pawliszyn, J. (1993). Kinetic model of supercritical fluid extraction. *J. Chromatogr. Sci.*, 31, 31–37.

Philips, N. (2010). Counteraction of skin inflammation and aging or cancer by polyphenols, Bentham Science.

Philips, N., Conte, J. and Chen, Y.J. (2009). Beneficial regulation of matrix metalloproteinases and their inhibitors, fibrillar collagens and transforming growth factor- β by *Polypodium leucotomos*, directly or in dermal fibroblasts, ultraviolet radiated fibroblasts, and melanoma cells. *Dermatological Research*, 301 (7), 487–495.

Philips, N., Keller, T. and C.Hendrix (2007). Regulation of the extracellular matrix remodeling by lutein in dermal fibroblasts, melanoma cells, and ultraviolet radiation exposed fibroblasts. *Archives of Dermatological Research*, 299 (8), 373–379.

Philips, N., Samuel, M. and Arena, R. (2010). Direct inhibition of elastase and matrix metalloproteinases and stimulation of biosynthesis of fibrillar collagens, elastin, and fibrillins by xanthohumol. *Journal of Cosmetic Science*, 61 (2), 125–132.

Philips, N., Smith, J., Keller, T. and Gonzalez, S. (2003). Predominant effects of *Polypodium leucotomos* on membrane integrity, lipid peroxidation, and expression of elastin and matrix metalloproteinase-1 in ultraviolet radiation exposed fibroblasts, and keratinocytes. *Journal of Dermatological Science*, 32 (1), 1–9.

- Pinnagoda, J. and Tupker, R.A. (1995). Measurement of transepidermal water loss. In: Noninvasive Methods and the Skin. J. Serup and G. B. E. Jemec. Boca Raton, FL, CRC Press: 173–178.
- Pongtip, S., Roongtawan, S. and Wandee, G. (2005). Antioxidant activity of Siamese neem tree (VP1209). *Journal of Ethnopharmacology*, 99, 109–122.
- Prakash, A. (2001). The BP and EP microbial limit test. CRC press.
- Reische, D.W., Lillard, D.A. and Eitenmiller, R.R. (1998). Food Lipids: Chemistry. New York, Marcel Dekker, Inc.
- Ricci, F., Berardi, V. and Risuleo, G. (2009). Differential Cytotoxicity of MEX: a Component of Neem Oil Whose Action Is Exerted at the Cell Membrane Level. *Molecules* 14, 122-132.
- Rice-Evans, C., Miller, N. and Paganga, G. (1996). Structure-antioxidant activity relationships of flavonoids and phenolic acids. *Free Radical Bio. Med.*, 20, 933–956.
- Ruksiriwanich, W., Manosroi, J., Abe, M., Manosroi, W. and Manosroi, A. (2011). 5alpha-reductase type 1 inhibition of *Oryza sativa* bran extract prepared by supercritical carbon dioxide fluid. *J. of Supercritical Fluids*, 59, 61–71.
- Ryan-Stoneham, T. and Tong, C.H. (2000). Degradation kinetics of chlorophyll in peas as a function of pH. *Journal of Food Science*, 65, 1296–1302.
- Saewan, N., John, D.S. and Kan, C. (2006). Antimalarial tetranortriterpenoid from the seeds of *Lansium domesticum* Corr. *Phytochemistry*, 67, 2288-2293.
- Sam, Z. and Chhandak, B. (2008). Plant terpenoids: applications and future potentials. *Biotechnology and Molecular Biology Reviews*, 3 (1), 1-7.

Serup, J. and Jemec, G. (1995). Handbook of Non-Invasive Methods and The Skin. London, UK, CRC Press.

Shahidi, F. and Wanasundara, P.K. (1992). Phenolic antioxidants. *Crit. Rev. Food Sci. Nutr.*, 32, 67–103.

Shahidi, F. and Zhong, Y. (2005). Antioxidants: Regulatory Status. Bailey's Industrial Oil and Fat Products. Canada, John Wiley and Sons, Inc. 491-509.

Sherma, J. (2000). Thin-layer chromatography in food and agricultural analysis. *Journal of Chromatography A*, 880, 129–147.

Sherwin, E. (1978). Oxidation and antioxidants in fat and oil processing. *Journal of the American Oil Chemists' Society*, 55 (11), 809–814.

Shimada, K., Fujikawa, K., Yahara, K. and Nakamura, T. (1992). Antioxidative properties of xanthan on the autoxidation of soybean oil in cyclodextrin emulsion. *J Agric Food Chem*, 40, 945–948.

Shimizu, K., Kondo, R., Sakai, K., Lee, S. and Sato, H. (1998). The inhibitory components from *Artocarpus incisus* on melanin biosynthesis. *Planta Medica*, 64, 408–412.

Siddiqui, B.S. and Rasheed, M. (2001). Three New Triterpenoids from *Azadirachta indica*. *Helv. Chim. Acta*, 84, 1962.

Simon, M. (1993). The Systematic Distribution of Tannins in the Leaves of Angiosperms: A Tool for Ecological Studies. *Biochemical Systematics and Ecology*, 21 (8), 833–846.

Singh, P., Shukla, R., Prakash, B., Kumar, A., Singh, S., Mishra, P.K. and Dubey, N.K. (2010). Chemical profile, antifungal, antiaflatoxic and antioxidant activity of *Citrus maxima* Burm. and *Citrus sinensis* (L.) Osbeck essential oils and their cyclic monoterpene, dl-limonene. *Food and Chemical Toxicology*, 48 (6), 1734–1740.

Singh, P., Shukla, R., Prakash, B., Kumar, A., Singh, S., Mishra, P.K. and Dubey, N.K. (2010). Chemical profile, antifungal, antiaflatoxic and antioxidant activity of *Citrus maxima* Burm. and *Citrus sinensis* (L.) Osbeck essential oils and their cyclic monoterpene, dl-limonene. *Food and Chemical Toxicology*, 48 (6), 1734–1740.

Singh, S. and Nalwa, H.S. (2007). Nanotechnology and health safety - Toxicity and risk assessments of nanostructured materials on human health. *J. Nanosci. Nanotechnol.*, 7 (9), 3048–3070.

Singhatong, S., Leelarungrayu, D. and Chaiyasut, C. (2010). Antioxidant and toxicity activities of *Artocarpus lakoocha* Roxb. heartwood extract. *Journal of Medicinal Plants Research*, 4 (10), 947–953.

Sirikatitham, A., Chuchom, T. and Itharat, A. (2007). Development of the chromatographic fingerprint analysis of dioscorealides and dioscoreanone from *Dioscorea membranacea* Pierre. *Songklanakarin J. Sci. Technol.*, 9 (1), 101–107.

Skehan, P., Storeng, R., Scudiero, D., Monks, A., McMahon, J., Vistica, D., Warren, J.T., Bokesch, H., Kenney, S. and Boyd, M.R. (1990). *J. Natl. Cancer Inst.*, 82, 1107–1112.

- Soong, Y.Y. and Barlow, P.J. (2006). Quantification of gallic acid and ellagic acid from longan (*Dimocarpus longan* Lour.) seed and mango (*Mangifera indica* L.) kernel and their effects on antioxidant activity. *Food Chemistry*, 97, 524–530.
- Spira, M., Gerow, F. and Hardy, S. (1974). Complications of chemical face peeling. *Plast Reconstr Surg*, 54, 397–403.
- Stegman, S.J. (1982). A comparative histologic study of the effects of three peeling agents and dermabrasion on normal and sundamaged skin. *Aesthetic Plast Surg*, 6, 123–135.
- Strongin, A.Y., Collier, I., Bannikov, G., Marmer, B.L., Grant, G.A. and Goldberg, G.I. (1995). Mechanism of cell surface activation of 72-kDa type IV collagenase. Isolation of the activated form of the membrane metalloprotease. *J. Biol. Chem.*, 270, 5331–5338.
- Suja, K., Jayalekshmy, A. and Arumughan, C. (2005). Antioxidant activity of sesame cake extract. *Food Chem*, 91, 213–219.
- Sultana, B., Anwar, F. and Ashraf, M. (2009). Effect of extraction solvent/technique on the antioxidant activity of selected medicinal plant extracts. *Molecules*, 14, 2167–2180.
- Sun, T., Xu, Z., Godber, J.S. and Prinyawiwatkul, W. (2006). Capabilities of oat extracts in inhibiting cholesterol and long chain fatty acid oxidation during heating. *Cereal Chem.*, 83, 451–454.
- Sung-Yum, S. and Niti, K.V.S. (2003). Mushroom tyrosinase: Recent prospects. *J Agric Food Chem*, 51, 2837–2853.

Suwannalert, P., Povichit, N., Puchadapirom, P. and Junking, M. (2012). Anti-aging activity and non-toxic dose of phytooxyresveratrol from *Artocarpus lakoocha* Roxb. *Trop J Pharm Res*, 11 (1), 69–74.

Tachakittirungrod, S., Okonogi, S. and Chowwanapoonpohn, S. (2007). Study on antioxidant activity of certain plants in Thailand: Mechanism of antioxidant action of guava leaf extract. *Food Chemistry*, 103, 318–388.

Tachibana, Y., Kikuzaki, H., Hj-Lajis, N. and Nakatani, N. (2001). Antioxidant activity of carbazoles from *Murraya koenigii* leaves. *J Agric Food Chem*, 49, 5589–5594.

Takema, Y., Yorimoto, Y., Kawai, M. and Imokawa, G. (1994). Age related changes in the elastic properties and thickness of human facial skin. *Br J Dermatol*, 131, 641–648.

Tanaka, T., Ishibashi, M., Fujimoto, H., Koyano, E.O.T., Kowithayakorn, T., Hayashi, M. and Komiyama, K. (2002). New onoceranoid triterpene constituents from *Lansium domesticum*. *J. Nat. Prod.*, 65, 1709–1711.

Tanaka, T., Ishibashi, M., Fujimoto, H., Okuyama, E., Koyano, T., Kowithayakorn, T., Hayashi, M. and Komiyama, K. (2002). New Onoceranoid Triterpene Constituents from *Lansium domesticum*. *journal of Natrural Products*, 65, 1709–1711.


Tengamnuay, P., Pengrungruangwong, K., Pheansri, I. and K. Likhitwitayawuid International Journal of Cosmetic Science, D.J.-X. (2006). *Artocarpus lakoocha* heartwood extract as a novel cosmetic ingredient: evaluation of the in vitro anti-tyrosinase and in vivo skin whitening activities. *International Journal of Cosmetic Science*, 28, 269–276.

- Tepe, B. and Sokman, A. (2007). Screening of the antioxidative properties and total phenolic contents of three endemic *Tanacetum* subspecies from Turkish flora. *Bioresource Technology*, 98 3076–3079.
- Thomas, G.T., Lewis, M.P. and Speight, P.M. (1999). Matrix metalloproteinases and oral cancer. *Oral Oncology*, 35 (3), 227–233.
- Tilaar, M., Wong, L., Ranti, A., Setiawati, E., Wasitaatmadja, S. and Maily, S. (2007a). Penelitian Bahan Pencerah dan Pelembab Kulit dari Tanaman. Kongres ISFI XV, Jakarta.
- Torskangerpoll, K. and Andersen, O.M. (2005). Colour stability of anthocyanin in aqueous solutions at various pH values. *Journal of Food Chemistry*, 89, 427–440.
- Touchstone, J.C. and Dobbins, M.F. (1978). Practical of Thin Layer Chromatography. New York, John Wiley Sons.
- Trueb, R. (2005). Aging of skin & hair. *Ther Umsch*, 62, 837.
- Tsai, T. and Lee, M. (2008). *Chromatographia*, 67 (5-6), 425–431.
- Tsuji, N., Moriwaki, S., Suzuki, Y., Takema, Y. and Imokawa, G. (2001). The role of elastases secreted by fibroblasts in wrinkle formation: Implication through selective inhibition of elastase activity. *Photochem Photobiol*, 74, 283–290.
- Ueda, C.T., Shah, V.P., Derdzinski, K., Ewing, G., Flynn, G., Maibach, H., Marques, M., Rytting, H., Shaw, S., Thakker, K. and Yacobi, A. (2009). Topical and Transdermal Drug Products. *Pharmacoepial Forum*, 35 (3), 750–764.
- Vanni, A., Gasraldi, D. and Giunata, G. (1990). Kinetic investigation on the double enzyme activity of the tyrosinase mushroom. *Annalli di Chimica*, 80, 35–60.

- Varani, J., Quan, T.H. and Fisher, G.J. (2011). Mechanisms and pathophysiology of photoaging and chronological skin aging. *Aging Skin: Current and Future Therapeutic Strategies*. Mumbai, Dattini book agency, 1–5.
- Vartanian, A.J. and Dayan, S.H. (2004). Facial rejuvenation using botulinum toxin A: review and update. Chicago, IL, U.S.A, Thiem Medical Publishers., Inc.
- Vasek, F.C. (1980). Creosote Bush: Long-Lived Clones in the Mojave Desert. *American Journal of Botany*, 67 (2), 246,255.
- Verheij, E. and Coronel, R. (1992). Edible fruits and nuts. *Plant Resources of South-East Asia II*. Bogor, Indonesia, Prosea Foundation: 186-190.
- Vikram, A., Jesudhasan, P.R., Jayaprakasha, G.K., Pillai, B.S. and Patil, B.S. (2010). Grapefruit bioactive limonoids modulate *E. coli* O157:H7 TTSS and biofilm. *International Journal of Food Microbiology*, 140 (2-3), 109–116.
- Wada, L. and Ou, B. (2002). Antioxidant activity and phenolic content of Oregon caneberries. *Journal of Agricultural and Food Chemistry*, 50 (12), 3495–3500.
- Wan, R., Mo, Y., Zhang, X., Chienc, S., Tolleruda, D.J. and Zhang, Q. (2008). Matrix metalloproteinase-2 and -9 are induced differently by metal nanoparticles in human monocytes: The role of oxidative stress and protein tyrosine kinase activation. *Toxicol Appl Pharmacol.*, 233 (2), 276–285.
- Wanasundara, P.K.J.P.D. and Shahidi, F. (2005). Antioxidants: Science, Technology, and Applications. *Bailey's Industrial Oil and Fat Products*. F. Shahidi. Canada, John Wiley & Sons, Inc: 431-488.
- Watson, A., Fray, T., Clarke, S., Yates, D. and Markwell, P. (2002). Reliable use of the ServoMed Evaporimeter EP-2 to assess transepidermal water loss in the canine. *J. Nutr.*, 132, 1661S–1664S.

- Weng, C. and Yen, G. (2012). Chemopreventive effects of dietary phytochemicals against cancer invasion and metastasis: Phenolic acids, monophenol, polyphenol, and their derivatives. *Cancer Treat Rev*, 38, 76–87.
- Wickett, R.R. (2001). Stretching the Skin Surface *Cosmetics and Toiletries*, 116 (3), 47-54.
- Widomska, J., Kostecka-Gugała, A., Latowski, D., Gruszecki, W.I. and Strzałka, K. (2009). Calorimetric studies of the effect of cis-carotenoids on the thermotropic phase behavior of phosphatidylcholine bilayers. *Biophysical Chemistry Volume*, 140 (1–3), 108–114.
- Williams, C.A. and Grayer, R.J. (2004). Anthocyanins and other flavonoids. *Nat. Prod. Rep.*, 21, 539-573.
- Wittern, K.P. (1991). Cosmetic preparation for skin care. *Cosmetics and toiletries development, production and use*. W. Umbach. New York, Ellis Horwood: 76.
- Yoshimura, K., Harii, K., Masuda, Y., Takahashi, M., Aoyama, T. and Iga, T. (2001). Usefulness of a narrow-band reflectance spectrophotometer in evaluating effects of depigmenting treatment. *Aesthetic Plast Surg*, 25, 129–133.
- Zhang, H.J., Zhao, W., Venkataraman, S., Robbins, M.E., Buettner, G.R., Kregel, K.C. and Oberley, L.W. (2002). Activation of matrix metalloproteinase-2 by overexpression of manganese superoxide dismutase in human breast cancer MCF-7 cells involves reactive oxygen species. *J Biol Chem*, 277 (23), 20919-20926.
- Zhang, Z., Li, D., Wang, L., Ozkan, N., Chen, X.D., Mao, Z. and Yang, H. (2007). Optimization of ethanol-water extraction of lignans from flaxseed. *Separation and Purification Technology*, 57 (1), 17-24.

Zhao, Y.L. and Nalwa, H.S. (2007). Nanotoxicology-Interactions of Nanomaterials with Biological Systems. Nanotoxicology. Los Angeles, California, American Scientific Publishers.



ลิขสิทธิ์มหาวิทยาลัยเชียงใหม่

Copyright© by Chiang Mai University
All rights reserved