

## REFERENCES

- Anderson, E. and G.L. Stebbins. 1954. Hybridization as an evolutionary stimulus. *Evolution*, 8: 378–388.
- Barbier, P. 1989a. Genetic variation and ecotypic differentiation in the wild rice species *Oryza rufipogon*. I. Population differentiation in life-history traits and isozymic loci. *Japanese Journal of Genetics*, 64: 259–271.
- Barbier, P. 1989b. Genetic variation and ecotypic differentiation in the wild rice species *Oryza rufipogon*. II. Influence of the mating system and life-history traits on the genetic structure of populations. *Japanese Journal of Genetics*, 64: 273–285.
- Barnaud, A., M. Deu, E. Garine, J. Chantereau, J. Bolteu, E.O. Koi'da, D. McKey and H.I. Joly. 2009. A weed–crop complex in sorghum: the dynamics of genetic diversity in a traditional farming system. *American Journal of Botany*, 96: 1869–1879.
- Barrett, S. C. H. 1983. Crop mimicry in weeds. *Economic Botany*, 37: 255-282.
- Bellue, M.K. 1932. Weeds of California seed rice. California. *Department of Agriculture Monthly Bull*, 21: 290-296.
- Bergelson, J. and C. B. Purrington, 1996. Surveying pattern in the cost of resistance in plants. *the American Naturalist*, 148:536-58.

- Burgos, N.R., J.K. Norsworthy, R.C. Scott and K.L. Smith. 2008. Red rice (*Oryza sativa*) status after 5 years of imidazolinone-resistant rice technology in Arkansas. *Weed Technology*, 22: 200–208.
- Campbell, L.G., A.A. Snow and C.E. Ridley. 2006. Weed evolution after crop gene introgression: greater survival and fecundity of hybrids in a new environment. *Ecology Letters*, 9: 1198–1209.
- Cao, Q., B.R. Lu, H. Xia, J. Rong, F. Sala, A. Spada, and F. Grassi. 2006. Genetic diversity and origin of weedy rice (*Oryza sativa* f. *spontanea*) population found in North-eastern China revealed by simple sequence repeat (SSR) markers. *Annals of Botany*, 98:1241-1252.
- Cavalli-Sforza, L.L. and A.W.F. Edwards. 1967. Phylogenetic analysis: models and estimation procedures. *Evolution*, 21(3):550-570.
- Chase M. W. and H. G. Hills. 1991. Silica gel: An ideal material for field preservation of leaf samples for DNA studies. *Taxon*, 40:215-220.
- Chen, L.J., D.S. Lee, Z.P. Song, H.S. Suh and B.R. Lu. 2004. Gene flow from cultivated rice (*Oryza sativa*) to its weedy and wild relatives. *Annals of Botany, (London)*. 93: 67–73.
- Chin, V.C., C.V. Hach, N.C. Thanh and N.T. Tai. 1999. Weedy rice situation in Vietnam. In FAO. Global workshop on red rice control, pp. 67-74. Report of Workshop, 30 August-3 September 1999, Varadero, Cuba. *Plant Production and Protection Division, Rome*, 155 pp.
- Chitrakon, S. 1995. Characterization, evaluation and utilization of wild rice germplasm in Thailand. Pathum Thani Rice Research Center. Thonburi, Pathum Thani, Thailand. 143 p.

- Coffey, K. 2002. Quantitative methods for the analysis of agrodiversity in Cultivating biodiversity. Edited by Brookfield, H., C. Padoch, H. Parsons and M. Stocking. *The Intermediate Technology Development Group Publishing*, pp. 78-94.
- Constantin, M.J. 1960. Characteristics of red rice in Louisiana. Louisiana State University, *Baton Rouge*, USA. (PhD dissertation)
- Cousens, R. and M. Mortimer. 1995. Dynamics of weed populations. *Cambridge University Press*. UK.
- De Datta, S.K. 1981. Principles and Practices of Rice Production. Los Baños, Philippines.
- Delouche, J.C., Burgos, N.R., Gealy, D.R., Gonzalo Aorrilla de San Martin. & Labrada, R. 2007. Weedy Rices-origin, biology, ecology and control. FAO, *Rome*, 144 p.
- De Wet, J.M.J and J.R. Harlan. 1975. Weeds and domesticates: evolution in the man-made habitat. *Economic Botany*, 29:99-107.
- Dodson, W.R. 1990. Rice weeds in Louisiana. St. Louis. *Annual Report of Agriculture Experiment Station Bulletin*, 61 (Part II).
- Dominguez, A.O. 1999. Situacion del arroz rojo en Venezuela. In FAO. Global workshop on red rice control, pp. 59-65. Report of Workshop, 30 August-3 September 1999, Varadero, Cuba. *Plant Production and Protection Division*, *Rome*, 155 pp.
- Doyle, J. J. and J. L. Doyle. 1987. A rapid DNA isolation procedure for small quantities of fresh leaf tissue. *Focus*, 12: 13-15.

- Ellstrand, N.C., H.C. Prentice and J.F. Hancock. 1999. Gene flow and introgression from domesticated plants into their wild relatives. *Annual Review of Ecology and Systematics*, 30:539-563.
- Ellstrand, N.C. and K.A. Schierenbeck. 2000. Hybridization as a stimulus for the evolution of invasiveness in plants? *Proceedings of the National Academy of Sciences, USA*. 97: 7043–7050.
- Gao, L.Z. 2004. Population structure and conservation genetics of wild rice *Oryza rufipogon* (Poaceae): a region-wide perspective from microsatellite variation. *Molecular Ecology*, 13:1009–1024.
- Ge, S., G.C.X. Oliveira, B.A. Schaal, L.Z. Gao and D.Y. Hong. 1999. RAPD variation within and between natural populations of the wild rice *Oryza rufipogon* Griff. from China and Brazil. *Heredity*, 82:638–644.
- Ghersa, C.M., M.L. Roush, S.R. Radosevich and S. Cordray. 1994. Co-evolution of agroecosystems and weed management. *BioScience*, 44: 85-94.
- Goudet J. 2001. FSTAT, a program to estimate and test gene diversities and fixation indices (version 2.9.3). <http://www.unil.ch/izea/softwares/fstat.html>. (1 December 2007).
- Guglielmini A.C., C.M. Ghersa and E.H. Satorre. 2007. Co-evolution of domesticated crops and associated weeds. *Ecología Austral*, 17:167-178.
- Harlan, J.R. 1982. Relationships between weeds and crops. In: W. Holzner and M. Numata (eds). *Biology and Ecology of weeds*, pp. 91-96. Dr. W. Junk Publishers. The Hague. Netherlands.
- Harlan, J.R. 1992. Crop and Man. 2<sup>nd</sup> edition. Madison, Wisconsin, USA.

- Harlan, D.R. and J.M.J. de Wet. 1971. Towards a rational classification of cultivated plants. *Taxon*, 20:509-517.
- Huey, B. and F. Baldwin. 1978. Red rice control. Pages 19-25. in E. F. Eastin, ed. Red rice: research and control. *Texas Agricultural Experiment Station Bulletin*, No. 1270.
- IRR-IBPGR. 1980. Descriptions for rice *Oryza sativa* L. IRRI, P.O. Box 993, Manila, Philippines, 21 p.
- Jamjod, S., C. Maneechote, T. Anakad, S. Niruntrayagul and B. Rekasem. 2003. Wild rice in Thailand: snapshots from 2002. Annual review 3, CMUPNlab working papers V, Chiang Mai University: 1-15.
- Jarvis D.I. and T. Hodgkin. 1999. Wild relatives and crop cultivars: detecting natural introgressions and farmer selection of new genetic combinations in agroecosystems. *Molecular Ecology*, 8:S159-S173.
- Jindaluang, S. 2009. Inheritance of Seedling Length in Weedy Rice. Master Thesis, Graduate School, Chiang Mai University, Chiang Mai.
- Kono, Y. and S. Konchan. 1996. Spread of Direct Seeded Lowland Rice in Northeast Thailand: Farmers' Adaptation to Economic Growth. *Southeast Asian Studies*, 33[4]:5-28.
- Kuroda, Y. 2004. Conservation biological studies on Asian wild rice (*O. rufipogon* and *O. nivara*) in the Vientiane Plain of Laos. Kyoto University. Japan.
- Langevin, S.A., K.Clay and J.B. Grace. 1990. The incidence and effect of hybridization between cultivated rice and its related weed red rice (*Oryza sativa* L.). *Evolution*, 44(4): 1000-1008.

- Liu, K. and S.V. Muse. 2005. POWERMARKER: integrated analysis environment for genetic marker data. *Bioinformatics*, 21:2128-2129.
- Maneechote, C. and S. Jamjod. 2005. Situation of weedy rice infestation in Thailand. Weedy rice symposium, Plant Protection Research and Development Office Department of Agricultural, Rama Gardens Hotel Bangkok, Thailand 21 October 2005. p 1-13.
- Maneechote, C., S. Jamjod and B. Rerkasem. 2004. Invasion of weedy rice in rice fields in Thailand: problems and management. *International Rice Research Notes*, 29(2): 20-22.
- Manly, K. F., R. H. Jr. Cudmore and J. M. Meer. 2001. *Mammalian Genome*, 12:930–932.
- Messeguer, J., C. Fogher, E. Guiderdoni, V. Marfa, M.M. Catala, G. Baldi and E. Mele. 2001. Field assessments of gene flow from transgenic to cultivated rice (*Oryza sativa L.*) using a herbicide resistance gene as tracer marker. *Theoretical and Applied Genetics*, 103: 1151-1159.
- Messeguer, J., V. Marfa, M.M. Catala, E. Guiderdoni and E. Mele. 2004 A field study of pollen-mediated gene flow from Mediterranean GM rice to conventional rice and red rice weed. *Molecular Breeding*, 13: 103-112.
- Michelmore, R., W. I. Paran and R. V. Kesseli. 1991. Identification of markers linked to disease-resistance genes by bulked segregant analysis: A rapid method to detect markers in specific genomic regions by using segregating populations. *Proceeding of the Natural Academy of Sciences, USA*, 88: 9828-9832.
- Morishima, H. 1986. Rice genetic. International Rice Research Institute. Manila Philippines.


- Morishima, H. and P. Barbier. 1990. Mating system and genetic structure of natural populations in wild rice *Oryza rufipogon* Griff. *Plant Species Biology*, 5:31-39.
- Morishima, H. and H.I. Oka. 1970. A survey of genetic variations in the populations of wild *Oryza* species and their cultivated relatives. *Japanese Journal of Genetics*, 45:371-385.
- Nagai, I. 1958. Japonica rice. Its breeding and culture. Yokendo, Tokyo, 843 pp.
- Nei, M. 1973. Analysis of gene diversity in subdivided populations. *Proceeding of the National Academy of Sciences, USA*, 70 (12):3321-3323.
- Niruntrayakul, S. 2008. Gene flow between cultivated and wild rice. Ph.D. Thesis, Graduate School, Chiang Mai University, Chiang Mai.
- Noldin, J.A. 2000. Red rice status and management in the Americas. In: Baki BB, Chin DV, Mortimer M, editors. Proceeding of wild and weedy rice in rice ecosystems in Asia: a review. Los Baños, Manila, Philippines: *International Rice Research Institute*, p 21-24.
- Noldin, J.A., J.M. Chandler and G.N. McCauley. 1999. Red rice (*Oryza sativa*) biology. I. Characterization of red rice ecotypes. *Weed Technology*, 13: 12-18.
- Oka, H. 1988. Origin of cultivated rice. Japan Scientific Societies Press, Tokyo.
- Agricultural Futures Trading Commission. 2007. Available from [http://www.aftc.or.th/itc/products\\_analyze\\_price\\_04.php?id=56&fgrp\\_id=5&fmnu\\_id=18](http://www.aftc.or.th/itc/products_analyze_price_04.php?id=56&fgrp_id=5&fmnu_id=18) (26 May 2008).
- Palumbi, S.R. 2001. Human as the world's greatest evolutionary force. *Science*, 293: 1786-1790.

- Panaud, O., Chen, X. and S. McCouch. 1996. Development of microsatellite markers and characterization of simple sequence length polymorphism (SSLP) in rice (*Oryza sativa L.*). *Molecular and General Genetics*, 252:597-607.
- Papa, R and P. Gepts. 2004. Gene flow between crops and their wild progenitors. *Encyclopedia of Plant and Crop Science*, DOI:10.1081/E-EPCS 120017095.
- Peakall, R. and P.E. Smouse. 2006. GenAIEx 6: genetic analysis in Excel. Population genetic software for teaching and research. *Molecular Ecology Notes*, 6(1):288-295.
- Postlethwait, J. 2009. Modern Biology. Holt, Rinehart and Winston, p. 317.
- Pritchard, J. K. and W. Wen. 2004. Documentation for Structure software: version 2. See <http://pritch.bsd.uchicago.edu/software.html>. (1 December 2006).
- Pritchard J.K., Stephens M., and P. Donnelly. 2000. Inference of population structure using multilocus genotype data. *Genetics*, 155: 945–959.
- Punyalue, A. 2006. Characterization of common wild rice populations from main rice growing region of Thailand. Master Thesis, Graduate School, Chiang Mai University
- Pusadee, T. 2009. Evolutionary dynamics of *Oryza sativa* in Thailand. Ph.D. Thesis, Graduate School, Chiang Mai University, Chiang Mai.
- Radosevich, S., J. Holt and C. Ghersa. 1997. Weed ecology: *Implications for management*. 2<sup>nd</sup> editor New York, USA, John Wiley and Sons, Inc.
- Robert, E.H., R.Q. Craufurd and F. Le Cochet. 1961. Estimation of percentage of natural cross-pollination: *experiment on rice*. *Nature*, 190: 1084-1085.
- Rohlf F.J. (1998). NTSYSpc: Numerical Taxonomy and Multivariate Analysis System. Version 2.02. Exeter Software, Setauket, New York.

- Rong, J., B.R. Lu, Z. Song, J. Su, A.A. Snow, X. Zhang, S. Sun, R. Chen and F. Wang. 2006. Dramatic reduction of crop-to-crop gene flow within a short distance from transgenic rice fields. *New Phytologist*, doi: 10.1111/j.1469-8137.2006.01906.x.
- Roumet M., M.F. Ostrowski, J. David, C. Tollon and M.H. Muller. 2012. Estimation of mating system parameters in an evolving gynodioecious population of cultivated sunflower (*Helianthus annuus* L.). *Heredity*, 108: 366–374.
- Semon, M., R. Nielsen, M.P. Jones and S.R. McCouch. 2005. The population structure of African cultivated rice *Oryza glaberrima* (Steu.): evidence for elevated levels of linkage disequilibrium caused by admixture with *O. sativa* and ecological adaptation. *Genetics*, 169:1639-1647.
- Shannon, C. E. and W. Weaver, 1949. The Mathematical Theory of Communication. Urbana, IL: University of Illinois Press.
- Sirabanchongkran, A., N. Yimyam, W. Boonma, K. Rerkasem, K. Coffey, M. Pinedo-Vasquez, and C. Padoch, 2004. Varietal turnover and seed exchange: implications for conservation of rice genetic diversity on-farm. *International Rice Research Note*, 29(2):12-14.
- Smith, R.J., Jr. 1968. Weed competition in rice. *Weed Science*, 16: 252-255.
- Song, Z.P., X. Xu, B. Wang, J.K. Chen and B.R. Lu. 2003. Genetic diversity in the northernmost *Oryza rufipogon* populations estimated by SSR markers. *Theoretical and Applied Genetics*, 107: 1492–1499.
- Sonnier, E. 1978. Cultural control of red rice. Pages 10-15. in E. F. Eastin, ed. Red rice: research and control. *Texas Agricultural Experiment Station Bulletin*, No. 1270.

- Spencer, C. and H. Barrett. 1983. Crop Mimicry in Weeds. *Economic Botany*, 37(3), 1983, pp. 255-282.
- Sun, C.Q., X.K. Wang, Z.C. Li and A. Yoshimura. 2001. Comparison of the genetic diversity of common wild rice (*Oryza rufipogon* Griff.) and cultivated rice (*O. sativa* L.) using RFLP markers. *Theor. Appl. Genet.*, 102: 157-162.
- Tamura, K., Dudley, J., Nei, M. and S. Kumar. 2007. MEGA4: molecular evolution genetics analysis (MEGA) software version 4.0. *Mol. Biol. Evol.*, 24(8):1596-1599.
- Vaughan, D.A. 1994. The wild relatives of rice: a genetic resources guide book. Los Baños, Philipines: *International Rice Research Institute*, 137 p.
- Vaughan, L.K., B.V. Ottis, A.M. Prazak-Havey, C. Sneller, J.M. Chandler and W.D. Park, 2001. Is all red rice found in commercial rice really *Oryza sativa*? *Weed Science*, 49:468-476.
- Vaughan, D.A., P.L. Sanchez, J. Ushiki, A. Kaga and N. Tomooka. 2005. Chapter 16 Asian rice and weedy rice evolutionary perspectives. pp 257-277. In Gressel J. (Ed.) Crop ferality and volunteerism. *CRC press, Taylor&Francis group*, Boca Raton, FL, USA.
- Vaughan D.A. and M. Morishima. 2003. Biosystematics of genus *Oryza*. In Smith, W.C. and R.H. Dilday, eds. Rice: origin, history, *technology and production*, pp. 27-65. Hoboken, USA, *John Wiley and Sons, Inc.*
- Wague, K. 1992. Comparison of seedling vigor and competitiveness in selected red rice and cultivated rice. Mississippi State University, USA. (MSc thesis).
- Weir, B.S. and C.C. Cockerham, 1984. Estimating F-statistics for the analysis of population structure. *Evolution*, 38: 1358-1370.

- Wiley R.H. 1981. Social structure and individual ontogenies: problems of description, mechanism, and evolution. P. 105-133. In Bateson P. P. G. and P. H. Klopfer (Eds), *Perspectives in ethology*, vol. 4. Plenum Press, New York.
- Wongtamee, A. 2008. Evaluation of exotic genes contamination from crop rice (*Oryza sativa* L.) in common wild rice (*Oryza rufipogon* Griff.) populations. Master Thesis, Graduate School, Chiang Mai University
- Yeh, F.C., Yang, R.C., and T.B.J. Boyle. 1999. POPGENE, version 1.32 Microsoft window-based freeware for population genetic analysis. (University of Alberta, Edmonton, Alberta Canada)
- Zhang, N., S. Linscombe and J. Oard. 2003. Out-crossing frequency and genetic analysis of hybrids between transgenic glufosinate herbicide-resistant rice and the weedy rice. *Euphytica*, 130: 35-45.



ลิขสิทธิ์มหาวิทยาลัยเชียงใหม่  
Copyright© by Chiang Mai University  
All rights reserved