

TABLE OF CONTENTS

	Page
Acknowledgements	iii
Abstract (English)	v
Abstract (Thai)	xi
List of Tables	xviii
List of Figures	xxv
Introduction	1
Chapter 1 Literature review	6
1.1 <i>Oryza</i> Genus	6
1.2 <i>Oryza sativa</i> primary gene pool	7
1.2.1 Common wild rice	8
<i>Genetic diversity of common wild rice</i>	8
1.2.2 Cultivated rice	9
1.2.3 Weedy rice	10
<u>Specific characters of weedy rice</u>	11
<i>Adaptation and crop mimicry</i>	11
<i>Shattering and dormancy</i>	12
<i>Competition</i>	13

	<i>Dispersion</i>	14
	<u>Gene flow or migration by hybridization in plants</u>	15
	<i>Crop-to-wild gene flow</i>	15
	<i>Crop-to-weed gene flow</i>	15
	<i>Crop-to-crop gene flow</i>	16
	<u>Selection: Agricultural practices as selection forces</u>	17
Chapter 2	Genetic structure of weedy rice populations from three rice production areas of Thailand	18
2.1	Introduction	18
2.2	Materials and methods	20
2.3	Results	33
	<i>Characterization in rice fields</i>	33
	<i>Morphological and physiological characterization</i>	37
	<i>Population genetic structure and spatial dynamics</i>	60
	<u><i>Populations from wet season of 2005</i></u>	60
	<u><i>Populations from wet season of 2008</i></u>	72
	<u><i>Populations from wet season of 2009</i></u>	84
	<u><i>Populations from dry season of 2009/10</i></u>	97
	<i>Population genetic structure and temporal dynamics</i>	109
2.4	Discussion	127

Chapter 3	Genetic structure and dynamics of sympatric wild-weed-crop rice populations in Thailand	132
	3.1 Introduction	132
	3.2 Materials and methods	135
	3.3 Results	143
	<i>Morphological and physiological characterization</i>	143
	<i>Population genetic structure and spatial dynamics</i>	154
	<u><i>Alleles and genotypes specific to crop and wild types</i></u>	154
	<u><i>Populations from wet season of 2005</i></u>	158
	<u><i>Populations from wet season of 2008</i></u>	168
	<u><i>Populations from wet season of 2009</i></u>	178
	<i>Population genetic structure and temporal dynamics</i>	188
	3.4 Discussion	203
Chapter 4	General Discussion	209
References		213
Appendix		224
Curriculum vitae		231

LIST OF TABLES

Table	Page
2.1.1 Location and description of weedy rice populations	21
2.1.2 Common wild rice accession, location and habitat in this study	24
2.1.3 Name, type, pedigree description and year released of cultivated rice in the present study	26
2.1.4 Microsatellite markers in the present study	28
2.2.1 Frequency (%) of type, plant height and seed characters of weedy rice populations of Thailand	35
2.2.2 Morphological characteristics and Shannon-Weaver Index (H') of weedy rice populations of Thailand	39
2.2.3 Physiological characteristics of weedy rice populations	48
2.3.1 Allele diversity of 8 weedy rice populations of Thailand in wet season of 2005 using 12 SSR markers	61
2.3.2 Genetic parameters of 8 weedy rice populations of Thailand in wet season of 2005 using 12 SSR markers	64
2.3.3 Summary of genetic parameters of weedy rice populations of Thailand in wet season of 2005 based on 12 SSR loci	64

2.3.4	Analysis of molecular variance (AMOVA) among populations and three rice production areas for 151 individuals of 8 weedy rice populations in wet season of 2005 based on 12 SSR loci	65
2.3.5	Pairwise of genetic differentiation (F_{ST}) within and between three rice production areas of 8 weedy rice populations in wet season of 2005 based on 12 SSR loci	65
2.3.6	Allele diversity of 20 weedy rice populations of Thailand in wet season of 2008 using 12 SSR markers	73
2.3.7	Genetic parameters of 20 weedy rice populations of Thailand in wet season of 2008 using 12 SSR markers	76
2.3.8	Summary of genetic parameters of weedy rice populations of Thailand in wet season of 2008 based on 12 SSR loci	77
2.3.9	Analysis of molecular variance (AMOVA) among populations and three rice production areas for 386 individuals of 20 weedy rice populations in wet season 2008 based on 12 SSR loci	77
2.3.10	Pairwise of genetic differentiation (F_{ST}) within and between three rice production areas of 20 weedy rice populations in wet season of 2008 based on 12 SSR loci	77
2.3.11	Allele diversity of 18 weedy rice populations of Thailand in wet season of 2009 using 12 SSR markers	85
2.3.12	Genetic parameters of 18 weedy rice populations of Thailand in wet season of 2009 using 12 SSR markers	88

2.3.13	Summary of genetic parameters of weedy rice populations of Thailand in wet season of 2009 based on 12 SSR loci	89
2.3.14	Analysis of molecular variance (AMOVA) among populations and three rice production areas for 180 individuals of 18 weedy rice populations in wet season of 2009 based on 12 SSR loci	89
2.3.15	Pairwise of genetic differentiation (F_{ST}) within and between three rice production areas of 18 weedy rice populations in wet season of 2009 based on 12 SSR loci	90
2.3.16	Allele diversity of 17 weedy rice populations of Thailand in dry season of 2009/10 using 12 SSR markers	98
2.3.17	Genetic parameters of 17 weedy rice populations of Thailand in dry season of 2009/10 using 12 SSR markers	101
2.3.18	Summary of genetic parameters of weedy rice populations of Thailand in dry season of 2009/10 based on 12 SSR loci	102
2.3.19	Analysis of molecular variance (AMOVA) among populations and three rice production areas for 170 individuals of 18 weedy rice populations in dry season of 2009/10 based on 12 SSR loci	102
2.3.20	Pairwise of genetic differentiation (F_{ST}) within and between three rice production areas of 17 weedy rice populations in dry season of 2009/10 based on 12 SSR loci	103

- 2.3.21 Allele diversity of 63 weedy rice populations collected from three rice production areas; Lower North, Northeast and Central Plain of Thailand in four growing seasons in wet season 2005, 2008, 2009 and dry season of 2009/10 using 12 SSR markers 111
- 2.3.22 Genetic parameters of 63 weedy rice populations collected from three rice production areas; Lower North, Northeast and Central Plain of Thailand in four growing seasons in wet season of 2005, 2008, 2009 and dry season of 2009/10 using 12 SSR markers 115
- 2.3.23 Analysis of molecular variance (AMOVA) among populations of each for 887 individuals of 63 weedy rice populations collected from three rice production areas; Lower North, Northeast and Central Plain of Thailand in four growing seasons in wet season of 2005, 2008, 2009 and dry season of 2009/10 using 12 SSR markers 115
- 2.3.24 Pairwise of genetic differentiation (F_{ST}) within and between four growing seasons in wet season of 2005, 2008, 2009 and dry season of 2009/10 of 63 weedy rice populations collected from three rice production areas; Lower North, Northeast and Central Plain of Thailand using 12 SSR Markers 116
- 3.1.1 Location and description of weedy rice, companion crop rice and native common wild rice populations with occur in sympatric rice field 136

3.1.2	Popular crop rice varieties, companion crop rice, weedy rice and native common wild rice populations in each rice growing region and year	142
3.2.1	Frequency (%) of six morphological characteristics of sympatric wild-weed-crop rice populations in Thailand	144
3.2.2	Physiological characteristics of sympatric wild-weed-crop rice populations in Thailand	149
3.3.1	Frequency of alleles specific to crop types and wild types in sympatric wild-weed-crop rice populations based on 12 microsatellite markers	156
3.3.2	Genotypic frequency of crop types, wild types and hybrid types in sympatric wild-weed-crop rice populations based on 12 microsatellite markers	157
3.3.3	Genetic parameters of sympatric wild-weed-crop rice populations in wet season of 2005 based on 12 microsatellite markers	160
3.3.4	Analysis of molecular variance (AMOVA) among sympatric wild-weed-crop rice populations in wet season of 2005 based on 12 microsatellite markers	161
3.3.5	Pairwise of genetic differentiation (F_{ST}) within and between sympatric wild-weed-crop rice populations in wet season of 2005 based on 12 microsatellite markers	161

3.3.6	Genetic parameters of sympatric wild-weed-crop rice populations in wet season of 2008 based on 12 microsatellite markers	170
3.3.7	Analysis of molecular variance (AMOVA) among sympatric wild-weed-crop rice populations in wet season of 2008 based on 12 microsatellite markers	171
3.3.8	Pairwise of genetic differentiation (F_{ST}) within and between sympatric wild-weed-crop rice populations in wet season of 2008 based on 12 microsatellite markers	171
3.3.9	Genetic parameters of sympatric wild-weed-crop rice populations in wet season of 2009 based on 12 microsatellite markers	180
3.3.10	Analysis of molecular variance (AMOVA) among sympatric wild-weed-crop rice populations in wet season of 2009 based on 12 microsatellite markers	181
3.3.11	Pairwise of genetic differentiation (F_{ST}) within and between sympatric wild-weed-crop rice populations in wet season of 2009 based on 12 microsatellite markers	181
3.3.12	Genetic parameters of 18 companion crop rice, 18 weedy rice and 18 native common wild rice populations collected from three growing seasons in 2005, 2008 and 2009, using 12 SSR markers	191
3.3.13	Analysis of molecular variance (AMOVA) among populations of each for 540 individuals of 18 companion crop rice, 18 weedy rice and 18 native common wild rice populations collected from three growing seasons in 2005, 2008 and 2009, using 12 SSR markers	192

- 3.3.14 Pairwise of genetic differentiation (F_{ST}) within and between 18 companion crop rice, 18 weedy rice and 18 native common wild rice populations collected from three growing seasons in 2005, 2008 and 2009, using 12 SSR markers

192



ลิขสิทธิ์มหาวิทยาลัยเชียงใหม่
Copyright© by Chiang Mai University
All rights reserved

LIST OF FIGURES

Figure	Page
2.1.1 Location of weedy rice sampling in Thailand of the present study	23
2.1.2 Location of common wild rice sampling in Thailand of the present study	25
2.2.1 Three types of weedy rice; Khao-harng, Khao Deed and Khao Lai	36
2.2.2 Morphological characteristics of weedy rice populations	43
2.2.3 Cluster analysis using NJ method on the basis of C.S. Chord (1967) genetic distance among pure line rice varieties, weedy rice and common wild rice populations in Thailand constructed based on 14 morphological traits	44
2.2.4 Distribution of days to flowering of weedy rice populations in Thailand	50
2.2.5 Distribution of tillers plant ⁻¹ of weedy rice populations in Thailand	51
2.2.6 Distribution of panicle plant ⁻¹ of weedy rice populations in Thailand	52
2.2.7 Distribution of culm length (cm) of weedy rice populations in Thailand	53
2.2.8 Distribution of panicle length (cm) of weedy rice populations in Thailand	54

2.2.9	Distribution of number of primary branches panicle ⁻¹ of weedy rice populations in Thailand	55
2.2.10	Distribution of number of spikelets panicle ⁻¹ of weedy rice populations in Thailand	56
2.2.11	Distribution of number of seeds panicle ⁻¹ of weedy rice populations in Thailand	57
2.2.12	Distribution of seed fertility (%) of weedy rice populations in Thailand	58
2.2.13	Distribution of seed shattering (%) of weedy rice populations in Thailand	59
2.3.1	Population assignment of 8 weedy rice populations of Thailand in wet season of 2005	68
2.3.2	Genetic proportion of 8 weedy rice populations of Thailand in wet season of 2005	69
2.3.3	Distribution of 151 individuals of 8 weedy rice populations of Thailand in wet season of 2005 formed by the principal coordinate analysis (PCA) on the basis of 12 microsatellite markers	70
2.3.4	Cluster analysis using NJ method based on C.S. Chord (1967) genetic distance among 8 weedy rice populations of Thailand in wet season of 2005	71
2.3.5	Population assignment of 20 weedy rice populations of Thailand in wet season of 2008	80

2.3.8	Genetic proportion of 20 weedy rice populations of Thailand in wet season of 2008	81
2.3.7	Distribution of 386 individuals of 20 weedy rice populations of Thailand in wet season of 2008 formed by the principal coordinate analysis (PCA) on the basis of 12 microsatellite markers	82
2.3.8	Cluster analysis using NJ method based on C.S. Chord (1967) genetic distance among 20 weedy rice populations of Thailand in wet season of 2008	83
2.3.9	Population assignment of 18 weedy rice populations of Thailand in wet season of 2009	93
2.3.10	Genetic proportion of 18 weedy rice populations of Thailand in wet of 2009	94
2.3.11	Distribution of 180 individuals of 18 weedy rice populations of Thailand in wet season of 2009 formed by the principal coordinate analysis (PCA) on the basis of 12 microsatellite markers	95
2.3.12	Cluster analysis using NJ method based on C.S. Chord (1967) genetic distance among 18 weedy rice populations of Thailand in wet season of 2009	96
2.3.13	Population assignment of 17 weedy rice populations of Thailand in dry season of 2009/10	105
2.3.14	Genetic proportion of 17 weedy rice populations of Thailand in dry season of 2009/10	106

2.3.15	Distribution of 170 individuals of 17 weedy rice populations of Thailand in dry season of 2009/10 formed by the principal coordinate analysis (PCA) on the basis of 12 microsatellite markers	107
2.3.16	Cluster analysis using NJ method based on C.S. Chord (1967) genetic distance among 17 weedy rice populations of Thailand in dry season of 2009/10	108
2.3.17	Observed heterozygosity (H_0), Genetic differentiation (F_{ST}), Inbreeding coefficient (F_{IS}) and Outcrossing rate (t) of 63 weedy rice populations of Thailand in four growing seasons using 12 SSR markers	114
2.3.18	Population assignment using STRUCTURE analysis of 63 weedy rice populations compared with 7 pure line cultivated rice varieties and 7 common wild rice populations	119
2.3.19	Genetic proportion of 63 weedy populations compared with 7 pure line cultivated rice varieties and 7 common wild rice populations	120
2.3.20	Principle component analysis (PCA) of 63 weedy populations compared with 7 pure line cultivated rice varieties and 7 common wild rice populations	122
2.3.21	Cluster analysis using NJ method based on C.S. Chord (1967) genetic distance among 63 weedy rice populations compared with 7 pure line cultivated rice varieties and 7 common wild rice populations	124

3.1.1	Location of weedy rice, companion crop rice and native common wild rice populations with occur in sympatric rice field	138
3.2.1	Cluster analysis using NJ method on the basis of C.S. Chord (1967) genetic distance among sympatric wild-weed-crop rice populations in Thailand	146
3.2.2	Distribution of days to flowering of sympatric wild-weed-crop rice populations in Thailand	150
3.2.3	Distribution of tillers plant ⁻¹ of sympatric wild-weed-crop rice populations in Thailand	151
3.2.4	Distribution of culm length (cm) of sympatric wild-weed-crop rice populations in Thailand	152
3.2.5	Distribution of seed shattering (%) of sympatric wild-weed-crop rice populations in Thailand	153
3.3.1	Population assignment of sympatric wild-weed-crop rice populations in wet 2005	164
3.3.2	Genetic proportion of sympatric wild-weed-crop rice populations in wet season 2005	165
3.3.3	Distribution of 180 individuals of sympatric wild-weed-crop rice populations in wet season of 2005 formed by the principal coordinate analysis (PCA) on the basis of 12 microsatellite markers	166
3.3.4	Cluster analysis using NJ method based on C.S. Chord (1967) genetic distance among sympatric wild-weed-crop rice populations in wet season of 2005	167

3.3.5	Population assignment of sympatric wild-weed-crop rice populations in wet season of 2008	174
3.3.6	Genetic proportion of sympatric wild-weed-crop rice populations in wet season of 2008	175
3.3.7	Distribution of 180 individuals of sympatric wild-weed-crop rice populations in wet season of 2008 formed by the principal coordinate analysis (PCA) on the basis of 12 microsatellite markers	176
3.3.8	Cluster analysis using NJ method based on C.S. Chord (1967) genetic distance among sympatric wild-weed-crop rice populations in wet season of 2008	177
3.3.9	Population assignment of sympatric wild-weed-crop rice populations in wet season of 2009	184
3.3.10	Genetic proportion of sympatric wild-weed-crop rice populations in wet season of 2009	185
3.3.11	Distribution of 180 individuals of sympatric wild-weed-crop rice populations in wet season of 2009 formed by the principal coordinate analysis (PCA) on the basis of 12 microsatellite markers	186
3.3.12	Cluster analysis using NJ method based on C.S. Chord (1967) genetic distance among sympatric wild-weed-crop rice populations in wet season of 2009	187

- 3.3.13 Observed heterozygosity (H_0), Genetic differentiation (F_{ST}), Inbreeding coefficient (F_{IS}) and Outcrossing rate (t) among companion crop rice, weedy rice and native common wild rice populations of Thailand collected from three growing seasons in 2005, 2008 and 2009, using 12 SSR markers 190
- 3.3.14 Population assignment using STRUCTURE analysis of 18 companion crop rice, 18 weedy rice and 18 native common wild rice populations in three growing seasons in 2005, 2008 and 2009 compared with 7 pure line cultivated rice varieties 196
- 3.3.15 Genetic proportion of 18 companion crop rice, 18 weedy rice and 18 native common wild rice populations in three growing seasons in 2005, 2008 and 2009 197
- 3.3.16 Principle component analysis (PCA) of 18 companion crop rice, 18 weedy rice and 18 native common wild rice populations in different three growing seasons in 2005, 2008 and 2009 compared with 7 pure line cultivated rice varieties 199
- 3.3.17 Cluster analysis using NJ method based on C.S. Chord (1967) genetic distance among 18 companion crop rice, 18 weedy rice and 18 native common wild rice in different three growing seasons in 2005, 2008 and 2009 compared with 7 pure line cultivated rice varieties populations 201