

REFERENCES

- Abadulla, E., Tzanov, T., Costa, S., Robra, K.H. and Cavaco-Paulo, A. (2000). Decolorization and detoxification of textile dyes with a laccase from *Trametes hirsute*. *Appl. Environ. Microbiol.*, 66: 3357-3362.
- Agbor, V.B., Cicek, N., Sparling, R., Berlin, A. and Levin, D.B. (2011). Biomass pretreatment: Fundamentals toward application. *Biotechnol. Adv.*, 29: 675-685.
- Aguilar, C.N., Augur, C. and Favela-Torres, E. (2001). Induction and repression patterns of fungal tannase in solid-state and submerged cultures. *Process Biochem.*, 36: 565-570.
- Aguilar, C.N., Favela-Torres, E., Viniegra-González, G. and Augur, C. (2002). Culture conditions dictate protease and tannase production in submerged and solid-state cultures by *Aspergillus niger* Aa-20. *Appl. Biochem. Biotechnol.*, 102-103: 407-414.
- Aguilar, C.N., Gutiérrez-Sánchez, G., Rado-Barragán, P.A., Rodríguez-Herrera, R., Martínez-Hernandez, J.L. and Contreras-Esquivel, J.C. (2008). Perspectives of solid state fermentation for production of food enzymes. *Am. J. Biochem. Biotechnol.*, 4: 354-366.
- Akay, K.U. (2007). A note on model selection in mixture experiments. *J. Math. Stat.*, 3: 93-99.

- Alessandro, B., Paola, G., Paolo, A., Giovanna, D.P., Ummacolata, M., Maria, T.V., Luigi, M., Lucia, A., Damiano, G.M. and Maria, M. (2008). Laccase treatment Impairs bisphenol A- induced cancer cell proliferation affecting estrogen receptor α -dependent rapid signals. *Life*, 60: 843-852.
- Alvira, P., Tomás-Pejó, E., Ballesteros, M. and Negro, M.J. (2010). Pretreatment technologies for an efficient bioethanol production process based on enzymatic hydrolysis: A review. *Bioresour. Technol.*, 101: 4851-4861.
- Ander, P. and Marzullo, L. (1997). Sugar oxidoreductases and veratryl alcohol oxidase as related to lignin degradation. *J. Biotechnol.*, 53: 115-131.
- Ariffin, H., Abdullah, N., Umi Kalsom, M.S. and Hassan, M.A. (2006). Production and characterization of cellulase by *Bacillus pumilus* EB3. *Int. J. Eng. Technol.*, 3: 47-53.
- Asit, D., Alan, B., and Kent, K. (1991). Identification of a specific manganese peroxidase among ligninolytic enzymes secreted by *Phanerochaete chrysosporium* during wood decay. *Appl. Environ. Microbiol.*, 57: 1453-1460.
- Atef, J., Francisco, G., Michel, J.P., Angel, T.M. and Maria, J.M. (2005). Role of *Pycnoporus coccineus* laccase in the degradation of aromatic compounds in olive oil mill wastewater. *Enzyme Microb. Technol.*, 36, 478-486.
- Avella, M., Casale, L., Dellerba, R., Focher, B., Martuscelli, E. and Marzetti, A. (1998). Broom fibres as reinforcing materials for polypropylene-based composites. *J. Appl. Polym. Sci.*, 68: 1077-1089.
- Aziz, N.H. and Mohsen, G.I. (2002). Bioconversion of acid- and gamma-ray-treated sweet potato residue to microbial protein by mixed cultures. *J. Ind. Microbiol. Biotechnol.*, 29: 264-267.

- Bajpai, P. (1999). Application of enzymes in the pulp and paper industry. *Biotechnol. Progr.*, 15: 147-157.
- Bastawde, K.B. (1992). Xylan structure, microbial xylanases, and their mode of action. *World J. Microbiol. Biotechnol.*, 8: 353-368.
- Bekalo, S.A. and Reinhardt, H.-W. (2010). Fiber of coffee husk and hull for the production of particleboard. *Mater. Struct.*, 43: 1049-1060.
- Betancur, G.J.V. and Pereira N.Jr. (2010). Sugar cane bagasse as feedstock for second generation ethanol production. Part I: diluted acid pretreatment optimization. *Electronic J. Biotechnol.*, 13: 1-9.
- Bigelow, M. and Wyman, C.E. (2002). Cellulase production on bagasse pretreated with hot water. *Appl. Biochem. Biotechnol.*, 98-100: 921-934.
- Blanchette, R.A. (1984). Manganese accumulation in wood decayed by white-rot fungi. *Phytopathol.*, 74: 153-160.
- Bradford, M.M. (1976). A rapid and sensitive method for the quantitation of microgram quantities of protein utilizing the principle of protein-dye binding. *Anal. Biochem.*, 72: 248-254.
- Brand, D., Pandey, A., Roussos, S. and Soccol, C.R. (2000). Biological detoxification of coffee husk by filamentous fungi using a solid state fermentation system. *Enzyme Microb. Technol.*, 27: 127-133.
- Brienzo, M., Monte, J.R. and Milagres, A.M.F. (2012). Induction of cellulase and hemicellulase activities of *Thermoascus aurantiacus* by xylan hydrolyzed products. *World J. Microbiol. Biotechnol.*, 28: 113–119.
- Brown, A. (1985). Review of lignin in biomass. *J. Appl. Biochem.*, 7: 371-387.

Browning, B.L. (1967). Method of Wood Chemistry. Vol. II. Wiley and Sons Interscience. New York. pp. 397.

Buddolla, V., Subhosh, M.C., Pallavi, H., and Rajasekher, B.R. (2008). Screening and assessment of laccase producing fungi isolated from different environmental samples. *Afr. J. Biotechnol.*, 7: 1129-1133.

Cambria, M.T., Cambria, A., Ragusa, S. and Rizzarelli, E. (2000). Production, purification, and properties of an extracellular laccase from *Rigidoporus lignosus*. *Protein Expression Purif.*, 18: 141-147.

Cameselle, C., Pazos, M., Lorenzo, M. and Sanromán, M.A. (2003). Enhanced decolourisation ability of laccase towards various synthetic dyes by an electrocatalysis technology. *Biotechnol. Lett.*, 25: 603-606.

Cara, C. Ruiz, E., Ballesteros, M., Manzanares, P., Negro, M.J. and Castro, E. (2008). Production of fuel ethanol from steam-explosion pretreated olive tree pruning. *Fuel*, 87: 692-700.

Chahal, D.S. (1991). Pretreatment of lignocelluloses. In: Chahal D.S. (ed) Food feed and fuel from biomass. Oxford and IBH, New Delhi. pp. 59-83.

Chai, W., Handa, Y., Suzuki, M., Saito, M., Kato, N. and Horiuchi, C.A. (2005). Biodegradation of bisphenol A by fungi. *Appl. Biochem. Biotechnol.*, 120: 175-182.

Chandra, R.P., Bura R., Mabey, W.E., Berlin, A., Pan, X. and Saddler, J.N. (2007). Substrate Pretreatment: The key to effective enzymatic hydrolysis of lignocellulosics. *Adv. Biochem. Eng/Biotechnol.*, 108: 67-93.

- Chawachart, N., Khanongnuch, C., Watanabe, T. and Lumyong, S. (2004). Rice bran as an efficient substrate for laccase production from thermotolerant basidiomycete *Coriolus versicolor* strain RC3. *Fungal Divers.*, 15: 23-32.
- Chivukula, M. and Renganathan, V. (1995). Phenolic azo dye oxidation by laccase from *Pyricularia oryzae*. *Appl. Environ. Microbiol.*, 61: 4374-4377.
- Christiane, G., Sabine, G., Clemens, K.P., Josef, S. and Dietmar, H. (2002). Characterization of the major laccase isoenzyme from *Trametes pubescens* and regulation of its synthesis by metal ions. *Microbiol.*, 148: 2159-2169.
- Claus, H., Faber, G. and König, H. (2002). Redox-mediated decolorization of synthetic dyes by fungal laccases. *Appl. Microbiol. Biotechnol.*, 59: 672-678.
- Couto, S.R., Rodríguez, A., Paterson, R.R.M., Lima, N. and Teixeira, J.A. (2006). Laccase activity from the fungus *Trametes hirsuta* using an air-lift bioreactor. *Lett. Appl. Microbiol.*, 42: 612-616.
- Cui, B.-K. Phylogenetic studies on *Phellinus*. NCBI/GenBank database Accession No. FJ627247.1
- Cui, B.-K. Phylogenetic studies on *Phellinus*. NCBI/GenBank database Accession No. FJ627248.1
- Cusano, A.M., Mekmouche, Y., Meglecz, E. and Tron, T. (2009). Plasticity of laccase generated by homeologous recombination in yeast. *FEBS. J.*, 276: 5471-5480.
- Dashtban, M., Schraft, H. and Qin, W. (2009). Fungal bioconversion of lignocellulosic residues; opportunities and perspectives. *Int. J. Biol. Sci.*, 5: 578-595.
- Dawson, L. and Boopathy, R. (2008). Cellulosic ethanol production from sugarcane bagasse without enzymatic saccharification. *Bioresources*, 3: 452-460.

- De Silva, R., Lago, E.S., Merheb, C.W., Macchione, M.M., Park, Y.K. and Gomes, E. (2005). Production of xylanase and CMCase on solid state fermentation in different residues by *Thermoascus aurantiacus* Mieh. *Braz. J. Microbiol.*, 36: 235-241.
- De Sousa, M.V., Monteiro, S.N. and d'Almeida, J.R.M. (2004). Evaluation of pretreatment, size and molding pressure on flexural mechanical behavior of chopped bagasse-polyester composites. *Polym. Testing*, 23: 253-258.
- Dong, J.L., Zhang, Y.W., Zhang, R.H., Huang, W.Z. and Zhang, Y.Z. (2005). Laccase gene from white rot fungus *Trametes gallica*. NCBI/GenBank database Accession No. AAW65485.1
- Dubois, M., Gilles, K. A., Hamilton, J. K., Rebers, P. A. and Smith, F. (1956). Colorimetric method for determination of sugars and related substances. *Anal. Chem.*, 28: 350–356.
- Durán, N., Rosa, M.A., D'Annibale, A. and Gianfreda, L. (2002). Applications of laccases and tyrosinases (phenoloxidases) immobilized on different supports: a review. *Enzyme Microb. Technol.*, 31: 907-931.
- Eggeman, T. and Elander, R.T. (2005). Process and economic analysis of pretreatment technologies. *Bioresour. Technol.*, 96: 2019-2025.
- Eggert, C., Temp, U. and Eriksson, K.E. (1996). The ligninolytic system of the white rot fungus *Pycnoporus cinnabarinus*: purification and characterization of the laccase. *Appl. Environ. Microbiol.*, 602: 1151-1158.

- Elisashvili, V., Kachlishvili, E. and Penninckx, M. (2008). Effect of growth substrate, method of fermentation, and nitrogen source on lignocelluloses-degrading enzymes production by white-rot basidiomycetes. *J. Ind. Microbiol. Biotechnol.*, 35: 1531-1538.
- Emert, G.H. (1991). Conversion of cellulose to chemicals including ethanol. In: Chahal D.S. (ed) Food Feed and Fuel from Biomass. Oxford and IBH. New Delhi, pp. 313-333.
- Emmel, A., Mathias, A.L., Wypych, F. and Ramos, L.P. (2003). Fractionation of *Eucalyptus grandis* chips by dilute acid-catalysed steam explosion. *Bioresour. Technol.*, 86: 105-115.
- Eken-Saraçoğlu, N. and Arslan, Y. (2000). Comparison of different pretreatments in ethanol fermentation using corn cob hemicellulosic hydrolysate with *Pichia stipitis* and *Candida shehatae*. *Biotechnol. Lett.*, 22: 855-858.
- Eriksson, K.E., Blanchette, R.A. and Ander, P. (1990). Microbial and Enzymatic Degradation of Wood and Wood Components. Springer-Verlag, Heidelberg, New York. pp. 397.
- Fang, H.Y., Chang, S.M., Hsieh, M.C. and Fang, T.J. (2007). Production, optimization growth conditions and properties of the xylanase from *Aspergillus carneus* M34. *J. Mol. Catal. B: Enzym.*, 49: 36-42.
- Farnet, A.M., Gil, G. and Ferre, E. (2008). Effects of pollutants on laccase activities of *Marasmius quercophilus*, a white-rot fungus isolated from Mediterranean sclerophyllous litter. *Chemosphere*, 70: 895-900.
- Faulon, J.-L., Carlson, G.A. and Hatcher, P.G. (1994). A three-dimensional model for lignocelluloses from gymnospermous wood. *Org. Geochem.*, 21: 1169-1179.

Fengel, D., and Wegener, G. (1983). Wood:Chemistry, Ultrastructure and Reactions. Walter de Gruyter & Cooperation. Berlin. pp. 613.

Fukuda, T., Uchida, H., Takashima, Y., Uwajima, T., Kawabata, T. and Suzuki, M. (2001). Degradation of bisphenol A by purified laccase from *Trametes villosa*. *Biochem. Biophys. Res. Commun.*, 284: 704-706.

Galbe, M. and Zacchi, G. (2002). A review of the production of ethanol from softwood. *Appl. Microbiol. Biotechnol.*, 59: 618-628.

Galbe, M. and Zacchi, G. (2007). Pretreatment of lignocellulosic materials for efficient bioethanol production. *Adv. Biochem. Eng. Biotechnol.*, 108: 41-65.

Galhaup, C., Goller, S., Peterbauer, C.K., Strauss, J. and Haltrich, D. (2002). Characterization of the major laccase isoenzyme from *Trametes pubescens* and regulation of its synthesis by metal ions. *Microbiol.*, 148: 2159-2169.

Gao, Z., Mori, T. and Kondo, R. (2012). The pretreatment of corn stover with *Gloeophyllum trabeum* KU-41 for enzymatic hydrolysis. *Biotechnol. Biofuels.*, 5: 28-38.

Gawande, P.V. and Kamat, M.Y. (1999). Production of *Aspergillus* xylanase by lignocellulosic waste fermentation and its application. *J. Appl. Microbiol.*, 879: 511-519.

Gerardi, V., Minelli, F. and Viggiano, D. (1998). Steam treated rice industry residues as an alternative feedstock for the wood based particle board industry in Italy. *Biomass Bioenergy*, 14: 295-299.

Germann, U.A., Muller, G., Hunziker, P.E. and Lerch, K. (1988). Characterization of two allelic forms of *Neurospora crassa* laccase. Amino- and carboxyl-terminal processing of a precursor. *J. Biol. Chem.*, 263: 885-896.

- Gianfreda, L., Xu, F. and Bollag, J.M. (1999). Laccase: a useful group of oxidoreductive enzymes. *Bioremediation J.*, 3: 1-25.
- Gilbert, H.J. and Hazlewood, G.P. (1993). Bacterial cellulases and xylanases. *J. Gen. Microbiol.*, 139: 187-194.
- Glenn, J.K., and Gold, M.H. (1983). Decolorization of several polymeric dyes by the lignin-degrading basidiomycete *Phanerochaete chrysosporium*. *Appl. Environ. Microbiol.*, 45: 1741-1747.
- Goh, T.K. (1999). Single-spore isolation using a hand-made glass needle. *Fungal Divers.*, 2: 47-63.
- Gomes, I., Gomes, J., Gomes, D.J. and Steiner, W. (2000). Simultaneous production of high activities of thermostable endoglucanase and beta-glucosidase by the wild thermophilic fungus *Thermoascus aurantiacus*. *Appl. Microbiol. Biotechnol.*, 53: 461-468.
- Gomes, A., Matsuo, T., Goda, K. and Ohgi, J. (2007). Development and effect of alkali treatment on tensile properties of curaua fiber green composites. *Composites: Part A*, 38: 1811-1820.
- Gong, C.S., Cao, N.J. Du, J. and Tsao, G.T. (1999). Ethanol production from renewable resources. *Adv. Biochem. Eng./Biotechnol.*, 65: 207-241.
- Gray, K.A., Zhao, L. and Emptage, M. (2006). Bioethanol. *Curr. Opin. Chem. Biol.*, 10: 141-146.
- Grohmann, K., Cameron, R.G. and Buslig, B.S. (1995). Fractionation and pretreatment of orange peel by dilute acid hydrolysis. *Bioresour. Technol.*, 54: 129-141.

- Gutierrez-Correa, M. and Tengerdy, R.P. (1998). Xylanase production by fungal mixed culture solid substrate fermentation on sugar cane bagasse. *Biotechnol. Lett.*, 20: 45-47.
- Han, L., Feng, J., Zhang, S., Ma, Z., Wang, Y. and Zhang, X. (2012). Alkali pretreatment of wheat straw and its enzymatic hydrolysis. *Braz. J. Microbiol.*, 43: 53-61.
- Hari, K.S., Janardhan, R.T. and Chowdary, G.V. (2001). Simultaneous saccharification and fermentation of lignocellulosic wastes to ethanol using a thermotolerant yeast. *Bioresour. Technol.*, 77: 193-196.
- Harnández, F.J.R., Marina, A., González, K., Vázquez, J. and Falcón, M.A. (2006). Production, partial characterization and mass spectrometric studies of the extracellular laccase activity from *Fusarium proliferatum*. *Appl. Microbiol. Biotechnol.*, 70: 212-221.
- Hans, E.S. and Klaus, P. (1996). On the interaction of lignin peroxidase with lignin. *Pure Appl. Chem.*, 66: 2089-2096.
- Harvey, P.J., Schoemaker, H.E., Bowen, R.M. and Palmer, J.M. (1985). Single-electron transfer processes and the reaction mechanism of enzymatic degradation of lignin. *FEBS Lett.*, 183: 13-16.
- Hawkins, G.M. and Doran-Peterson, J. (2011). A strain of *Saccharomyces cerevisiae* evolved for fermentation of lignocellulosic biomass displays improved growth and fermentative ability in high solids concentrations and in the presence of inhibitory compounds. *Biotechnol. Biofuels*, 4: 49-62.

Héla, Z.M., Tahar, M., Abdelhafidh, D., Sami, S. and Angel, T.M. (2006). Laccase purification and characterization from *Trametes trogii* isolated in Tunisia: Decolorization of textile dyes by the purified enzyme. *Enzyme Microb. Technol.*, 39: 141-148.

Hendriks, A.T.W.M. and Zeeman, G. (2009). Pretreatments to enhance the digestibility of lignocelluloses biomass. *Bioresour. Technol.*, 100: 10-18.

Hankin, L. and Anagnostakis, S. (1977). Solid media containing carboxy methyl cellulose to detect CM cellulase activity of microorganisms. *J. Gen. Microbiol.*, 98: 109 -115.

Hewitt, E.J. and Smith, T.A. (1975). Plant Mineral Nutrition. English, London University Press.

Higgins, D.G., Bleasby, A.J., Fuchs, R. (1992). CLUSTAL W: improved software for multiple sequence alignment. *Comput. Appl. Biosci.*, 8: 189-191.

Hongman, H., Jiti, Z., Jing, W., Cuihong, D. and Bin, Y. (2004). Enhancement of laccase production by *Pleurotus ostreatus* and its use for the decolorization of anthraquinone dye. *Process Biochem.*, 39: 1415-1419.

Hoshida, H., Nakao, M., Kanazawa, H., Kubo, K. and Hakukawa, T. (2001). Isolation of five laccase gene sequences from the white-rot fungus *Trametes sanguinea* by PCR, and cloning, characterization and expression of the laccase cDNA in yeasts. *J. Biosci. Bioeng.*, 92: 372-380.

Houtman, C.J. and Atalla, R.H. (1995). Cellulose-lignin interactions. *Plant Physiol.*, 107: 977-984.

- Hu, G., Heitmann, J.A. and Rojas O.J. (2008). Feedstock pretreatment strategies for producing ethanol from wood, bark, and forest residues. *Bioresources*, 3: 270-294.
- Imai, M., Ikari, K. and Suzuki, I. (2004). High-performance hydrolysis of cellulose using mixed cellulase species and ultrasonication pretreatment. *Biochem. Eng. J.*, 17: 79-83.
- Ismail, H., Nizam, J.M. and Abdul Khalil, H.P.S. (2001). The effect of a compatibilizer on the mechanical properties and mass swell of white rice husk ash led natural rubber/linear low density polyethylene blends. *Polym. Testing*, 20: 125-33.
- Jang, K.-Y., Cho, S.-M., Seok, S.-J., Kong, W.-S., Kim, G.-H. and Sung, J.-M. (2009). Screening of biodegradable function of indigenous ligno-degrading mushroom using dyes. *Mycobiology*, 37: 53-61.
- Jaouani, A., Tabka, M.G. and Penninckx, M.J. (2006). Lignin modifying enzymes of *Coriopsis polyzona* and their role in olive oil mill wastewaters decolourisation. *Chemosphere*, 62: 1421-1430.
- Jayawardhana, K. and Van, W.G.P. (2004). Modeling of carbonic acid pretreatment process using ASPEN-Plus. *Appl. Biochem. Biotechnol.*, 113-116: 1087-1102.
- Jecu, L. (2000). Solid state fermentation of agricultural; wastes for endoglucanase production. *Ind. Crops Prod.*, 11: 1-5.
- Jeewon, L. (1997). Biological conversion of lignocellulosic biomass to ethanol. *J. Biotechnol.*, 56: 1-24.

- Jia, L.D., and Yi, Z.Z. (2004). Purification and characterization of two laccase isoenzymes from ligninolytic fungus *Trametes gallica*. *Preparative Biochem. Biotechnol.*, 34: 179-194.
- Jianmin, G., Haibo, W., Daheng, Z., Mingxue, Y., Fangxia, G. and Yu, X. (2008). Production and characterization of cellulolytic enzymes from the thermoacidophilic fungal *Aspergillus terreus* M11 under solid-state cultivation of corn stover. *Bioresour. Technol.*, 99: 7623-7629.
- Jing, D., Li, P., Stagnitti, F. and Xiong, X. (2007). Optimization of laccase production from *Trametes versicolor* by solid fermentation. *Can. J. Microbiol.*, 53: 245-251.
- Jolival, C., Madzak, C., Brault, A., Caminade, E. and Malosse, C. (2005). Expression of laccase IIIb from the white-rot fungus *Trametes versicolor* in the yeast *Yarrowia lipolytica* for environmental applications. *Appl. Microbiol. Biotechnol.*, 66: 450-456.
- Jommuengbout, P., Phitsuwan, P., Pinitglang, S., Kyu, K.L. and Ratanakhanokchai, K. (2009). Xylooligosaccharide production from agricultural wastes by purified xylanase from alkaliphilic *Bacillus firmus* strain K-1. *Agri. Sci. J.*, 40: 301-304.
- José, M.O., María, J.N., Felicia, S., Ignacio, B., Paloma, M., Alberto, G., and Mercedes, B. (2006). Effects of acetic acid, furfural and catechol combinations on ethanol fermentation of *Kluyveromyces marxianus*. *Process Biochem.*, 41: 1223-1228.
- Joshi, B., Bhatt, M.R., Sharma, D., Joshi, J., Malla, R. and Sreerama, L. (2011). Lignocellulosic ethanol production: current practices and recent developments. *Biotechnol. Mol. Biol. Rev.*, 6: 172-182.

- Juliano, B.O. (1985). Rice Chemistry and Technology. American Association of Cereal Chemistry, New York.
- Jung, H., Xu, F. and Li, K. (2002). Purification and characterization of laccase from wood-degrading fungus *Trichophyton rubrum* LKY-7. *Enzyme Microb. Technol.*, 30: 161-168.
- Justo, A. and Hibbett, D.S. (2011). Phylogenetic classification of *Trametes* (*Basidiomycota*, *Polyporales*) based on a five-marker dataset. *Taxon*, 60: 1567-1583.
- Kabel, M.A. (2007). Effect of pretreatment severity on xylan solubility and enzymatic breakdown of the remaining cellulose from wheat straw. *Bioresour. Technol.*, 98: 2034-2042.
- Kalogeris, E., Christakopoulos, P., Kekos, D. and Macris, B.J. (1998). Studies on the solid-state production of thermostable endoxylanases from *Thermoascus aurantiacus*: Characterization of two isozymes. *J. Biotechnol.*, 60: 155-163.
- Kamal, G.M., Anwar, F., Hussain, A.I., Sarri, N. and Ashraf, M.Y. (2011). Yield and chemical composition of Citrus essential oils as affected by drying pretreatment of peels. *Inter. Food Res. J.*, 18: 1275-1282.
- Kandelbauer, A. and Güebitz, G.M. (2004). Bioremediation for the decolorization of textile dyes – a review, in Environmental Chemistry, ed. By Lichtfouse E., Schwarzbauer J. and Robert D. Springer, Berlin, pp. 269-288.
- Kang, S.-W., Park Y.-S., Lee J.-S., Hong S.-I. and Kim S.-W. (2004). Production of cellulase and hemicellulases by *Aspergillus niger* KK2 from lignocellulosic biomass. *Bioresour Technol.*, 91: 153-156.

Kang, H.-K. and Kim, D. (2012). Efficient bioconversion of rice straw to ethanol with TiO₂/UV Pretreatment. *Bioprocess Biosyst. Eng.*, 35: 43-48.

Kevin, A.G., Lishan, Z. and Mark, E. (2006). Bioethanol. *Curr. Opin. Chem. Biol.*, 10: 141-146.

Khanongnuch, C., Wanphrut, N., Lumyong, S. and Watanabe, T. (2004). Thermotolerant wood rotting fungi isolated from northern Thailand and their potential uses in lignin degrading applications. *Fungal Divers.*, 15: 187-196.

Khurana, S., Kapoor, M., Gupta, S. and Kuhad, R.C. (2007). Statistical optimization of alkaline xylanase production from *Streptomyces violaceoruber* under submerged fermentation using response surface methodology. *Indian J. Microbiol.*, 47: 144-152.

Kiiskinen, L.L. and Saloheim, M. (2004). Molecular cloning and expression in *Saccharomyces cerevisiae* of a laccase gene from the ascomycete *Melanocarpus albomyces*. *Appl. Environ. Microbiol.*, 70: 137-144.

Kim, Y.-K., Kwak, M.-S., Lee, W.H. and Choi, J.-W. (2000). Ultrasonic pretreatment for thermophilic aerobic digestion in industrial waste activated sludge treatment. *Biotechnol. Bioprocess Eng.*, 5: 469-474.

Kim, K.H., Tucker, M.P. and Nguyen, Q.A. (2002). Effects of pressing lignocellulosic biomass on sugar yield in two-stage dilute-acid hydrolysis process. *Biotechnol. Prog.*, 18: 489-494.

Kirk, T.K. (1987). Enzymatic 'combustion': the microbial degradation of lignin. *Ann. Rev. Microbiol.*, 41: 465-505.

Kocyigit, A., Baris, P.M., Yasa, I., Ozdemir, G. and Karaboz, I. (2012). Production of laccase from *Trametes trogii* TEM H2: a newly isolated white-rot fungus by air sampling. *J. Basic Microbiol.*, 52: 1-9.

Kongbuntad, W., Khanongnuch, C. and Lumyong, S. (2006^a). Efficacy of xylanase supplementation produced from *Thermoascus aurantiacus* SL16W in diet on Thai native chicken performance. *Int. J. Poultry Sci.*, 5: 463-469.

Kongbuntad, W., Seanphet, K., Khanongnuch, C. and Lumyong, S. (2006^b). Determination of toxic effects of crude xylanase derived from *Thermoascus aurantiacus* SL16W by hematology and blood biochemistry in Albino rats. *Pak. J. Biol. Sci.*, 9: 2255-2260.

Kornilłowicz-Kowalska, T. and Rybczyńska, K. (2012). Decolorization of Remazol Brilliant Blue (RBBR) and Poly R-478 dyes by *Bjerkandera adusta* CCBAS 930. *Central Eur. J. Biol.*, 7: 948-956.

Kuch, H.M. and Ballschmiter, K. (2001). Determination of endocrine disrupting phenolic compounds and estrogens in surface and drinking water by JRG- (NCI)-MS in the pictogram per liter range. *Environ. Sci. Technol.*, 35: 3201-3206.

Kuhad, R.C., Gupta, R. and Singh, A. (2011). Microbial cellulases and their industrial applications. *Enzyme Res.*, 1-10.

Kumar, P., Barrett, D.M., Delwiche, M.J. and Stroeve, P. (2009). Method for pretreatment of lignocellulosic biomass for efficient hydrolysis and biofuel production. *Ind. Eng. Chem. Res.*, 48: 3713-3729.

- Laura, L., Eliana, M. and Araceli, M.R. (2010). Effect of nitrogen sources and vitamins on ligninolytic enzyme production by some white-rot fungi. Dye decolorization by selected culture filtrates. *Bioresour. Technol.*, 12: 4554-4563.
- Lavarack, B.P., Giffin, G.J. and Rodman, D. (2002). The acid hydrolysis of sugarcane bagasse hemicelluloses to produce xylose, arabinose, glucose, and other products. *Biomass Bioenergy*, 23: 367-380.
- Lee, J. (1997). Biological conversion of lignocelluloses biomass to ethanol. *J. Biotechnol.*, 56: 1-24.
- Lee, Y.-J. and Shin, K.-S. (1999). Purification and properties of laccase of the white-rot basidiomycete *Coriolus hirsutus*. *J. Microbiol.*, 37: 148-153.
- Lee, J.-W., Gwak, K.-S., Park, J.-Y., Park, M.-J., Choi, D.-H., Kwon, M. and Choi, I.-G. (2007). Biological pretreatment of softwood *Pinus densiflora* by three white rot fungi. *J. Microbiol.*, 45: 485–491.
- Lee, Y.-J., Kim, B.-K., Lee, B.-H., Jo, J.-I., Lee, N.-K., Chung, C.-H., Lee, Y.-C. and Lee, J.-W. (2008). Purification and characterization of cellulase produced by *Bacillus amyoliquefaciens* DL-3 utilizing rice hull. *Bioresour. Technol.*, 99: 378-386.
- Lemos, J.L.S., Fontes, M.C.A. and Pereira, N.J., (2001). Xylanase production by *Aspergillus awamori* in solid-state fermentation and influence of different nitrogen sources. *Appl. Biochem. Biotechnol.*, 91: 681-689.
- Leonowicz, A., Matuszewska, A., Luterek, J., Ziegenhagen, D., Wojtaś-Wasilewska, M., Cho, N.-S., Hofrichter, M. and Rogalski, J. (2011). Biodegradation of lignin by white rot fungi. *Fungal Genet. Biol.*, 27: 175-185.

- Li, K., Helm, R.F. and Erikssen, K.E.L. (1998). Mechanistic studies of the oxidation of a non-phenolic lignin model compound by the laccase/1-hydroxybenzotriazole redox system. *Biotechnol. Appl. Biochem.*, 27: 239-243.
- Li, L., Wenkui, D., Peng, Y., Jian, Z. and Yinbo, Q. (2009). Decolorisation of synthetic dyes by crude laccase from *Rigidoporus lignosus* W1. *J. Chem. Technol. Biotechnol.*, 84: 399-404.
- Lin, Y. and Tanaka, S. (2006). Ethanol fermentation from biomass resources: current state and prospects. *Appl. Microbiol. Biotechnol.*, 69: 627-642.
- Lin, J.-Y., Wu, T.-Z. and Chou, J.-C. (2006). *In vitro* induction of fruiting body in *Antrodia cinnamomea*- a medicinally important fungus. *Microbiology*, 47: 267-272.
- Lin, C.-W., Tran, D.-T., Lai, C.-Y., I, Y.-P. and Wu, C.-H. (2010). Response surface optimization for ethanol production from *Pennisetum Alopecoider* by *Klebsiella oxytoca* THLC0409. *Biomass Bioenergy*, 34: 1922-1929.
- Linde, M., Galbe, M. and Zacchi, G. (2007). Simultaneous saccharification and fermentation of steam-pretreated barley straw at low enzyme loadings and low yeast concentration. *Enzyme Microb. Technol.*, 40: 1100-1107.
- Liou, T.-H. (2004). Preparation and characterization of nano-structured silica from rice husk. *Mat. Sci. Eng. A-Struct.*, 364: 313-323.
- Liu, W., Chao, Y., Liu, S., Bao, H. and Qian, S., (2003). Molecular cloning and characterization of a laccase gene from the basidiomycete *Fome lignosus* and expression in *Pichia pastoris*. *J. Appl. Microbiol. Biotechnol.*, 63: 174-181.

- Lloyd, T.A. and Wyman, C.E. (2005). Combined sugar yields for dilute sulfuric acid pretreatment of corn stover followed by enzymatic hydrolysis of the remaining solids. *Bioresour. Technol.*, 96: 1967-1977.
- Lomascolo, A., Cayol, J.-L., Roche, M., Guo, L., Robert, J.-L., Record, E., Lesage-Meessen, L., Ollivier, B., Sigoillot, J.-C. and Asther, M. (2002). Molecular clustering of *Pycnoporus* strains from various geographic origins and isolation of monokaryotic strains for laccase hyperproduction. *Mycol. Res.*, 106: 1193-1203.
- Ludueña, L., Fasce, D., Alvarez, V.A. and Stefani, P.M. (2011). Nanocellulose from rice husk following alkaline treatment to remove silica. Feasibility of ethanol production from coffee husks. *Bioresources*, 6: 1440-1453.
- Lynd, L.R., Zyl, W.H., McBride J.E. and Laser, M. (2005). Consolidated bioprocessing of cellulosic biomass: an update. *Curr. Opin. Biotechnol.*, 16: 577-583.
- Madhavi, S.R. and Lale, S.S. (2006). Increased production of extracellular laccase by the white rot fungus *Coriolus versicolor* MTCC 138. *World J. Microbiol. Biotechnol.*, 22: 921-926.
- Mais, U., Esteghlalian, A.R., Saddler, L.N. and Mansfield, S.D. (2002). Enhancing the enzymatic hydrolysis of cellulosic materials using simultaneous ball milling. *Appl. Biochem. Biotechnol.*, 98: 815-832.
- Mani, S., Tabil, L.G. and Sokhansanj, S. (2004). Grinding performance and physical properties of wheat and barley straws, corn stover and switchgrass. *Biomass Bioenergy*, 27: 339-352.

- Manjunathan, J., Kumar, M., Jenefar, S. and Kaviyarasan, V. (2010). Screening of inducers for laccase production by *Lentinus tuberregium* (fr) in liquid medium. *J. Biosci. Res.*, 1: 88-93.
- Mansur, M., Arias, M.E. Flärdh, M. and González, A.E. (2003). The white-rot fungus *Pleurotus ostreatus* secretes laccase isozymes with different substrate specificities. *Mycologia.*, 95: 1013-1020.
- Manzanares, P., Fajardo, S., and Martín, C. (1995). Production of ligninolytic activities when treating paper pulp effluents by *Trametes versicolor*. *J. Biotechnol.*, 43: 125-132.
- Maria, T.C., Antonio, C., Santa, R. and Enrico, R. (2000). Production, purification, and properties of an extracellular laccase from *Rigidoporus lignosus*. *Protein Expres. Purif.*, 18: 141-147.
- Mario, C.N.S., Francisco. G., Angélica, M.A., Angel, T.M. and María, J.M. (2002). Induction, isolation, and characterization of two laccase from the white rot basidiomycete *Coriolopsis rigida*. *Appl. Environ. Microbiol.*, 68: 1534-1540.
- Martinello, T., Kaneko, T.M., Velasco, M.V.R., Taqueda, M.E.S. and Consiglieri, V.O. (2006). Optimization of poorly compactable drug tablets manufactured by direct compression using the mixture experimental design. *Int. J. Pharm.*, 322: 87-95.
- Mariam, L., Diego, M. and Ángeles, S. (2006). Effect of heavy metals on the production of several laccase isoenzymes by *Trametes versicolor* and on their ability to decolourise dyes. *Chemosphere*, 63: 912-917.

- Marzullo, L., Cannio, R., Giardina, P., Santini, M.T. and Sannia, G. (1995). Veratryl alcohol oxidase from *Pleurotus ostreatus* participates in lignin degradation and prevents polymerization of laccase-oxidized substrates. *J. Biol. Chem.*, 270: 3823-3827.
- Masutti, D.C., Borgognone, A. and Setti, L. (2012). Production of enzymes from rice husks and wheat straw in solid state fermentation. *Chem. Eng. Trans.*, 27: 133-138.
- Michizoe, J., Ichinose, H., Kamiya, N., Maruyama, T. and Goto, M. (2005). Biodegradation of phenolic environmental pollutants by a surfactant-laccase complex in organic media. *J. Biosci. Bioeng.*, 99: 642-647.
- Mikhail, M.A., Daba, A.S. and El-Assar, S.A. NCBI/GenBank database Accession No. JN854132.1
- Mikiashvili, N., Wasser, S.P., Nevo, E. and Elisashvili, V. (2006). Effects of carbon and nitrogen sources on *Pleurotus ostreatus* ligninolytic enzyme activity. *World J. Microbiol. Biotechnol.*, 22: 999-1002.
- Milala, M.A., Shugaba, A., Gidado, A., Ene, A.C. and Wafar, J.A. (2005). Studies on the use of agricultural wastes for cellulase enzyme production by *Aspergillus niger*. *Res. J. Agric. & Biol. Sci.*, 1: 325-328.
- Miller, G.L. (1959). Use of dinitrosalicylic acid reagent for determination of reducing sugar. *Anal. Chem.*, 31: 426-428.
- Minnussi, R.C., Pastore, G.M. and Duran, N. (2002). Potential application of laccase in the food industry. *Trends Food Sci. Technol.*, 13: 205-216.

Moros, A.S., Gozávez-Zafrilla, J.M., Lora-García, J. and García-Díaz, J.C. (2005).

Mixture design applied to describe the influence of ionic composition on the removal of nitrate ions using nanofiltration. *Desalination*, 185: 289-296.

Mosier, N.S., Wyman C. Dale, B., Elander, R., Lee Y.Y., Holtzapple, M. and Ladisch, M.R. (2005). Features of promising technologies for pretreatment of lignocellulosic biomass. *Bioresour. Technol.*, 96: 673-686.

Mtui, G.Y.S. (2009). Recent advances in pretreatment of lignocellulosic wastes and production of value added products. *Afr. J. Biotechnol.*, 8: 1398-1415.

Mudgett, R. (1986). Manual of Industrial Microbiology and Biotechnology. Demain, A. and Solomon, N. (Eds.). American Society for Microbiology, Washington, AD., pp: 66-83.

Murthy, P.S. and Manonmani, H.K. (2008). Bioconversion of coffee industry wastes with white rot fungi *Pleurotus florida*. *Res. J. Environ. Sci.*, 2: 145-150.

Murthy, P.S. and Naidu. M.M. (2010). Production and application of xylanase from *Penicillium* sp. utilizing coffee by-products. *Food Bioprocess Technol.*, 5: 657-664.

Murugesan, K., Arulmani, M., Nam, I.-H., Kim, Y.-M. and Chang, Y.-S. (2006). Purification and characterization of laccase produced by a white rot fungus *Pleurotus sajor-caju* under submerged culture condition and its potential in decolorization of azo dyes. *Appl. Microbiol. Biotechnol.*, 72: 939-946.

Nagai, M., Sato, T., Watanabe, H., Saito, K. and Kawata, M. (2002). Purification and characterization of an extracellular laccase from the edible mushroom *Lentinula edodes*, and decolorization of chemically different dyes. *Appl. Microbiol. Biotechnol.*, 60: 327-335.

- Naki, A. and Varfolomeev, S.D. (1981). Inhibition mechanism of *Polyporus versicolor* laccase by halide ions. *Biochem.*, 46: 1694-1702.
- Navia, D.P., Velasco, M.R.J. and Hoyos, C.J.L. (2011). Production and evaluation of ethanol from coffee processing by-products. *J. Vitae.*, 18: 287-294.
- Nicole, M., Chamberland, H., Geiger, J.P., Lecours, N. and Valéro, J. (1992). Immunocytochemical localization of laccase in wood decayed by *Rigidoporus lignosus*. *Appl. Environ. Microbiol.*, 58:1727-1739.
- Nitheranont, T., Watanabe, A. and Asada, Y. (2011^a). Extracellular laccase produced by an edible basidiomycetous mushroom, *Grifola frondosa*: purification and characterization. *Biosci. Biotechnol. Biochem.*, 75: 538-543.
- Nitheranont, T., Watanabe, A., Suzuki, T., Katayama, T., and Asada, Y. (2011^b). Decolorization of synthetic dyes and biodegradation of bisphenol A by laccase from the edible mushroom, *Grifola frondosa*. *Biosci. Biotechnol. Biochem.*, 75: 1845-1847.
- Núñez, M and Ryvarden, L. (2001). East Asian Polypores Vol. 2. Polyporaceae Synopsis Fungorum 14. Fungiflora, Oslo
- Okamoto, K., Yanagi, S.O. and Sakai, T. (2000). Purification and characterization of extracellular laccase from *Pleurotus ostreatus*. *Mycosci.*, 41: 7-13.
- Oscar, J.S. and Carlos, A.C. (2008). Trends in biotechnological production of fuel ethanol from different feedstocks. *Bioresour Technol.*, 99: 5270-5295.
- Osma, J.F., Toca, J.L. and Couto, S.R. (2007). Banana skin: A novel waste for laccase production by *Trametes pubescens* under solid-state conditions. Application to synthetic dye decolouration. *Dyes Pigm.*, 75: 32-37.

- Palmqvist, E. and Hahn-Hägerdal, B. (2000). Fermentation of lignocellulosic hydrolysates I: inhibition and detoxification. *Bioresour. Technol.*, 74: 17-24.
- Pandey, A. (1994). Solid state fermentation: An overview. In: Solid Fermentation, Ashok Pandey (ed.), Wiley Eastern: New Delhi.
- Pandey, A., Selvakumar, P., Soccol, C.R. and Nigam, P. (1999). Solid state fermentation for the production of industrial enzymes. *Curr. Sci.*, 77:149-162.
- Pandey, A., Soccol, C.R., Nigam, P., Brand, D., Mohan, R. and Roussos, S. (2000). Biotechnological potential of coffee pulp and coffee husk for bioprocesses. *Biochem. Eng. J.*, 6: 153-162.
- Papinutti, V.L. and Forchiassin, F. (2007). Lignocellulolytic enzymes from *Fomes sclerodermeus* growing in solid-state fermentation. *J. Food Eng.*, 81: 54-59.
- Parani, K. and Eyini, M. (2010). Effect of co-fungal treatment on biodegradation of coffee pulp waste in solid state fermentation. *Asian J. Exp. Biol. Sci.*, 1: 352-359.
- Parani, K. and Eyini, M. (2012). Production of ligninolytic enzymes during solid state fermentation of coffee pulp by selected fungi. *Sci. Res. Rep.*, 2: 202-209.
- Park, Y.S., Kang, S.W., Lee, J.S., Hong, S.I. and Kim, S.W. (2002). Xylanase production in solid state fermentation by *Aspergillus niger* mutant using statistical experimental design. *Appl. Microbiol. Biotechnol.*, 58: 762-766.
- Patle, S. and Lal, B. (2007). Ethanol production from hydrolysed agricultural wastes using mixed culture of *Zymomonas mobilis* and *Candida tropicalis*. *Biotechnol. Lett.*, 29: 1839-1843.
- Patrick, F., Mtui, G., Mshandete, A.M., Johansson, G. and Kivaisi, A. (2009). Purification and characterization of a laccase from the basidiomycete *Funalia trogii* (Berk.) isolated in Tanzania. *Afr. J. Biochem. Res.*, 3: 250-258.

- Pietrobon, V.C., Monteiro, R.T.R., Pompeu, G.B., Borges, E.P., Lopes, M.L., Amorim, H.V., Cruz, S.H. and Viégas, E.K.D. (2011). Enzymatic hydrolysis of sugarcane bagasse pretreated with acid or alkali. *Braz. Arch. Biol. Technol.*, 54: 229-233.
- Pointing, S.B. (1999). Qualitative methods for the determination of lignocellulolytic enzyme production by tropical fungi. *Fungal Divers.*, 2: 17-33.
- Pothast, A., Rosenau, T. and Fischer, K. (2001). Oxidation of benzyl alcohols by the laccase-mediator system (LMS)- a comprehensive kinetic description. *Holzforschung*, 55: 47-56.
- Potumarthi, R., Baadhe, R.R., Nayak, P., and Jetty, A. (2012). Simultaneous pretreatment and saccharification of ricehusk by *Phanerochete chrysosporium* for improved production of reducing sugars. *Bioresour. Technol.*, 22 (in press)
- Pradeep, P., Reddy, O.V.S. and Ko, P.R.M.S. (2012). Process optimization for ethanol production from very high gravity (VHG) finger millet medium using response methodology. *Iranian J. Biotechnol.*, 10: 168-174.
- Raeder, U. and Broda, P. (1985). Rapid preparation of DNA from filamentous fungi. *Lett. Appl. Microbiol.*, 1: 17-20.
- Ramos, J., Rojas, T., Navarro, F., Valos, F.D., Sanjuaán, R., Rutaga, J., Young, R.A., 2004. Enzymatic and fungal treatments on sugarcane bagasse for the production of mechanical pulps. *J. Agric. Food Chem.*, 52: 5057-5062.
- Raval, V.H., Sanghvi, G.V., Gohil, A.R., Rajput, K.S., Faterpekar, S. and Deshmukh, S.K. NCBI/GenBank database Accession No. GQ906769.1

- Rezende, C.A., Lima, M.A., Maziero, P., Azevedo, E.R., Garcia, W. and Polikarpov, I. (2011). Chemical and morphological characterization of sugarcane bagasse submitted to delignification process for enhanced enzymatic digestibility. *Biotechnol. Biofuels*, 4: 54-71.
- Rim, K., Lasssd, B., Steve, W., Mariem, E., Abdelhafidh, D., Sami, S. and Tahar, M. (2010). Decolourization and detoxification of textile industry wastewater by the laccase-mediator system. *J. Hazard. Mat.*, 175: 802-808.
- Rodríguez, C.S., and Sanromán, M.A. (2005). Application of solid-state fermentation to ligninolytic enzyme production. *Biochem. Eng. J.*, 22: 211-219.
- Rodríguez, C.S., Sanromán, Ma. and Gübitz, G.M. (2005). Influence of redox mediators and metal ions on synthetic dye decolorization by crude laccase from *Trametes hirsute*. *Chemosphere*, 58: 417-422.
- Rosales, E., Couto, S.R. and Sanromán, A., 2002. New uses of food waste: application to laccase production by *Trametes hirsuta*. *Biotechnol. Lett.*, 24: 701-704.
- Rosana, C.M., Márcio, A.M., José, A.S., Carmen, V.F. and Hiroshi, A. (2007). Purification, characterization and application of laccase from *Trametes versicolor* for colour and phenolic removal of olive mil wastewater in the presence of 1-hydroxybenzotriazole. *Afr. J. Biotechnol.*, 6: 1248-1254.
- Royer, J.C. and Nakas, J.P. (1989). Xylanase production by *Trichoderma longibrachiatum*. *Enzyme Microb. Technol.*, 11: 405-410.
- Ryu, S.H., Lee, A.Y. and Kim, M. (2008). Molecular characteristics of two laccase from the basidiomycete fungus *Polyporus brumalis*. *J. Microbiol.*, 46: 62-69.

- Saito, T., Kato, K., Yokogawa, Y., Nishida, M. and Yamashita, N. (2004). Detoxification of bisphenol A and nonylphenol by purified extracellular laccase from a fungus isolated from soil. *J. Biosci. Bioeng.*, 98: 64-66.
- Saitou, N., and Nei, M. (1987). The neighbor-joining method: a new method for reconstructing phylogenetic trees. *Mol. Biol. Evol.*, 4: 406-425.
- Salvachúa, D., Prieto, A., López-Abelairas, M., Lu-Chau, T., Martínez, A.T. and Martínez, M.J. (2011). Fungal pretreatment: An alternative in second-generation ethanol from wheat straw. *Bioresour. Technol.*, 102: 7500-7506.
- Sánchez, C. (2009). Lignocellulosic residues: Biodegradation and bioconversion by fungi. *Biotechnol. Adv.*, 27: 185-194.
- Sanghvi, G.V., Koyani, R.D. and Rajput, K.S. (2010). Thermostable xylanase production and partial purification by solid-state fermentation using agricultural waste wheat straw. *Mycology*, 1: 106-112.
- Sanjeev, K.S., Krishan, L.K., and Harmeet, S.G. (2002). Fermentation of enzymatically saccharified sunflower stalks for ethanol production and its scale up. *Bioresour. Technol.*, 85: 31-33.
- Saparrat, M.C.N., Guillén, F., Arambarri, A.M., Martínez, A.T. and Martínez, M.J. (2002). Induction, isolation, and characterization of two laccase from the white rot basidiomycete *Coriolopsis rigida*. *Appl. Environ. Microbiol.*, 68: 1534-1540.
- Sasaki, C., Takada, R., Watanabe, T., Honda, Y., Karita, S., Nakamura, Y. and Watanabe, T. (2011). Surface carbohydrate analysis and bioethanol production of sugarcane bagasse pretreated with the white rot fungus, *Ceriporiopsis subvermispora* and microwave hydrothermolysis. *Bioresour. Technol.*, 21: 9942-9946.

- Sasikumar, E. and Viruthagiri, T. (2008). Optimization of process conditions using response surface methodology (RSM) for ethanol production from pretreated sugarcane bagasse: kinetics and modeling. *Bioenerg. Res.*, 1: 239-247.
- Sasikumar, E. and Viruthagiri, T. (2010). Simultaneous saccharification and fermentation (SSF) of sugarcane bagasse - kinetics and modeling. *Int. J. Chem. Biol. Eng.*, 3: 57-64.
- Sathish, T., Lakshmi, G.S., Rao, Ch.S., Brahmaiah, P. and Prakasham, R.S. (2008). Mixture design as first step for improved glutaminase production in solid-state fermentation by isolated *Bacillus* sp. RSP-GLU. *Lett. Appl. Microbiol.*, 47: 256-262.
- Schoemaker, H.E., Harvey, P.J., Bowen, R.M., and Palmer, J.M. (1985). On the mechanism of enzymatic lignin breakdown. *FEBS Lett.*, 183: 7-12.
- Schurz, J. (1978). Bioconversion of cellulosic substances into energy chemicals and microbial protein. Ghose, T.K.; Ed.: New Delhi. pp. 37.
- Selvankumar, T., Govarthanan, M. and Govindaraju, M. (2011). Endoglucanase production by *Bacillus amyloliquefaciens* using coffee pulp as substrate in solid state fermentation. *Int. J. Pharm. Bio. Sci.*, 2: 355-362.
- Shaul, G.M., Holdsworth, T.J., Dempsey, C.R. and Dostal, K.A. (1991). Fate of water soluble azo dyes in the activated sludge process. *Chemosphere*, 22: 107-119.
- Shi, J., Sharma-Shivappa, R.R., Chinn, M. and Howell, N. (2009). Effect of microbial pretreatment on enzymatic hydrolysis and fermentation of cotton stalks for ethanol production. *Biomass Bioenergy*, 33, 88-96.

Silverstein, R.A., Chen, Y., Sharma, -S.R.R., Boyette, M.D. and Osborne, J. (2007).

A comparison of chemical pretreatment methods for improving saccharification of cotton stalks. *Bioresour. Technol.*, 98: 3000-3011.

Slavin, J.L., Brauer, P.M. and Marlett, J.A. (1981). Neutral detergent fiber, hemicellulose and cellulose digestibility in human subjects. *J. Nutrition*, 111: 287-297.

Snedecor, G.W. and Cochran, W.G. (1989). Statistical Methods, Eighth Edition, Iowa State University Press. pp. 503.

Solomon, E.I., Sundaram, U.M. and Machonkin, T.E. (1996). Multicopperoxidases and oxygenases. *Chem. Rev.*, 96:2563-2605.

Sonali, P. and Banwari, L. (2007). Ethanol production from hydrolysed agricultural waste using mixed culture of *Zymomonas mobilis* and *Candida tropicalis*. *Biotechnol. Lett.*, 29: 1839-1843.

Souza, J.V.B., Silva, E.S., Cavallazzi, J.R.P. and Sobrinho, A.S. (2010). Formulation of a liquid medium with wheat bran for the production of laccase by *Trametes versicolor* in an air-lift bioreactor. *J. Food Agri. Environ.*, 8: 394-396.

Sugumaran, M. (1995). A caution about the azide inhibition of enzymes associated with electrophilic metabolites. *Biochem. Biophys. Res. Com.*, 212: 834-839.

Sun, Y. and Cheng, J. (2002). Hydrolysis of lignocellulosic materials for ethanol production: a review. *Bioresour. Technol.*, 83: 1-11.

Sun, W.-L. and Tao, W.-Y. (2010). Comparison of cell growth and ethanol productivity on different pretreatment of rice straw hemicelluloses hydrolysate by using *Candida shehatae* CICC 1766. *Afr. J. Microbiol. Res.*, 4: 1105-1109.

- Sung, Y.J. and Shin, S.-J. (2011). Compositional changes in industrial hemp biomass (*Cannabis sativa* L.) induced by electron beam irradiation pretreatment. *Biomass Bioenergy*, 35: 3267-3270.
- Svarachorn, A. (1999). Production of fungal-xylanase using agricultural waste by solid state fermentation. *J. Sci. Res. Chulalongkorn Univ.*, 24: 13-20.
- Syafri, R., Ahmad, I. and Abdullah, I. (2011). Effect of rice husk surface modification by LENR on the mechanical properties of ND/HDPE reinforced rice husk composite. *Sains Malaysiana*, 40: 749-756.
- Taherzadeh, M.J. and Karimi, K. (2007). Acid-based hydrolysis processes for ethanol from lignocellulosic materials: A review. *Bioresources*, 2: 472-499.
- Taherzadeh, M.J. and Karimi, K. (2008). Pretreatment of lignocellulosic wastes to improve ethanol and biogas production: A review. *Int. J. Mol. Sci.*, 9: 1621-1651.
- Taiz, L. and Zieger, E. (2010). *Plant Physiology*. Sinauer Associates, Inc., Publishers. Sunderland. pp. 782.
- Tamura, K., Dudley, J., Nei, M. and Kumar, S. (2007). MEGA4: molecular evolutionary genetics analysis (MEGA) software version 4.0. *Mol. Biol. Evol.*, 24: 1596-1599.
- Tanabe, N., Sagawa, I., Ohtsubo, K., Iijima, Y. and Yanagi, S.O. (1989). Comparison of phenoloxidase activities during the cultivation. *Agric. Biol. Chem.*, 53: 3061-3063.
- Tanahashi, M., Takada, S., Aoki, T., Goto, T., Higuchi, T., Hanai, S. 1983. Characterization of explosion wood. 1. Structure and physical properties. *Wood Res.*, 69: 36-51.

Tarley, C.R.T., Ferreira, S.L.C. and Arruda, M.A.Z. (2004). Use of modified rice husks as a natural solid adsorbent of trace metals: characterization and development of an on-line preconcentration system for cadmium and lead determination by FAAS. *Microchem. J.*, 77: 163-175.

Tassinari, T. and Macy, C. (1997). Differential speed two roll mill pretreatment of cellulosic materials for enzymatic hydrolysis. *Biotechnol. Bioeng.*, 19: 1321-1330.

Tebka, M.G., Herpoel-Gimbert, I., Monod, F., Asther, M. and Sigoillot, J.C. (2006). Enzymatic saccharification of wheat straw for bioethanol production by a combined cellulase xylanase and feruloyl esterase treatment. *Enzyme Microb. Technol.*, 39: 897-902.

Techapun, C., Sinsuwongwat, S., Watanabe, M., Sasaki, K. and Poosaran, N. (2002). Production of cellulase-free xylanase by a thermotolerant *Streptomyces* sp. grown on agricultural waste and media optimization using mixture design and Plackett – Burman experimental design methods. *Biotechnol. Lett.*, 24:1437–1442.

Teymouri, F., Laureano, -P.L., Alizadeh, H. and Dale, B.E. (2005). Optimization of the ammonia fiber explosion (AFEX) treatment parameters for enzymatic hydrolysis of corn stover. *Bioresour. Technol.*, 96: 2014-2018.

Thomsen, M.H., Thygesen, A. and Thomsen, A.B. (2008). Hydrothermal treatment of wheat straw at pilot plant scale using a three-step reactor system aiming at high hemicelluloses recovery, high cellulose digestibility and low lignin hydrolysis. *Bioresour. Technol.*, 99: 4221-4228.

Thurston, C.F. (1994). The structure and function of fungal laccases. *Microbiol.* 140: 19-26.

Tien, M. and Kirk, T.K. (1983). Lignin-Degrading enzyme from Hymenomycetes *Phanerochaete chrysosporium* Burds. *Science*, 221: 661-663.

Tišma, M., Žnidaršič-Plazl, P., Vasić-Rački, Đ. and Zelić, B. (2012). Optimization of laccase production by *Trametes versicolor* cultivated on industrial waste. *Appl. Biochem. Biotechnol.*, 166: 36-46.

Uchida, H., Fukuda, T., Miyamoto, H., Kawabata, T., Suzuki, M. and Uwajima, T. (2001). Polymerization of bisphenol A by purified laccase from *Trametes villosa*. *Biochem. Biophys. Res. Commun.*, 287: 355-358.

Verga, E., Reczey, K. and Zacchi, G. (2004). Optimization of steam pretreatment of corn stover to enhance enzymatic digestibility. *Appl. Biochem. Biotechnol.*, 113: 509-523.

Ververis, C., Georghiou, K., Danielidis, D., Hatzinilolaou, D.G., Santas, P., Santas, R. and Corleti, V. (2007). Cellulose, hemicelluloses, lignin and ash content of some organic materials and their suitability for use as paper pulp supplements. *Bioresour. Technol.*, 98: 296-301.

Viikari, L., Alapuranen, M., Puranen, T., Vehmaanpera, J., Siika-Aho, M. (2007). Thermostable enzymes in lignocellulose hydrolysis. *Adv. Biochem. Eng. Biotechnol.* 108: 121-145.

Vimalashanmugam, K. and Viruthagiri T. (2012). Statistical optimization of media components for xylanase production by *aspergillus fumigatus* using SSF. *Int. J. Eng. Res. Appl.*, 2: 1320-1329.

Wells, A., Teria, M. and Eve, T. (2006). Green oxidations with laccase-mediator systems. *Biochem. Soc. Trans.*, 34: 304-308.

- Wen, Z., Liao, W. and Chen, S. (2005). Production of cellulase/ β -glucosidase by the mixed fungi culture *Trichoderma reesie* and *Aspergillus phoenicis* on dairy manure. *Process Biochem.*, 40: 3087-3094.
- Wenster-Botz, D. 2000. Experimental design for fermentation media development: Statistical design or global random search. *J. Biosc. Bioeng.*, 90: 473-483.
- White, T.J., Bruns, T., Lee, S. and Taylor, J.W. (1999). Amplification and direct sequencing of fungal ribosomal RNA genes for phylogenetics. In: PCR protocols: A guide to methods and applications, eds. Innis M.A., Gelfand D.H., Sninsky J.J., White T.J. Academic Press, Inc. New York, pp. 315-322.
- Wong, K.K.Y., Nelson, S.L. and Saddler, J.N. (1996). Xylanase treatment for the peroxide bleaching of oxygen delignified kraft pulps derived from three softwood species. *J. Biotechnol.*, 48: 137-145.
- Xiao, W. and Clarkson, W.W. (1997). Acid solubilization of lignin and bioconversion of treated newsprint to methane. *Biodegrad.*, 8: 61-66.
- Xu, Z., Wang, Q., Jiang, Z., Yang, X.-X. and Ji, Y. (2007). Enzymatic hydrolysis of pretreated soybean straw. *Biomass Bioenergy*, 31: 162-167.
- Yang, C., Shen, Z., Yu, G. and Wang, J. (2008). Effect and after effect of γ radiation pretreatment on enzymatic hydrolysis of wheat straw. *Bioresour. Technol.*, 99: 6240-6245.
- Yoon, L.W., Ngoh, G.C., Chua, A.S.M., Hashim, M.A. (2011). Comparison of ionic liquid, acid and alkali pretreatments for sugarcane bagasse enzymatic saccharification. *J. Chem. Technol. Biotechnol.*, 86: 1342-1348.

- Yu, J., Zhang, X. and Tan, T. (2009). Optimization of media conditions for the production of ethanol from sweet sorghum juice by immobilized *Saccharomyces cerevisiae*. *Biomass Bioenergy*, 33: 521-526.
- Yue, F., Lan, W., Zhang, A., Liu, C., Sun, R. and Ye, J. (2012). Dissolution of holocellulose in ionic liquid assisted with ball-milling pretreatment and ultrasound irradiation. *Bioresources*, 7: 2199-2208.
- Zaldivar, J., Nielsen, J. and Olsson, L. (2001). Fuel ethanol production from lignocellulose: a challenge for metabolic engineering and process integration. *Appl. Microbiol. Biotechnol.*, 56:17–34.
- Zemnukhova, L.A., Tomshich, S.V., Mamontova, V.A., Komandrova, N.A., Fedorishcheva, G.A. and Sergienko, V.I. (2004). Composition and properties of polysaccharides from rice husk. *Russ. J. Appl. Chem.*, 77: 1883-1887.
- Zhang, X., Xu C. and Wang, H. (2007). Pretreatment of bamboo residues with *Coriolus versicolor* for enzymatic hydrolysis. *J. Biosci. Bioeng.*, 104: 149-151.
- Zhao, X., Zhang, L. and Lui, D. (2007). Comparative study on chemical pretreatment methods for improving enzymatic digestibility of crofton weed stem. *Bioresour. Technol.*, 99: 3729-3736.
- Zheng, Y., Pan, Z. and Zhang, R. (2009). Overview of biomass pretreatment for cellulosic ethanol production. *Int. J. Agric. Biol. Eng.*, 2: 51-68.
- Zimbardi, F., Viola, E., Nanna, F., Larocca, E., Cardinale, M. and Barisano, D. (2007). Acid impregnation and steam explosion of corn stover in batch processes. *Ind. Crops Prod.*, 26: 195-206.

Zorba, O. and Kurt, S. (2006). Optimization of emulsion characteristics of beef, chicken and turkey meat mixtures in model system using mixture design. *Meat Science*, 73: 611-618.

Zhu, S.D., Yu, Z.N., Wu, Y.X., Zhang, X., Li, H. and Gao, M. (2005). Enhancing enzymatic hydrolysis of rice straw by microwave pretreatment. *Chem. Eng. Commun.*, 192: 1559-1566.

Zhu, Z., Sathitsuksanoh, N., Vinzant, T., Schell, D.J., McMillan, J.D. and Zhang, Y.H. (2009). Comparative study of corn stover pretreated by dilute acid and cellulose solvent-based lignocellulose fractionation: enzymatic hydrolysis, supramolecular structure, and substrate accessibility. *Biotechnol. Bioeng.*, 103: 715-724.