

BIBLIOGRAPHY

1. Sohn W, Ismail AI. Regular dental visits and dental anxiety in an adult dentate population. *J Am Dent Assoc* 2005; 136: 58-66.
2. van Wijk AJ, Hoogstraten J. Anxiety and pain during dental injections. *J Dent* 2009; 37: 700-4.
3. Marsac ML, Funk JB. Relationships among psychological functioning, dental anxiety, pain perception, and coping in children and adolescents. *J Dent Child (Chic)* 2008; 75: 243-51.
4. Malamed SF. Sedation: A Guide to Patient Management. 5 ed: Mosby, 2009.
5. Weaver JM. ADA sedation-anesthesia guidelines revisions deserve our support. *Anesth Prog* 2007; 54: 99.
6. Morris Clark AB. Handbook of Nitrous Oxide and Oxygen Sedation. MOSBY, 2008.
7. Holroyd I. Conscious sedation in pediatric dentistry. A short review of the current UK guidelines and the technique of inhalational sedation with nitrous oxide. *Paediatr Anaesth* 2008; 18: 13-7.
8. Clark MS, Campbell SA, Clark AM. Technique for the administration of nitrous oxide/oxygen sedation to ensure psychotropic analgesic nitrous oxide (PAN) effects. *Int J Neurosci* 2006; 116: 871-7.
9. Clark M. Handbook of Nitrous Oxide and Oxygen Sedation: MOSBY 1999.
10. Wilson S. A survey of the American Academy of Pediatric Dentistry membership: nitrous oxide and sedation. *Pediatr Dent* 1996; 18: 287-93

11. Burton JH, Auble TE, Fuchs SM. Effectiveness of 50% nitrous oxide/50% oxygen during laceration repair in children. *Acad Emerg Med* 1998; 5: 112-7.
12. Gerhardt RT, King KM, Wiegert RS. Inhaled nitrous oxide versus placebo as an analgesic and anxiolytic adjunct to peripheral intravenous cannulation. *Am J Emerg Med* 2001; 19: 492-4.
13. Fauroux B, Onody P, Gall O, Tourniaire B, Koscielny S, Clement A. The efficacy of premixed nitrous oxide and oxygen for fiberoptic bronchoscopy in pediatric patients: a randomized, double-blind, controlled study. *Chest* 2004; 125: 315-21.
14. Rosen MA. Nitrous oxide for relief of labor pain: a systematic review. *Am J Obstet Gynecol* 2002; 186: S110-26.
15. Collado V, Hennequin M, Faulks D, Mazille MN, Nicolas E, Koscielny S, Onody P. Modification of behavior with 50% nitrous oxide/oxygen conscious sedation over repeated visits for dental treatment a 3-year prospective study. *J Clin Psychopharmacol* 2006; 26: 474-81.
16. Hennequin M, Collado V, Faulks D, Koscielny S, Onody P, Nicolas E. A clinical trial of efficacy and safety of inhalation sedation with a 50% nitrous oxide/oxygen premix (Kalinox) in general practice. *Clin Oral Investig* 2010; 16: 633-42.
17. Collado V, Nicolas E, Faulks D, Tardieu C, Maniere MC, Droz D, Onody P, Hennequin M. Evaluation of safe and effective administration of nitrous oxide after a postgraduate training course. *BMC Clin Pharmacol* 2008; 8: 3.

18. Paterson SA, Tahmassebi JF. Paediatric dentistry in the new millennium: 3. Use of inhalation sedation in paediatric dentistry. *Dent Update* 2003; 30: 350-6, 8.
19. Gamis AS, Knapp JF, Glenski JA. Nitrous oxide analgesia in a pediatric emergency department. *Ann Emerg Med* 1989; 18: 177-81.
20. O'Sullivan I, Benger J. Nitrous oxide in emergency medicine. *Emerg Med J* 2003; 20: 214-7.
21. Quarnstrom FC, Mar RS. A premix of 50% nitrous oxide - 50% oxygen for sedation during dental procedures. *Anesth Prog* 1983; 30: 197-8.
22. Onody P, Gil P, Hennequin M. Safety of inhalation of a 50% nitrous oxide/oxygen premix: a prospective survey of 35 828 administrations. *Drug Saf* 2006; 29: 633-40.
23. Gall O, Annequin D, Benoit G, Glabeke E, Vrancea F, Murat I. Adverse events of premixed nitrous oxide and oxygen for procedural sedation in children. *Lancet* 2001; 358: 1514-5.
24. Jeske AH, Whitmire CW, Freels C, Fuentes M. Noninvasive assessment of diffusion hypoxia following administration of nitrous oxide-oxygen. *Anesth Prog* 2004; 51: 10-3.
25. Jackson DL, Johnson BS. Conscious sedation for dentistry: risk management and patient selection. *Dent Clin North Am* 2002; 46: 767-80.
26. Jackson DC. A trial of pre-mixed nitrous oxide and oxygen in dental anaesthesia for children. *Br J Anaesth* 1967; 39: 161-4.
27. Wright GZ, Chiasson RC. The use of sedation agents by Canadian pediatric dentists. *Pediatr Dent* 1987; 9: 308-11.

28. Davis MJ. Conscious sedation practices in pediatric dentistry: a survey of members of the American Board of Pediatric Dentistry College of Diplomates. *Pediatr Dent* 1988; 10: 328-9.
29. Lawrence SM, McTigue DJ, Wilson S, Odom JG, Waggoner WF, Fields HW, Jr. Parental attitudes toward behavior management techniques used in pediatric dentistry. *Pediatr Dent* 1991; 13: 151-5.
30. Eaton JJ, McTigue DJ, Fields HW, Jr., Beck M. Attitudes of contemporary parents toward behavior management techniques used in pediatric dentistry. *Pediatr Dent* 2005; 27: 107-13.
31. Quarnstrom FC, Milgrom P, Bishop MJ, DeRouen TA. Clinical study of diffusion hypoxia after nitrous oxide analgesia. *Anesth Prog* 1991; 38: 21-3.
32. Dunn-Russell T, Adair SM, Sams DR, Russell CM, Barenie JT. Oxygen saturation and diffusion hypoxia in children following nitrous oxide sedation. *Pediatr Dent* 1993; 15: 88-92.
33. Emmanouil DE, Quock RM. Advances in understanding the actions of nitrous oxide. *Anesth Prog* 2007; 54: 9-18.
34. Primosch RE, Buzzi IM, Jerrell G. Effect of nitrous oxide-oxygen inhalation with scavenging on behavioral and physiological parameters during routine pediatric dental treatment. *Pediatr Dent* 1999; 21: 417-20.
35. Holroyd I, Roberts GJ. Inhalation sedation with nitrous oxide: a review. *Dent Update* 2000; 27: 141-2, 4, 6.
36. Guideline on appropriate use of nitrous oxide for pediatric dental patients. *Pediatr Dent* 2008; 30: 140-2.

37. Kupietzky A, Tal E, Shapira J, Ram D. Fasting state and episodes of vomiting in children receiving nitrous oxide for dental treatment. *Pediatr Dent* 2008; 30: 414-9.
38. Policy on minimizing occupational health hazards associated with nitrous oxide. *Pediatr Dent* 2008; 30: 64-5.
39. Practice guidelines for sedation and analgesia by non-anesthesiologists. A report by the American Society of Anesthesiologists Task Force on Sedation and Analgesia by Non-Anesthesiologists. *Anesthesiology* 1996; 84: 459-71.
40. Ong KS, Seymour RA, Tan JM. A prospective randomized crossover study of the preemptive analgesic effect of nitrous oxide in oral surgery. *Oral Surg Oral Med Oral Pathol Oral Radiol Endod* 2004; 98: 637-42.
41. Luhmann JD, Kennedy RM, Porter FL, Miller JP, Jaffe DM. A randomized clinical trial of continuous-flow nitrous oxide and midazolam for sedation of young children during laceration repair. *Ann Emerg Med* 2001; 37: 20-7.
42. Houpt MI, Limb R, Livingston RL. Clinical effects of nitrous oxide conscious sedation in children. *Pediatr Dent* 2004; 26: 29-36.
43. Faulks D, Hennequin M, Albecker-Grappe S, Maniere MC, Tardieu C, Berthet A, Wolikow M, Droz D, Koscielny S, Onody P. Sedation with 50% nitrous oxide/oxygen for outpatient dental treatment in individuals with intellectual disability. *Dev Med Child Neurol* 2007; 49: 621-5.
44. Annequin D, Carbajal R, Chauvin P, Gall O, Tourniaire B, Murat I. Fixed 50% nitrous oxide oxygen mixture for painful procedures: A French survey. *Pediatrics* 2000; 105: E47.

45. Faddy SC, Garlick SR. A systematic review of the safety of analgesia with 50% nitrous oxide: can lay responders use analgesic gases in the prehospital setting? *Emerg Med J* 2005; 22: 901-8.
46. Babl FE, Oakley E, Seaman C, Barnett P, Sharwood LN. High-concentration nitrous oxide for procedural sedation in children: adverse events and depth of sedation. *Pediatrics* 2008; 121: e528-32.
47. Collado V, Nicolas E, Faulks D, Hennequin M. A review of the safety of 50% nitrous oxide/oxygen in conscious sedation. *Expert Opin Drug Saf* 2007; 6: 559-71.
48. Becker DE, Rosenberg M. Nitrous oxide and the inhalation anesthetics. *Anesth Prog* 2008; 55: 124-30.
49. Constant I, Abbas M, Boucheseiche S, Laude D, Murat I. Non-invasive assessment of cardiovascular autonomic activity induced by brief exposure to 50% nitrous oxide in children. *Br J Anaesth* 2002; 88: 637-43.
50. Zacny JP, Hurst RJ, Graham L, Janiszewski DJ. Preoperative dental anxiety and mood changes during nitrous oxide inhalation. *J Am Dent Assoc* 2002; 133: 82-8.
51. Conry JP, Feigal RJ, Beniak TE. Assessment of recovery from nitrous oxide-oxygen sedation using neuropsychometry. *Anesth Prog* 1989; 36: 15-20.
52. Ayer WA, Getter L. Psychomotor responses to nitrous oxide-oxygen sedation during dental treatment. *Anesth Prog* 1974; 21: 71-3.
53. Trieger N, Newman MG, Miller JC. An objective measure of recovery. *Anesth Prog* 1969; 16: 4-7.

54. Babl FE, Puspitadewi A, Barnett P, Oakley E, Spicer M. Preprocedural fasting state and adverse events in children receiving nitrous oxide for procedural sedation and analgesia. *Pediatr Emerg Care* 2005; 21: 736-43.
55. Gan TJ, Meyer T, Apfel CC, Chung F, Davis PJ, Eubanks S, Kovac A, Philip BK, Sessler DI, Temo J, Tramer MR, Watcha M. Consensus guidelines for managing postoperative nausea and vomiting. *Anesth Analg* 2003; 97: 62-71.
56. Corah NL. Development of a dental anxiety scale. *J Dent Res* 1969; 48: 596.
57. Kindler CH, Harms C, Amsler F, Ihde-Scholl T, Scheidegger D. The visual analog scale allows effective measurement of preoperative anxiety and detection of patients' anesthetic concerns. *Anesth Analg* 2000; 90: 706-12.
58. Millar K, Jelacic M, Bonke B, Asbury AJ. Assessment of preoperative anxiety: comparison of measures in patients awaiting surgery for breast cancer. *Br J Anaesth* 1995; 74: 180-3.
59. Dixon JS, Bird HA. Reproducibility along a 10 cm vertical visual analogue scale. *Ann Rheum Dis* 1981; 40: 87-9.
60. Ahearn EP. The use of visual analog scales in mood disorders: a critical review. *J Psychiatr Res* 1997; 31: 569-79.
61. Williams VS, Morlock RJ, Feltner D. Psychometric evaluation of a visual analog scale for the assessment of anxiety. *Health Qual Life Outcomes* 2010; 8: 57.
62. Maxwell C. Sensitivity and accuracy of the visual analogue scale: a psychophysical classroom experiment. *Br J Clin Pharmacol* 1978; 6: 15-24.
63. Weimann J. Toxicity of nitrous oxide. *Best Pract Res Clin Anaesthesiol* 2003; 17: 47-61.

64. Yagiela JA. Health hazards and nitrous oxide: a time for reappraisal. *Anesth Prog* 1991; 38: 1-11.
65. Axelsson G, Ahlborg G, Jr., Bodin L. Shift work, nitrous oxide exposure, and spontaneous abortion among Swedish midwives. *Occup Environ Med* 1996; 53: 374-8.
66. Rowland AS, Baird DD, Weinberg CR, Shore DL, Shy CM, Wilcox AJ. Reduced fertility among women employed as dental assistants exposed to high levels of nitrous oxide. *N Engl J Med* 1992; 327: 993-7.
67. Brodsky JB, Cohen EN, Brown BW, Jr., Wu ML, Whitcher CE. Exposure to nitrous oxide and neurologic disease among dental professionals. *Anesth Analg* 1981; 60: 297-301.
68. Sweeney B, Bingham RM, Amos RJ, Petty AC, Cole PV. Toxicity of bone marrow in dentists exposed to nitrous oxide. *Br Med J (Clin Res Ed)* 1985; 291: 567-9.
69. Control of nitrous oxide in dental operatories. National Institute for Occupational Safety and Health. *Appl Occup Environ Hyg* 1999; 14: 218-20.
70. Szymanska J. Environmental health risk of chronic exposure to nitrous oxide in dental practice. *Ann Agric Environ Med* 2001; 8: 119-22.
71. Middendorf PJ, Jacobs DE, Smith KA, Mastro DM. Occupational exposure to nitrous oxide in dental operatories. *Anesth Prog* 1986; 33: 91-7.
72. McGlothlin JD, Jensen PA, Fischbach TJ, Hughes RT, Jones JH. Control of anesthetic gases in dental operatories. *Scand J Work Environ Health* 1992; 18: 2: 103-5.

73. Nitrous oxide in the dental office. ADA Council on Scientific Affairs; ADA Council on Dental Practice. *J Am Dent Assoc* 1997; 128: 364-5.
74. Howard WR. Nitrous oxide in the dental environment: assessing the risk, reducing the exposure. *J Am Dent Assoc* 1997; 128: 356-60.
75. Hulland SA, Freilich MM, Sandor GK. Nitrous oxide-oxygen or oral midazolam for pediatric outpatient sedation. *Oral Surg Oral Med Oral Pathol Oral Radiol Endod* 2002; 93: 643-6.
76. McCann W, Wilson S, Larsen P, Stehle B. The effects of nitrous oxide on behavior and physiological parameters during conscious sedation with a moderate dose of chloral hydrate and hydroxyzine. *Pediatr Dent* 1996; 18: 35-41.
77. Mace SE, Brown LA, Francis L, Godwin SA, Hahn SA, Howard PK, Kennedy RM, Mooney DP, Sacchetti AD, Wears RL, Clark RM. Clinical policy: Critical issues in the sedation of pediatric patients in the emergency department. *Ann Emerg Med* 2008; 51: 378-99.
78. Gayet S, Bernit E, Sati H, Veit V, Mazodier K, Schleinitz N, Kaplanski G, Harle JR. [Pain prevention with fixed 50% nitrous oxide-oxygen mixture during bone-marrow biopsy]. *Rev Med Interne* 2009; 30: 208-14.
79. German M, Pavo MR, Palacios A, Ordonez O. Use of fixed 50% nitrous oxide-oxygen mixture for lumbar punctures in pediatric patients. *Pediatr Emerg Care* 2011; 27: 244-5.
80. James MF, Manson ED, Dennett JE. Nitrous oxide analgesia and altitude. *Anaesthesia* 1982; 37: 285-8.