REFERENCES

- Bates, B.C., Z.W. Kundzewicz, S. Wu and J.P. Palutikof, Eds. (2008): Climate Change and Water. TechnicalPaper of the Intergovernmental Panel on Climate Change, IPCC Secretariat, Geneva.
- Beau Damen. BEFS Thailand: Key results and policy recommendations for future bioenergy development. Food and Agriculture Organization of the United Nations. Rome; 2010.
- Berndes, G. (2002). Bioenergy and water the implications of large-scale bioenergy production for water use and supply. Global Environmental Change-Human and Policy Dimensions, 12(4), 253-271.
- BSI (British Standards Institution). 2011. PubliclyAvailable Specification PAS 2050:
 2011 Specification for the assessment of the life cycle greenhouse gas emissions of goods and services. London, BSI.
- Carbon Trust, (2008). Code of good practice for product greenhouse gas emissions and reduction claims.
- Chapagain, A. K., Hoekstra, A. Y., Savenije, H. H. G., & Gautam, R. (2006). The water footprint of cotton consumption: An assessment of the impact of worldwide consumption of cotton products on the water resources in the cotton producing countries. Ecological Economics, 60(1), 186-203.
- Contreras, A. M., Rosa, E., Perez, M., Van Langenhove, H., & Dewulf, J. (2009).
 Comparative Life Cycle Assessment of four alternatives for using by-products of cane sugar production. Journal of Cleaner Production, 17(8), 772-779.
- Dalla Marta, A., Mancini, M., Natali, F., Orlando, F., & Orlandini, S. (2012). From water to bioethanol: The impact of climate variability on the water footprint. Journal of Hydrology, 444, 180-186.
- EPLCA, 2007. Carbon Footprint-What It Is and How to Measure It. European Platform on Life Cycle Assessment, European Commission.

- Ercin, A.E., Hoekstra, A.Y. (2012). Carbon and Water Footprints; concepts, Methodologies and Policy Response. The United Nations Educational, Scientific and Cultural Organization. France.
- FAO. 1986. Yield response to water. By: J. Doorenbos and A.H. Kassam. FAO Irrigation and Drainage Paper 33. Rome, Italy.
- FAO. 1998. Crop evapotranspiration: Guidelines for computing crop water requirements. By: Richard Allen, Luis Pereira, Dirk Raes and Martin Smith. FAO Irrigation and Drainage Paper 56. Rome, Italy.
- Food and Agriculture Organization (2009). CROPWAT 8.0 Model. Last visit in February 2013. Available in http://www.fao.org/nr/water /infores_databases_cropwat.html.
- Gerbens-Leenes, P.W., and Hoekstra, A. Y. (2012). The water footprint of sweeteners and bio-ethanol. Environment International, 40, 202-211.
- Gerbens-Leenes, P. W., Hoekstra, A. Y., & van der Meer, T. (2009). The water footprint of energy from biomass: A quantitative assessment and consequences of an increasing share of bio-energy in energy supply. Ecological Economics, 68(4), 1052-1060.
- Gerbens-Leenes, P. W., Van Lienden, A. R., Hoekstra, A. Y., & van der Meer, T. H. (2012). Biofuel scenarios in a water perspective: The global blue and green water footprint of road transport in 2030. Global Environmental Change-Human and Policy Dimensions, 22(3), 764-775.
- Hannah Kosow and Robert Gaβner Methods of future and scenario analysis;Overview, assessment, and selection criteria (2008) www.diegdi.de/CMS...nsf/.../Studies%2039.2008.pdf
- Hoekstra, A. Y. (ed) (2003) 'Virtual water trade: Proceedings of the International Expert Meeting on Virtual Water Trade', 12–13 December 2002, Value of Water Research Report Series No 12, UNESCO-IHE, Delft, Netherlands, www.waterfootprint.org/Reports/Report12.pdf.
- Hoekstra, A.Y., Chapagain, A.K., (2007). Water footprint of nations: water use buy people as a function of their consumption pattern. Water Resour. Manage 21, 35-48.

- Hoekstra, A.Y., Chapagain, A.K., (2008). Globalization of water. Sharing the Planet's Freshwater Resources. Blackwell, Oxford, UK.
- Hoekstra, A.Y., Chapagain, A.K., Aldaya, M.M., Mekonnen, M.M. (2011). The water footprint assessment manual: setting the global standard. Water Footprint Network, the Netherlands.
- How will climate change impact on fresh water security?. Last visit April 2013. http://www.guardian.co.uk/ environment/2012/nov/30/climate-change-water.
- IPCC. (2006). In: Eggleston, H.S., Buendia, L., Miwa, Ngara, T., Tanabe, K. (Eds.),2006 IPCC Guidelines for National Greenhouse Gas Inventories, The National Greenhouse Gas Inventories Programme. IGES, Japan.
- Iribarren, D., Hospido, A., Moreira, M. T., & Feijoo, G. (2010). Carbon footprint of canned mussels from a business-to-consumer approach. A starting point for mussel processors and policy makers. Environmental Science & Policy, 13(6), 509-521.
- ISO, 2006a. ISO 14040:2006. Environmental Management Life Cycle Assessment -Principles and Framework. International Organization for Standardization.
- ISO, 2006b. ISO 14044:2006. Environmental Management Life Cycle Assessment Requirements and guidelines. International Organization for Standardization.
- ISO, 2006c. ISO 14064-1:2006. Greenhouse Gases. Part 1. Specification with Guidance at the Organization Level for Quantification and Reporting of Greenhouse Gas Emissions and Removals. International Organization for Standardization.
- ISO, 2006d. ISO 14025:2006. Environmental management Type III environmental declarations – Principles and procedures.
- Japan, (2009). Japanese Technical Specification "General principles for the assessment and labeling of Carbon Footprint of products".
- Kongboon, R., Samputtagul, S. (2012). The water footprint of sugarcane and cassava in northern Thailand. Procedia Social and Behavioral Sciences, 40, 451 460.
- Kongboon, R., Samputtagul, S. (2012). Water Footprint of Bioethanol Production from Sugarcane in Thailand. Journal of Environment and Earth Science, 2(11), 61-67.

- Land Development Department. Soil series data. Last visit in March 2012. Available in http://giswebldd.ldd. go.th/.
- Leenes, G., Hoekstra, A. Y. and Meer, H. V. D. 2009. The Water footprint of energy from biomass: a quantitative assessment and consequences of an increasing share of bio-energy in energy supply. Ecological economics, (68), 1052-1060 pp.
- Liu, C., Kroeze, C., Hoekstra, A. Y., & Gerbens-Leenes, W. (2012). Past and future trends in grey water footprints of anthropogenic nitrogen and phosphorus inputs to major world rivers. Ecological Indicators, 18, 42-49.
- Liu, J. G., Williams, J. R., Zehnder, A. J. B. and Yang, H. (2007) 'GEPIC: Modelling wheat yield and crop water productivity with high resolution on a global scale', Agricultural Systems, vol. 94, no 2, pp 478–493.
- Luo, L., van der Voet, E., & Huppes, G. (2009). Life cycle assessment and life cycle costing of bioethanol from sugarcane in Brazil. Renewable & Sustainable Energy Reviews, 13(6-7), 1613-1619.
- Mekonnen, M. M., & Hoekstra, A. Y. (2011). The green, blue and grey water footprint of crops and derived crop products. Hydrology and Earth System Sciences, 15(5), 1577-1600.
- Natural Resources Defense Council. Climate change, water, and risk: Current water demands are not sustainable. Last visit in April 2013. Available in http://www.nrdc.org/globalWarming/watersustain- ability/.
- O, A. (2010). Water use impact of ethanol at a gasoline substitution ratio of 5% from cassava in Nigeria. Biomass and Bioenergy, 34(7), 985-992.
- Office of Agricultural Economics. Agricultural statistic of Thailand. Last visit in September 2010. Available in http://www.oae.go.th/download/download _journal/yearbook2552.pdf.
- Ometto, A. R., Hauschild, M. Z., & Roma, W. N. L. (2009). Lifecycle assessment of fuel ethanol from sugarcane in Brazil. International Journal of Life Cycle Assessment, 14(3), 236-247.

Pollution control department. Water Quality standard. Last visit in Februarys 2013. Available in http://www.pcd.go.th/info_serv/reg_std_water05.html.

- Renouf, M. A., Pagan, R. J., & Wegener, M. K. (2011). Life cycle assessment of Australian sugarcane products with a focus on cane processing. International Journal of Life Cycle Assessment, 16(2), 125-137.
- Richard G. Allen, Luis S. Pereira, Dirk Raes, Martin Smith. (1998). FAO Irrigation and Drainage Paper No. 56 Crop evapotranspiration: guidelines for computing crop water requirements. Food and Agriculture Organization of the United Nations. Rome.
- Royal Irrigation Department. Crop coefficient. Last visit in August 2011. Available in http://water.rid.go.th/hwm/ cropwater/CWRdata/Kc/kc_th.pdf.
- SETAC Europe LCA Steering Committee, 2008. Standardisation efforts to measure greenhouse gases and 'carbon footprinting' for products. The International Journal of Life Cycle Assessment 13 (2), 87–88.
- Sevigne, E., Gasol, C. M., Brun, F., Rovira, L., Pages, J. M., Camps, F., et al. (2011).
 Water and energy consumption of Populus spp. bioenergy systems: A case study in Southern Europe. Renewable & Sustainable Energy Reviews, 15(2), 1133-1140.
- Scientific Applications International Corporation (SAIC). Life Cycle Assessment: Principles and Practice. Last visit in May 2010. Available in http://www.epa.gov/nrmrl/std/lca/lca.html.
- Siebert, S., Doll, P., (2010). Quantifying blue and green virtual water contents in global crop production as well as potential production losses without irrigation. Science 319(5867), 1238-1240.
- Thailand Greenhouse Gas Management Organization (Public Organization, TGO). Emission factor, Thailand. Last visit January 2010. Available in: http://thaicarbonlabel.tgo.or.th/download/Emission_Factor.pdf.
- Thai Meteorological Department. Climate Data. Last visit in June 2011. Available online in http://www.tmd.go.th/.
- The National Technical Committee on Product Carbon Footprint (Thailand), 2010. The National Guideline on Product Carbon Footprint. Amarin Publishing Bangkok, Thailand.
- UN-Water Statistics. Last visit in February 2013. Available in www.unwater.org /statistcs_res.html.

- Williams, J. R., Jones, C.A., Kiniry, J. R. and Spanel, D. A. (1989) 'The EPIC crop growth-model', Transactions of the ASAE, vol 32, no 2, pp497–511.
- Williams, J. R. (1995) 'The EPIC model', in V. P. Singh (ed) Computer Models of Watershed Hydrology, Water Resources Publisher, CO, pp909–1000.
- W. Michael, H. Jeongwoo, D. Jennifer B, C. Hao, E. Amgad. (2012). Well-to-wheels energy use and greenhouse gas emissions of ethanol from corn, sugarcane and cellulosic biomass for US use. Environment Research Letters. 7, 1-13.

Yuttitham, M., Gheewala, S. H., & Chidthaisong, A. (2011). Carbon footprint of sugar produced from sugarcane in eastern Thailand. Journal of Cleaner Production, 19(17-18), 2119-2127.

ลิ<mark>ปสิทธิ์มหาวิทยาลัยเชียงใหม่</mark> Copyright[©] by Chiang Mai University All rights reserved