

TABLE OF CONTENTS

	Page
Acknowledgement	III
Abstract (Thai)	IV
Abstract (English)	VII
List of Table	XII
List of Figure	XV
Chapter 1 Introduction	1
Chapter 2 Literature reviews	5
Chapter 3 Materials and Methods	26
Chapter 4 Results	39
Chapter 5 Discussion	141
Chapter 6 Conclusion	156
References	159
Appendices	176
Appendix A	177
Appendix B	184
Curriculum Vitae	187

LIST OF TABLES

Table		Page
3-1	Description of collecting site where snail collections were performed during April 2010 – June 2012 from 12 provinces, northern Thailand.	31
3-2	List of material and sequences of mtCOI used for generated phylogenetic relationship of heterophyid trematodes	38
4-1	The total number of each snail species which collecting in 12 provinces of northern Thailand	56
4-2	Biodiversity indices of freshwater snail calculating form northern Thailand	59
4-3	Composition of the snail species and number of cercarial infections in Chiang Mai province	99
4-4	Composition of the snail species and number of cercarial infections in Chiang Rai province	104
4-5	Composition of the snail species and number of cercarial infections in Mae Hong Son province	105
4-6	Composition of the snail species and number of cercarial infections in Lampang province	106
4-7	Composition of the snail species and number of cercarial infections in Lamphun province	108

LIST OF TABLES

Table		Page
4-8	Composition of the snail species and number of cercarial infections in Nan province	109
4-9	Composition of the snail species and number of cercarial infections in Phayao province	110
4-10	Composition of the snail species and number of cercarial infections in Phrae province	111
4-11	Composition of the snail species and number of cercarial infections in Phitsanulok province	112
4-12	Composition of the snail species and number of cercarial infections in Sukhothai province	113
4-13	Composition of the snail species and number of cercarial infections in Tak province	114
4-14	Composition of the snail species and number of cercarial infections in Uttaradit province	115
4-15	List of the sequences of mtCOI producing significant data of <i>H. taichui</i> from NCBI databases	122
4-16	List of the sequences of mtCOI producing significant data of <i>H. pumilio</i> from NCBI databases	125

LIST OF TABLES

Table		Page
4-17	Specification of the <i>H. taichui</i> and <i>H. pumilio</i> specific primers designed based on the partial sequence of mtCOI	126
4-18	Summaries of positive results for the molecular identification using multiplex PCR	136

LIST OF FIGURES

Figure		Page
2-1	General pattern of heterophyid trematodes life cycle by using thiarid snails and cyprinoid fishes as a first and second intermediate host	12
2-2	Type of cercaria	13
2-3	Important features of a dextral, or right-turned, spire of gastropod shell	21
3-1	Map of the 64 sampling sites of northern Thailand	29
4-1	Morphological characteristic of <i>Haplorchis taichui</i>	41
4-2	Morphological characteristic of <i>Haplorchis pumilio</i>	41
4-3	Morphological characteristic of <i>Centrocestus caninus</i>	44
4-4	Morphological characteristic of <i>Stellantchasmus falcatus</i>	44
4-5	Morphological characteristic of <i>Haplorchoides</i> sp.	47
4-6	Morphological characteristic of <i>Ganeo tigrinus</i>	47
4-7	Morphological characteristic of <i>Prostorchigenes majeedi</i>	50
4-8	Morphological characteristic of <i>Fasciola gigantica</i>	50
4-9	Morphological characteristic of <i>Fischoederius. elongatus</i>	53
4-10	Morphological characteristic of <i>Orthocoelium streptocoelium</i>	53
4-11	The proportion of freshwater snail families were collected from 12 provinces of northern Thailand	55

LIST OF FIGURES (CONTINUED)

Figure		Page
4-12	Shell characteristics of <i>Adamietta housei</i>	60
4-13	Shell characteristics of <i>Bithynia siamensis siamensis</i>	61
4-14	Shell characteristics of <i>Bithynia funiculata</i>	62
4-15	Shell characteristics of <i>Brotia costula costula</i>	63
4-16	Shell characteristics of <i>Brotia citrina</i>	64
4-17	Shell characteristics of <i>Brotia wykoffi</i>	65
4-18	Shell characteristics of <i>Clea helana</i>	66
4-19	Shell characteristics of <i>Eyriesia eyriesi</i>	67
4-20	Shell characteristics of <i>Filopaludina doliaris</i>	68
4-21	Shell characteristics of <i>Filopaludina martensi martensi</i>	69
4-22	Shell characteristics of <i>Filopaludina sumatrensis polygramma</i>	70
4-23	Shell characteristics of <i>Filopaludina sumatrensis speciosa</i>	71
4-24	Shell characteristics of <i>Indoplanorbis exustus</i>	72
4-25	Shell characteristics of <i>Lymnaea auricularia rubiginosa</i>	73
4-26	Shell characteristics of <i>Melanoides tuberculata</i>	74
4-27	Shell characteristics of <i>Mekongia swainsoni</i>	75
4-28	Shell characteristics of <i>Paludomus siamensis</i>	76
4-29	Shell characteristics of <i>Pomacea canaliculata</i>	77
4-30	Shell characteristics of <i>Tarebia granifera</i>	78

LIST OF FIGURES (CONTINUED)

Figure		Page
4-31	Shell characteristics of <i>Thiara scabra</i>	79
4-32	Shell characteristics of <i>Sinotaia mandahlbarthi</i>	80
4-33	The proportion of cercarial infected of each snail families	82
4-34	Cercarial infections of freshwater snails in northern Thailand	83
4-35	Morphological characteristic of distome cercaria	85
4-36	Morphological characteristic of cercariae	86
4-37	Morphological characteristic of gymnocephalous cercaria	87
4-38	Morphological characteristic of <i>Transversotrema</i> cercaria	88
4-39	Morphological characteristic of strigea cercaria	89
4-40	Morphological characteristic of megalurous cercaria	90
4-41	Morphological characteristic of monostome cercaria	91
4-42	Morphological characteristic of ophthalmoxiphidiocercaria	92
4-43	Morphological characteristic of parapleurolophocercous cercaria	93
	Type I	
4-44	Morphological characteristic of parapleurolophocercous cercaria	94
	Type II	
4-45	Morphological characteristic of parapleurolophocercous cercaria	95
	Type III	
4-46	Morphological characteristic of pleurolophocercous cercaria	96
4-47	Morphological characteristic of xiphidiocercaria	97

LIST OF FIGURES (CONTINUED)

Figure		Page
4-48	Morphological characteristic of virgulate cercaria	98
4-49	Overview of distribution of cercarial infected in freshwater snails in 12 provinces of northern Thailand	116
4-50	Distribution of each type of cercaria, which infected in freshwater snails in 12 provinces of northern Thailand	117
4-51	The mtCOI product of ten species trematodes	121
4-52	The multiple alignment of ten species of trematodes	127
4-53	PCR product of <i>H. taichui</i> and <i>H. pumilio</i> specific primers by various annealing temperatures	128
4-54	Sensitivity tests of <i>Haplorchis taichui</i> and <i>H. pumilio</i> specific primers	130
4-55	Specificity tests of <i>H. taichui</i> specific primer by various species of trematode and freshwater snails	131
4-56	Specificity tests of <i>H. pumilio</i> specific primer by various species of trematode and freshwater snails	131
4-57	Multiplex PCR detection the <i>H. taichui</i> and <i>H. pumilio</i>	132
4-58	Molecular identification by multiplex PCR reaction	133

LIST OF FIGURES (CONTINUED)

Figure		Page
4-59	The rooted phylogeny from partial mtCOI sequences of 17 trematode species using NJ method based on the Kimura two-parameter model	140