

## TABLE OF CONTENTS

	<b>Page</b>
ACKNOWLEDGEMENT	iii
ABSTRACT (ENGLISH)	v
ABSTRACT (THAI)	vii
LIST OF TABLES	xviii
LIST OF FIGURES	xx
ABBREVIATIONS AND SYMBOLS	xxiii
CHAPTER 1 INTRODUCTION	1
1.1 Statement and significance of the problems	1
1.2 Objective	3
1.3 Scope of the study	4
1.4 Literature reviews	5
1.4.1 Thai flowers	5
1.4.1.1 Introduction	5
1.4.1.2 <i>Gardenia jasminoides</i>	6
1.4.1.3 <i>Millingtonia hortensis</i>	8
1.4.1.4 <i>Mimusops elengi</i>	17
1.4.2 Extraction processes	22

1.4.2.1	Enfleurage	22
1.4.2.2	Solvent extraction	22
1.4.2.3	Supercritical carbon dioxide fluid extraction (scCO <sub>2</sub> )	23
1.4.3	Phytochemical determination	25
1.4.3.1	Alkaloids	25
1.4.3.2	Flavonoids	25
1.4.3.3	Saponins	26
1.4.3.4	Tannins	27
1.4.3.5	Reducing sugars	28
1.4.3.6	High performance liquid chromatography(HPLC)	30
1.4.4	Nanovesicles	31
1.4.4.1	Classification of nanovesicles	31
	A. Liposomes	32
	A.1 Liposomes in the basic sciences	34
	A.2 Liposomes in medicines	34
	A.3 Liposomes in cosmetics	36
	A.4 Liposomes in bioengineering	36
	A.5 Liposomes in agro-food industries	37
	B. Niosomes	38
	C. Transfersomes	40

D. Ethosomes	41
1.4.4.2 Preparation of nanovesicles	42
A. Ether injection method	42
B. Hand shaken method	43
C. Reverse phase evaporation method	44
D. Transmembrane pH gradient (inside acidic) drug uptake process (remote loading)	44
E. Formation of proliposomes and liposomes from proliposomes	45
F. The bubble method	45
G. High pressure homogenization and Microfluidization	46
H. Supercritical carbon dioxide fluid technique (scCO <sub>2</sub> )	46
1.4.4.3 Physicochemical characteristics of nanovesicles	51
A. Vesicular size	52
B. Morphology	52
C. Charge	53
D. Entrapment efficiency	53
1.4.5 Biological assays	54

1.4.5.1	<i>In vivo</i> skin irritation assay in rabbits	54
1.4.5.2	<i>In vitro</i> assays for cosmetic uses	56
A.	DPPH radical scavenging activity	56
B.	Ferrous metal chelating activity assay	56
C.	Tyrosinase inhibition	57
D.	Lipid peroxidation inhibition assay	58
E.	Cell proliferation activity	59
F.	Gelatinolytic activity on MMP-2 inhibition (zymography)	61
CHAPTER 2 MATERIALS AND METHODS		66
2.1	Materials and equipments	66
2.1.1	Chemicals	66
2.1.2	Cell lines	67
2.1.3	Animals	67
2.1.4	Equipments	67
2.2	Methods	68
2.2.1	Biological activities of the Thai flower extracts	68
2.2.1.1	Collection of the raw materials	69
2.2.1.2	Preparation of the flower extracts	69
A.	Flower samples	69
B.	Maceration process	69
C.	Supercritical carbon dioxide fluid	70

(scCO <sub>2</sub> ) extraction	
2.2.1.3 Determination of the characteristics and phytochemicals of the extracts	70
2.2.1.4 Chemical identification of the extracts by GC/HPLC	71
2.2.1.5 Biological activity determination of the Extracts	73
A. DPPH radical scavenging activity	73
B. Metal chelating activity	74
C. Tyrosinase inhibition assay	75
D. Cell proliferation activity on aged human skin fibroblasts by SRB assay	76
E. Gelatinolytic activity	77
2.2.1.6 Selection of the extracts	78
2.2.2 Characteristics and biological activities of Thai flower extracts loaded in niosomes	78
2.2.2.1 Preparation of niosomal formulations loaded with the extracts / maximum loading determination	78
2.2.2.2 Morphology, vesicular size and zeta potential determination	79
2.2.2.3 Biological activities of niosomes loaded	79

with the extracts	
A. Cytotoxicity on human skin fibroblast by SRB assay	79
B. Lipid peroxidation activity	80
C. Metal ion chelating activity	81
D. Gelatinolytic activity (zymography) on MMP-2 inhibition	82
2.2.2.4 Physical stability of the flower extracts loaded and not loaded in niosomes	83
2.2.3 Irritation study on rabbits' skin of the extracts and niosomes loaded with the extracts	83
2.2.3.1 Preparation of the animals	83
2.2.3.2 Method	84
2.2.4 Cost estimation	85
2.2.4.1 The selected extracts	85
2.2.4.2 The selected niosmes loaded with the flower extract	85
CHAPTER 3 RESULTS AND DISCUSSION	86
3.1 Biological activities of the Thai flower extracts	86
3.1.1 Physical characteristics of the extracts	86
3.1.2 Biological activities of the extracts	87
3.1.2.1 DPPH radical scavenging activity	87

3.1.2.2 Tyrosinase inhibition activity	89
3.1.2.3 Cell proliferation activity	90
3.1.2.4 Zymography of MMP-2 inhibition	90
3.2 Characteristics and biological activity of Thai flower extracts loaded in niosomes	91
3.2.1 Physical stability of the flower extracts loaded and non-loaded in niosomes	91
3.2.2 Cytotoxicity of the extracts loaded and non-loaded in niosomes	97
3.2.3 Biological activities of the flower extracts loaded and non-loaded in niosomes	98
3.2.3.1 Antioxidant activities	98
3.2.3.2 Gelatinolytic activity of the flower extracts loaded and non-loaded in niosomes	99
3.3 Skin irritation on rabbits' skin of the flower extracts loaded and non-loaded in niosomes	105
3.4 Cost estimation	107
3.4.1 The selected extracts	107
3.4.2 The selected niosomes loaded with the flower extracts	107

REFERENCES	112
APPENDICES	132
Appendix A	133
Appendix B	137
Appendix C	152
Appendix D	154
Appendix E	156
CURRICULUM VITAE	160



## LIST OF TABLES

<b>Table</b>		<b>Page</b>
1	The chemical compositions found in the extract of <i>G. jasminoides</i> flowers	9
2	The chemical compositions found in the extract of <i>M hortensis</i> flowers	16
3	The chemical compositions found in the extract of <i>M. elengi</i> flowers	19
4	Classifications of the liposome based on their structural properties and the preparation methods	35
5	Liposomal cosmetic formulations currently in the market	37
6	The vesicles formation ability, HLB values and chemical structure of some non-ionic surfactants	39
7	The critical temperature, pressure and density of substances common used as supercritical fluids	47
8	The advantages and disadvantages of different methods for the separation of the vesicles from the untrapped drug	55
9	Body distribution of collagen types	63
10	Types of MMPs	63

11	Comparison of appearance, percentage yields, antioxidative, tyrosinase inhibition activities and the percentages of cell growth on human skin fibroblasts (15 <sup>th</sup> passage) of the three flower extracts prepared by the two non-heated methods (scCO <sub>2</sub> and hexane maceration)	88
12	Stability of the extract using $\gamma$ linoleic acid percent remaining as the marker determination by HPLC	97
13	The antioxidant activities including DPPH radical scavenging (SC <sub>50</sub> ), metal chelating (MC <sub>50</sub> ) and lipid peroxidation inhibition (LC <sub>50</sub> ) of the extracts loaded and non-loaded in niosomes	101
14	Rabbit skin erythema measured by Mexameter of various samples	106
15	The estimated cost of the extracts and the extracts loaded in niosomes	108

## LIST OF FIGURES


<b>Figure</b>		<b>Page</b>
1	The flower of <i>G. jasminoides</i>	6
2	The flowers of <i>M. hortensis</i>	8
3	The flowers of <i>M. elengi</i>	17
4	Chemical structures of some common alkaloids	26
5	Chemical structures of flavonoid classes	27
6	Chemical structures of saponins	28
7	Chemical structures of condensed (proanthocyanins) and hydrolysable (corilagin and geraniin) tannins	29
8	Chemical structures of reducing forms of glucose (left) and fructose (right)	31
9	The critical packing parameter (CPP) of lipids and their structure form (DGDG: digalactosyl diglyceride and diglucosyl diglyceride; MGDG: monogalactosyl diglyceride and monoglucosyl diglyceride)	33
10	Pressure-temperature phase diagram and supercritical fluid region	47
11	Reaction of the DPPH radical in the presence of the antioxidant during the DPPH assay	57
12	Melanogenesis pathway	58

- 13 The reaction of lipid peroxidation in ferric thiocyanate assay 59
- 14 Structure of MTT and their corresponding reaction products 60
- 15 Gelatin zymography 65
- 16 The percentage yields of *M. elengi*, *M. moso* and *G. jasminoides* 87  
by Hexane Maceration and scCO<sub>2</sub> with the various concentrations  
of ethanol
- 17 MMP-2 inhibition activity of the flower extracts from *M. elengi* 91  
(Bullet Wood) prepared by scCO<sub>2</sub> with 33% (w/v) ethanol, *M.*  
*hortensis* (Cork Tree) by scCO<sub>2</sub> with 20% (w/v) ethanol and *G.*  
*jasminoides* (Gardenia) by scCO<sub>2</sub> with 33% (w/v) ethanol  
compared to vitamin C and E. (A) zymograms and (B) the  
percentages of MMP-2 inhibition.  $MMP-2 \text{ inhibition } (\%) = 100 -$   
 $[(MMP-2 \text{ content}_{\text{sample}} / MMP-2 \text{ content}_{\text{control}}) \times 100]$ . The density  
was the total intensity of all the pixels in the volume divided by the  
area of the volume
- 18 Physical appearances at initial of (A) the blank niosomes, the 93  
niosomes loaded with the extracts of (B) *Mimusops elengi* (C)  
*Gardenia jasminoides* and (D) *Millingtonia hortensis*
- 19 Morphology of niosomes loaded with (A) *Mimusops elengi*, (B) 94  
*Millingtonia hortensis* and (C) *Gardenia jasminoides* extract. The  
niosomal dispersion stained with 1% methylene blue was observed  
under 100x optical microscope

20	Vesicular sizes of niosomes loaded with <i>Mimusops elengi</i> (A), <i>Millingtonia hortensis</i> (B) and <i>Gardenia jasminoides</i> (C) extracts kept at 4±2, 27±2 and 45±2°C for 3 months	95
21	Zeta potential values of niosomes loaded with <i>Mimusops elengi</i> (A), <i>Millingtonia hortensis</i> (B) and <i>Gardenia jasminoides</i> (C) extracts kept at 4±2, 27±2 and 45±2°C for 3 months	96
22	Percentages of cell viability of human skin fibroblasts treated with the flower extracts loaded and non-loaded in niosomes. Vitamin C was used as a standard	100
23	Zymograms of MMP-2 inhibition of the non-loaded and loaded flower extracts in niosomes in comparing to the negative (untreated cells) and positive (vitamin C) control systems at various concentrations (0.01, 0.1 and 1 mg/ml)	102
24	The percentages of MMP-2 inhibition of the <i>Mimusops elengi</i> (A), <i>Millingtonia hortensis</i> (B) and <i>Gardenia jasminoides</i> (C) loaded and non-loaded in niosomes	103

## ABBREVIATIONS AND SYMBOLS

BHT	butylated hydroxytoluene
D-MEM	dulbecco's modified eagle's medium
DMSO	dimethyl sulfoxide
DLS	dynamic light scattering
EDTA	ethylenediaminetetraacetic acid
FBS	fetal bovine serum
FeCl <sub>2</sub>	ferrous chloride
h	hour
HPLC	high performance liquid chromatography
IC <sub>50</sub>	concentration providing 50% inhibition
MEM	eagle's minimal essential medium
MTT	3-(4,5-dimethyl-thiazol-2-yl)-2,5-diphenyl-2 <i>H</i> -tetrazolium bromide
mg	milligram
ml	milliliter
mM	millimolar
nm	nanometer
PB	phosphate-buffered
PBS	phosphate-buffered saline
rpm	revolutions per minute
s	second
SC <sub>50</sub>	concentration providing 50% scavenging

The background of the page features a large, faint watermark of the Chiang Mai University logo. The logo is circular and contains a central figure of an elephant standing under a traditional Thai umbrella. The text 'CHIANG MAI UNIVERSITY 1964' is written around the bottom inner edge of the circle, and Thai script is written around the top inner edge. There are also decorative floral motifs on the left and right sides of the circle.

SRB	sulphorodamine B
TEM	transmission electron microscopy
TLC	thin layer chromatography
Tween 61	polyoxyethylene sorbitan monostearate
$\lambda$	wavelength in nanometer
$\mu\text{g}$	microgram
$\mu\text{L}$	microliter
$^{\circ}\text{C}$	celcius degree

ลิขสิทธิ์มหาวิทยาลัยเชียงใหม่

Copyright© by Chiang Mai University  
All rights reserved