

## TABLE OF CONTENTS

|   | <b>Page</b> |
|---|-------------|
| ACKNOWLEDGEMENTS                              | iii         |
| ABSTRACT (ENGLISH)                            | v           |
| ABSTRACT (THAI)                               | ix          |
| TABLE OF CONTENT                              | xii         |
| LIST OF TABLES                                | xvi         |
| LIST OF FIGURES                               | xxi         |
| ABBREVIATIONS AND SYMBOLS                     | xxvii       |
| <br>CHAPTER I: INTRODUCTION                   | <br>1       |
| 1.1 Statement and significance of the problem | 1           |
| 1.2 Purposes of the study                     | 4           |
| 1.3 Education advantages                      | 5           |
| <br>CHAPTER II: LITERATURE REVIEW             | <br>6       |
| 2.1 Nanodelivery                              | 6           |
| 2.2 Microemulsion                             | 6           |
| 2.2.1 Definition                              | 6           |
| 2.2.2 Components of microemulsion             | 8           |
| 2.2.2.1 Oil phase                             | 8           |
| 2.2.2.2 Surfactant                            | 9           |

|  |           |
|--|-----------|
| 2.2.2.3 Cosurfactant   | 10        |
| 2.2.2.4 Aqueous phase  | 12        |
| 2.3 Phase behavior   | 12        |
| 2.4 Topical microemulsion and recent researches on microemulsion         | 13        |
| 2.5 Antityrosinase activity  | 16        |
| 2.6 Antioxidant activity   | 18        |
| <b>CHAPTER III: MATERIALS AND METHODS</b>                                | <b>21</b> |
| 3.1 Instruments  | 21        |
| 3.2 Chemicals  | 23        |
| 3.3 Plants materials   | 26        |
| 3.4 Extraction of essential oil  | 28        |
| 3.5 Study of outer appearance and density of oil                         | 31        |
| 3.6 Study of chemical components by gas chromatography/mass spectrometry | 31        |
| 3.7 Study of the biological activities of the oils                       | 32        |
| 3.7.1 Determination of antioxidant activity                              | 32        |
| 3.7.1.1 Ferric Reducing Antioxidant Power (FRAP)                         | 32        |
| 3.7.1.2 ABTS Method  | 33        |
| 3.7.2 Determination of tyrosinase inhibitory activity                    | 33        |
| 3.8 Solubility study   | 36        |
| 3.9 Refractive index and surface tension of oil                          | 38        |
| 3.10 Cytotoxicity tests  | 40        |
| 3.11 The study of phase diagram  | 40        |
| 3.11.1 Preliminary study for phase diagram construction                  | 41        |

|  |           |
|--|-----------|
| 3.11.2 Pseudoternary phase diagram construction                          | 47        |
| 3.12 Microemulsion base and drug loaded microemulsion preparation        | 50        |
| 3.13 Drug release study  | 52        |
| 3.14 Stability study of the microemulsion                                | 53        |
| <b>CHAPTER IV: RESULTS AND DISCUSSION</b>                                | <b>55</b> |
| 4.1 Extraction of essential oils   | 55        |
| 4.2 Study of outer appearance and density of oil                         | 58        |
| 4.3 Study of chemical components by gas chromatography/mass spectrometry | 63        |
| 4.4 The study of the biological activities of the oils                   | 116       |
| 4.4.1 Antioxidant activity   | 116       |
| 4.4.1.1 Ferric Reducing Antioxidant Power (FRAP)                         | 116       |
| 4.4.1.2 ABTS Method  | 118       |
| 4.4.2 Tyrosinase inhibitory activity                                     | 121       |
| 4.5 Solubility test of the oil   | 124       |
| 4.6 Refractive index and surface tension of the oil                      | 126       |
| 4.7 Cytotoxicity tests   | 130       |
| 4.8 Study of phase diagram   | 132       |
| 4.8.1 Preliminary study for phase diagram construction                   | 132       |
| 4.8.2 Phase diagram construction   | 152       |
| 4.8.2.1 Effect of surfactant type  | 152       |
| 4.8.2.2 Effect of type and proportion of cosurfactant                    | 153       |
| 4.8.2.3 Effect of type and concentration of electrolyte                  | 158       |
| 4.8.2.1 Effect of pH   | 158       |

|                       |  |     |
|-----------------------|--|-----|
| 4.9                   | Microemulsion base and drug loaded microemulsion preparation | 172 |
| 4.10                  | Stability study of microemulsion                             | 181 |
| 4.10.1                | Stability of the formulations without drug                   | 181 |
| 4.10.2                | Stability of clotrimazole loaded formulations                | 181 |
| 4.11                  | Drug release study   | 200 |
| CHAPTER V: CONCLUSION |  | 207 |
| REFERENCES            |  | 211 |
| APPENDICES            |  | 238 |
| APPENDIX I            |  | 239 |
| APPENDIX II           |  | 241 |
| CURRICULUM VITAE      |  | 242 |

## LIST OF TABLES

| Table |  | page |
|-------|--|------|
| 3.1   | Plants for oil extraction used in this research  | 26   |
| 3.2   | The part of the plant used in extraction of oil  | 28   |
| 3.3   | Polarity, dielectric constant and solubility parameter of the solvents used in this experiment | 37   |
| 3.4   | The amount of substances used in a system of Tween 20: ethanol = 1:1                           | 42   |
| 3.5   | The amount of substances used in a system of Tween 20: ethanol = 2:1                           | 43   |
| 3.6   | The amount of substances used in a system of Tween 20: ethanol = 3:1                           | 44   |
| 3.7   | The amount of substances used in a system of Tween 20: ethanol = 9:1                           | 45   |
| 3.8   | The amount of substances used in a system of Tween 80: ethanol = 1:1                           | 46   |
| 3.9   | The amount of substances used in a system of Tween 80: ethanol = 2:1                           | 47   |
| 3.10  | The amount of substances used in a system of Tween 80: ethanol = 3:1                           | 48   |
| 3.11  | The amount of substances used in a system of Tween 80: ethanol = 9:1                           | 49   |
| 3.12  | Microemulsion systems used in this study   | 52   |
| 4.1   | The amount oil extracted as percent yield (% v/w) relative to fresh weight                     | 56   |
| 4.2   | The explanation of the letter in Figure 4.1  | 59   |
| 4.3   | The density of oil used in this experiment   | 61   |
| 4.4   | Chemical components of essential oil from <i>Apium graveolens</i> Linn.                        | 64   |
| 4.5   | Chemical components of essential oil from <i>Anethum graveolens</i> Linn.                      | 65   |

|      |  |     |
|------|--|-----|
| 4.6  | Chemical components of essential oil from <i>Centella asiatica</i> Urban.        | 67  |
| 4.7  | Chemical components of essential oil from <i>Coriandrum sativum</i> Linn.        | 69  |
| 4.8  | Chemical components of essential oil from <i>Eryngium foetidum</i> Linn.         | 71  |
| 4.9  | Chemical components of essential oil from <i>Polyscias fruticosa</i> Harms.      | 72  |
| 4.10 | Chemical components of essential oil from <i>Eupatorium odoratum</i> Linn.       | 74  |
| 4.11 | Chemical components of essential oil from <i>Spilanthes acmella</i> Murr.        | 76  |
| 4.12 | Chemical components of essential oil from <i>Cymbopogon citratus</i> Stapf.      | 78  |
| 4.13 | Chemical components of essential oil from <i>Coleus amboinicus</i> Lour.         | 80  |
| 4.14 | Chemical components of essential oil from <i>Melissa officinalis</i> Linn.       | 82  |
| 4.15 | Chemical components of essential oil from <i>Ocimum basilicum</i> Linn.          | 84  |
| 4.16 | Chemical components of essential oil from <i>Ocimum canum</i> Sims.              | 86  |
| 4.17 | Chemical components of essential oil from <i>Ocimum gratissimum</i> Linn.        | 88  |
| 4.18 | Chemical components of essential oil from <i>Ocimum sanctum</i> Linn.            | 90  |
| 4.19 | Chemical components of essential oil from <i>Cinnamomum bejolghota</i><br>Sweet. | 92  |
| 4.20 | Chemical components of essential oil from <i>Piper sarmentosum</i> Roxb.         | 94  |
| 4.21 | Chemical components of essential oil from <i>Polygonum odoratum</i> Lour.        | 96  |
| 4.22 | Chemical components of essential oil from <i>Citrus aurantifolia</i> Swing.      | 98  |
|      | (Leaf)   |     |
| 4.23 | Chemical components of essential oil from <i>Citrus aurantifolia</i> Swing.      | 100 |
|      | (Peel)   |     |
| 4.24 | Chemical components of essential oil from <i>Citrus maxima</i> (Burm.) Merr.     | 102 |
|      | (Leaf)   |     |

|      |   |     |
|------|---|-----|
| 4.25 | Chemical components of essential oil from <i>Citrus maxima</i> (Burm.) Merr. (peel)     | 104 |
| 4.26 | Chemical components of essential oil from <i>Houttuynia cordata</i> Thunb               | 107 |
| 4.27 | Chemical components of essential oil from <i>Boesenbergia pandurata</i> (Roxb.) Schltr. | 109 |
| 4.28 | Chemical components of essential oil from <i>Curcuma longa</i> Linn                     | 110 |
| 4.29 | Chemical components of essential oil from <i>Curcuma zedoaria</i> (Berg) Roscoe.        | 113 |
| 4.30 | Chemical components of essential oil from <i>Zingiber cassumunar</i> Roxb               | 115 |
| 4.31 | Chemical components of essential oil from <i>Zingiber officinale</i> Roscoe.            | 117 |
| 4.32 | Ferric reducing power (EC1) of oils   | 120 |
| 4.33 | Trolox Equivalent Antioxidant Capacity (TEAC) values of oils                            | 123 |
| 4.34 | Tyrosinase inhibition activity of oils  | 125 |
| 4.35 | The solubility of sesame oil and lemongrass oil in various solvents                     | 128 |
| 4.36 | The refractive index of sesame oil and lemongrass oil                                   | 129 |
| 4.37 | Surface tension of sesame oil, lemongrass oil and various solvents                      | 130 |
| 4.38 | % w/w of Tween20, ethanol, oil and water (Tween 20: ethanol = 1:1)                      | 134 |
| 4.39 | % w/w of Tween20, ethanol, oil and water (Tween 20: ethanol = 2:1)                      | 135 |
| 4.40 | % w/w of Tween20, ethanol, oil and water (Tween 20: ethanol = 3:1)                      | 136 |
| 4.41 | % w/w of Tween20, ethanol, oil and water (Tween 20: ethanol = 9:1)                      | 137 |
| 4.42 | % w/w of Tween80, ethanol, oil and water (Tween 80: ethanol = 1:1)                      | 138 |
| 4.43 | % w/w of Tween80, ethanol, oil and water (Tween 80: ethanol = 2:1)                      | 139 |
| 4.44 | % w/w of Tween80, ethanol, oil and water (Tween 80: ethanol = 3:1)                      | 140 |
| 4.45 | % w/w of Tween80, ethanol, oil and water (Tween 80: ethanol = 9:1)                      | 141 |



|      |  |     |
|------|--|-----|
| 4.46 | Physical appearance of formulation of Tween 20, ethanol, oil and water<br>(Tween 20: ethanol = 1:1)            | 142 |
| 4.47 | Physical appearance of formulation of Tween 20, ethanol, oil and water<br>(Tween 20: ethanol = 2:1)            | 143 |
| 4.48 | Physical appearance of formulation of Tween 20, ethanol, oil and water<br>(Tween 20: Ethanol = 3:1)            | 144 |
| 4.49 | Physical appearance of formulation of Tween 20, ethanol, oil and water<br>(Tween 20: ethanol = 9:1)            | 145 |
| 4.50 | Physical appearance of formulation of Tween 80, ethanol, oil and water<br>(Tween 80: ethanol = 1:1)            | 146 |
| 4.51 | Physical appearance of formulation of Tween 80, ethanol, oil and water<br>(Tween 80: ethanol = 2:1)            | 147 |
| 4.52 | Physical appearance of formulation of Tween 80, ethanol, oil and water<br>(Tween 80: Ethanol = 3:1)            | 148 |
| 4.53 | Physical appearance of formulation of Tween 80, ethanol, oil and water<br>(Tween 80: ethanol = 9:1)            | 149 |
| 4.54 | Co-surfactants used in this study  | 155 |
| 4.55 | The formulation of microemulsion and conventional emulsion used in<br>this study                               | 175 |
| 4.56 | Droplet size and size distribution of internal phase and conductivity of<br>microemulsion systems without drug | 179 |
| 4.57 | Droplet size and size distribution of internal phase and conductivity of<br>microemulsion systems with drug    | 180 |



|      |   |     |
|------|---|-----|
| 4.58 | The amount of drug remaining in formula-1 of lemongrass oil<br>and sesame oil emulsion system   | 185 |
| 4.59 | The amount of drug remaining in formula-2 of lemongrass oil<br>and sesame oil emulsion system   | 186 |
| 4.60 | The amount of drug remaining in formula-3 of lemongrass oil<br>and sesame oil emulsion system   | 187 |
| 4.61 | The amount of drug remaining in formula-4 of lemongrass oil<br>and sesame oil emulsion system   | 188 |
| 4.62 | The amount of drug remaining in formula-5 of lemongrass oil<br>and sesame oil emulsion system   | 189 |
| 4.63 | Amount of drug released from lemongrass oil microemulsion systems   | 203 |
| 4.64 | Amount of drug released from sesame oil microemulsion systems   | 204 |
| 4.65 | Release rate of clotrimazole from the microemulsion systems and their<br>respective correlation coefficient of the linear equations obtained from<br>the plot between the amount of drug release vs. time | 205 |

## LIST OF FIGURES

| Figure |   | Page |
|--------|---|------|
| 2.1    | A hypothetical pseudo-ternary phase diagram of an oil/surfactant/water system | 13   |
| 2.2    | The pathway of melanin biosynthesis   | 17   |
| 3.1    | Essential oil hydrodistillation   | 30   |
| 3.2    | Tyrosinase inhibitory determination   | 35   |
| 3.3    | Vortex mixer  | 38   |
| 3.4    | Multipoint stirrer  | 38   |
| 3.5    | Whilhelmy type surface tensiometer  | 39   |
| 4.1    | The physical appearance of the oil used in this study                         | 58   |
| 4.2    | GC chromatogram of essential oil from <i>Apium graveolens</i> Linn.           | 63   |
| 4.3    | GC chromatogram of essential oil from <i>Anethum graveolens</i> Linn.         | 65   |
| 4.4    | GC chromatogram of essential oil from <i>Centella asiatica</i> Urban.         | 67   |
| 4.5    | GC chromatogram of essential oil from <i>Coriandrum sativum</i> Linn.         | 69   |
| 4.6    | GC chromatogram of essential oil from <i>Eryngium foetidum</i> Linn.          | 70   |
| 4.7    | GC chromatogram of essential oil from <i>Polyscias fruticosa</i> Harms.       | 72   |
| 4.8    | GC chromatogram of essential oil from <i>Eupatorium odoratum</i> Linn.        | 74   |
| 4.9    | GC chromatogram of essential oil from <i>Spilanthes acmella</i> Murr.         | 76   |
| 4.10   | GC chromatogram of essential oil from <i>Cymbopogon citratus</i> Stapf.       | 78   |

|      |   |     |
|------|---|-----|
| 4.11 | GC chromatogram of essential oil from <i>Coleus amboinicus</i> Lour.  | 80  |
| 4.12 | GC chromatogram of essential oil from <i>Melissa officinalis</i> Linn.  | 82  |
| 4.13 | GC chromatogram of essential oil from <i>Ocimum basilicum</i> Linn.   | 84  |
| 4.14 | GC chromatogram of essential oil from <i>Ocimum canum</i> Sims.   | 86  |
| 4.15 | GC chromatogram of essential oil from <i>Ocimum gratissimum</i> Linn.   | 88  |
| 4.16 | GC chromatogram of essential oil from <i>Ocimum sanctum</i> Linn.   | 90  |
| 4.17 | GC chromatogram of essential oil from <i>Cinnamomum bejolghota</i> Sweet.   | 92  |
| 4.18 | GC chromatogram of essential oil from <i>Piper sarmentosum</i> Roxb.  | 94  |
| 4.19 | GC chromatogram of essential oil from <i>Polygonum odoratum</i> Lour.   | 96  |
| 4.20 | GC chromatogram of essential oil from <i>Citrus aurantifolia</i> Swing. (Leaf)  | 98  |
| 4.21 | GC chromatogram of essential oil from <i>Citrus aurantifolia</i> Swing. (Peel)  | 100 |
| 4.22 | GC chromatogram of essential oil from <i>Citrus maxima</i> (Burm.) Merr. (Leaf)                                       | 102 |
| 4.23 | GC chromatogram of essential oil from <i>Citrus maxima</i> (Burm.) Merr. (Peel)                                       | 104 |
| 4.24 | GC chromatogram of essential oil from <i>Houttuynia cordata</i> Thunb.  | 106 |
| 4.25 | GC chromatogram of essential oil from <i>Boesenbergia pandurata</i> (Roxb.)   | 108 |
| 4.26 | GC chromatogram of essential oil from <i>Curcuma longa</i> Linn.  | 110 |
| 4.27 | GC chromatogram of essential oil from <i>Curcuma zedoaria</i> (Berg) Rosc.  | 113 |
| 4.28 | GC chromatogram of essential oil from <i>Zingiber cassumunar</i> Roxb.  | 115 |
| 4.29 | GC chromatogram of essential oil from <i>Zingiber officinale</i> Rosc.  | 117 |
| 4.30 | Cytotoxicity of the oil of <i>C. citratus</i> (lemongrass, A) and <i>S. indicum</i> (sesame, B) on normal human PBMCs | 131 |

|      |  |     |
|------|--|-----|
| 4.31 | Effect of Tween 20 and ethanol with water and lemongrass oil<br>( <i>Cymbopogon citratus</i> Stapf.)                                       | 150 |
| 4.32 | Effect of Tween 80 and ethanol with water and lemongrass oil<br>( <i>Cymbopogon citratus</i> Stapf.)                                       | 151 |
| 4.33 | Microemulsion area (M) of lemongrass oil in various surfactant<br>systems with ethanol (1 : 1)   | 156 |
| 4.34 | Microemulsion area (M) of lemongrass oil in various surfactant<br>systems with ethanol (1 : 2)   | 156 |
| 4.35 | Microemulsion area (M) of sesame oil in various surfactant<br>systems with ethanol (1 : 1)   | 157 |
| 4.36 | Microemulsion area (M) of sesame oil in various surfactant<br>systems with ethanol (1 : 2)   | 157 |
| 4.37 | Microemulsion area (M) of lemongrass oil in Tween 20 and sesame oil<br>in Tween 85 without cosurfactant                                    | 158 |
| 4.38 | Effect of carbon atom in co-surfactant molecules on the phase diagram<br>of lemongrass oil (the ratio of surfactant: co-surfactant is 1:2) | 160 |
| 4.39 | Effect of carbon atom in co-surfactant molecules on the phase diagram<br>of lemongrass oil (the ratio of surfactant: co-surfactant is 1:1) | 161 |
| 4.40 | Effect of carbon atom in co-surfactant molecules on the phase diagram<br>of lemongrass oil (the ratio of surfactant: co-surfactant is 2:1) | 162 |
| 4.41 | Effect of carbon atom in co-surfactant molecules on the phase diagram<br>of sesame oil (the ratio of surfactant: co-surfactant is 1:2)     | 163 |
| 4.42 | Effect of carbon atom in co-surfactant molecules on the phase diagram<br>of sesame oil (the ratio of surfactant: co-surfactant is 1:1)     | 164 |

|      |   |     |
|------|---|-----|
| 4.43 | Effect of carbon atom in co-surfactant molecules on the phase diagram of sesame oil (the ratio of surfactant: co-surfactant is 2:1)   | 165 |
| 4.44 | Effect of surfactant : co-surfactant on the microemulsion area (M) of lemongrass oil  | 166 |
| 4.45 | Effect of surfactant : co-surfactant on the microemulsion area (M) of sesame oil  | 167 |
| 4.46 | Effect of NaCl on the microemulsion area (M) of lemongrass oil (surfactant system is Tween 20 : Ethanol = 2:1)  | 168 |
| 4.47 | Effect of CaCl <sub>2</sub> on the microemulsion area (M) of lemongrass oil (surfactant system is Tween 20 : Ethanol = 2:1)   | 168 |
| 4.48 | Effect of NaCl on the microemulsion area (M) of sesame oil (surfactant system is Tween 85 : Ethanol = 2:1)  | 169 |
| 4.49 | Effect of CaCl <sub>2</sub> on the microemulsion area (M) of sesame oil (surfactant system is Tween 85 : Ethanol = 2:1)   | 169 |
| 4.50 | Effect of chemical structure on the microemulsion area (M) of lemongrass oil (surfactant system is Tween 20 : Ethanol = 2:1) and sesame oil (surfactant system is Tween 85 : Ethanol = 2:1) | 170 |
| 4.51 | Effect of pH on the microemulsion area (M) of lemongrass oil (surfactant system is Tween 20 : Ethanol = 2:1)  | 171 |
| 4.52 | Effect of pH on the microemulsion area (M) of sesame oil (surfactant system is Tween 85 : Ethanol = 2:1)  | 171 |
| 4.53 | Appearance of the lemongrass oil microemulsion without drug   | 176 |
| 4.54 | Appearance of the lemongrass oil microemulsion with drug  | 176 |
| 4.55 | Appearance of the sesame oil microemulsion without drug   | 177 |

|      |   |     |
|------|---|-----|
| 4.56 | Appearance of the sesame oil microemulsion with drug  | 177 |
| 4.57 | The amount of drug remaining in lemongrass oil microemulsion system<br>(HT = 45°C, RT = Room temperature about 30°C, LT = 4°C)  | 190 |
| 4.58 | The amount of drug remaining in sesame oil microemulsion system<br>(HT = 45°C, RT = Room temperature about 30°C, LT = 4°C)  | 191 |
| 4.59 | The outer appearance of lemongrass oil microemulsion systems<br>without drug which were kept in various conditions<br>(HT = 45°C, RT = Room temperature about 30°C, LT = 4°C,<br>HCC = heating and cooling cycle) | 192 |
| 4.60 | The outer appearance of lemongrass oil microemulsion systems<br>with drug which were kept in various conditions<br>(HT = 45°C, RT = Room temperature about 30°C, LT = 4°C,  | 193 |
| 4.61 | The outer appearance of sesame oil microemulsion systems<br>without drug which were kept in various conditions<br>(HT = 45°C, RT = Room temperature about 30°C, LT = 4°C,   | 194 |
| 4.62 | The outer appearance of sesame oil microemulsion systems<br>with drug which were kept in various conditions<br>(HT = 45°C, RT = Room temperature about 30°C, LT = 4°C,  | 195 |
| 4.63 | Comparison of characteristic appearance of various formulations<br>stored at 45°C   | 196 |
| 4.64 | Comparison of characteristic appearance of various formulations<br>stored at 30°C   | 197 |
| 4.65 | Comparison of characteristic appearance of various formulations<br>stored at 4°C  | 198 |

|      |  |     |
|------|--|-----|
| 4.66 | Comparison of characteristic appearance of various formulations<br>stored in heating-cooling cycle | 199 |
| 4.67 | The product of miroemulsion of sesame oil and lemongrass oil                                       | 200 |
| 4.68 | The releasing profile of drug from lemongrass oil<br>microemulsion systems                         | 206 |
| 4.69 | The releasing profile of drug from sesame oil microemulsion systems                                | 206 |



## ABBREVIATIONS AND SYMBOLS

|       |   |  |
|-------|---|--|
| Abs   | = | Absorbance                               |
| °C    | = | Degree celcius                           |
| Fe    | = | Iron                                     |
| g     | = | Gram                                     |
| GC-MS | = | Gas Chromatography and Mass Spectrometer |
| hr    | = | Hour                                     |
| µg    | = | Microgram                                |
| µL    | = | Microliter                               |
| µM    | = | Micromolar                               |
| mL    | = | Milliliter                               |
| mM    | = | Millimolar                               |
| M     | = | Molar                                    |
| min   | = | Minute                                   |
| nm    | = | Nanometer                                |
| %     | = | Percent                                  |
| v/v   | = | Volume/Volume                            |
| w/w   | = | Weight/Weight                            |