

REFERENCE

- [1] R. Feynman, "Simulating physics with computers," *Int. J. Theor. Phys.*, **21**, 467 (1982).
- [2] D. P. DiVincenzo, "Quantum computation," *Science*, **270**, 255 (1995).
- [3] P.W. Shor, "Algorithms for quantum computation: Discrete logarithms and factoring," *Proc. 35th Annual Symposium on Foundation of Computer Science*, 124 (1994).
- [4] D. R. Simon, "On the power of quantum computation," *35th Annual Symposium on Foundations of Computer Science*, 116 (1994).
- [5] D. S. Abrams, S. Lloyd, "Simulation of many-body Fermi systems on a universal quantum computer," *Phys. Rev. Lett.*, **79**, 2586 (1997).
- [6] A. Aspuru-Guzik, A. D. Dutoi, P. J. Love, and M. Head-Gordon, "Simulated quantum computation of Molecular energies," *Science*, **309**, 1704 (2005).
- [7] R. Dumke, et al., "Micro-optical realization of arrays of selectively addressable dipole traps: A scalable configuration for quantum computation with atomic qubits," *Phys. Rev. Lett.*, **89**, 097903 (2002).
- [8] D. Schrader, et al., "Neutral atom quantum register," *Phys. Rev. Lett.*, **93**, 150501 (2004).
- [9] D. N. Karl, X. Li, and D. S. Weiss, "Imaging single atoms in a three-dimensional array," *Nature Phys.*, **3**, 556 (2007).
- [10] M. Karski, et al., "Nearest-neighbor detection of atoms in a 1D optical lattice by fluorescence imaging," *Phys. Rev. Lett.*, **102**, 053001 (2009).

- [11] D. Jaksch, J. I. Cirac, P. Zoller, S. L. Rolston, R. R. Cote, and M. D. Lukin, “Fast quantum gates for neutral atoms,” *Phys. Rev. Lett.* **85**, 2208–2211 (2000).
- [12] L. Isenhower, E. Urban, X. L. Zhang, A. T. Gill, T. Henage, T. A. Johnson, T. G. Walker, and M. Saffman, “Demonstration of a Neutral Atom Controlled-NOT Quantum Gate,” *Phys. Rev. Lett.*, **104**, 010503 (2010).
- [13] T. Wilk, A. Gaetan, C. Evellin, J. Wolters, Y. Miroshnychenko, P. Grangier, and A. Browaeys, “Entanglement of two individual neutral atoms using Rydberg blockade,” *Phys. Rev. Lett.*, **104**, 010502 (2010).
- [14] R. Gehr, J. Volz, G. Dubois, T. Steinmetz, Y. Colombe, B. L. Lev, R. Long, J. Estève, and J. Reichel, “Cavity-Base Single Atom Preparation and High-Fidelity Hyperfine State Readout,” *Phys. Rev. Lett.*, **104**, 203602 (2010).
- [15] C. J. Hood, T. W. Lynn, A. C. Doherty, A. S. Parkins, and H. J. Kimble, “The Atom-Cavity Microscope: Single Atom Bound in Orbit by Single Photons,” *Science*, **287**, 1447 (2000).
- [16] W. S. Bakr, A. Peng, M. E. Tai, R. Ma, J. Simon, J. I. Gillen, S. Fölling, L. Pollet, and M. Greiner, “Probing the superfluid-to-Mott-insulator transition at the single-atom level,” *Science*, **329**, 547 (2010).
- [17] C. Weitenberg, M. Endres, J. F. Sherson, M. Cheneau, P. Schauß, T. Fukuhara, I. Bloch, and S. Kuhr, “Single-Spin Addressing in an Atomic Mott Insulator,” *Nature*, **471**, 319 (2011).
- [18] M. T. DePue, C. McCormick, S. L. Winoto, S. Oliver, and D. S. Weiss, “Unity occupation of sites in a 3D optical lattice,” *Phys. Rev. Lett.*, **82**, 2262 (1999).
- [19] N. Schlosser, G. Reymond, I. Protsenko, P. Grangier, “Sub-Poissonian loading of single atoms in a microscopic dipole trap,” *Nature*, **411**, 1024 (2001).

- [20] W. Alt, D. Schrader, S. Kuhr, M. Müller, V. Gomer, and D. Meschede, “Single atoms in a standing-wave dipole trap,” *Phys. Rev. A*, **67**, 033403 (2003).
- [21] T. Grünzweig, A. Hilliard, M. McGovern, and M. F. Andersen, “Near-deterministic preparation of a single atom in an optical microtrap,” *Nature Phys.*, **6**, 951 (2010).
- [22] L. Marcassa, V. Bagnato, Y. Wang, C. Tsao, J. Weiner, O. Dulieu, Y. B. Band, and P. S. Julienne, “Collisional loss rate in a magneto-optical trap for sodium atoms: light intensity dependence,” *Phys. Rev. A*, **47**, R4563 (1993).
- [23] T. Walker and P. Feng, “Measurements of collisions between laser-cooled atoms,” *Adv. At. Mol. Opt. Phys.*, **34**, 125 (1994).
- [24] T. Walker and D. R. Pritchard, “Effects of hyperfine structure on alkali trap-loss collisions,” *Laser Phys.*, **4**, 1085 (1994).
- [25] L. Marcassa, S. R. Muniz, and J. Flemming, “Optical catalysis in a Na-vapor cell MOT,” *Braz. J. Phys.*, **27**, 238 (1997).
- [26] B. Ueberholz, S. Kuhr, D. Frese, D. Meschede, and V. Gomer, “Counting cold collisions,” *J. Phys. B*, **33**, L135 (2000).
- [27] B. Ueberholz, S. Kuhr, D. Frese, V. Gomer, and D. Meschede, “Cold collisions in a high-gradient magneto-optical trap,” *J. Phys. B*, **35**, 4899 (2002).
- [28] E. Raab, M. Prentiss, A. Cable, S. Chu, and D. Pritchard, “Trapping of neutral-Sodium atoms with radiation pressure,” *Phys. Rev. Lett.*, **59**, 2631 (1987).
- [29] C. Kittel and H. Kroemer, *Thermal Physics*, W. H. Freeman and Company, 1998.
- [30] H. J. Straten and P. Metcalf, *Laser Cooling and Trapping*, Springer-Verlag New York, Inc., 2002.

- [31] D. J. Griffiths, *Introduction to Quantum Mechanics*, 2nd Ed. Prentice Hall, 2005.
- [32] J. Dalibard and C. Cohen-Tannoudji, “Dress-atom approach to atomic motion in laser light: the dipole force revisited,” *J. Opt. Soc. Am. B.*, **2**, 1701 (1985).
- [33] C. Cohen-Tannoudji, J. Dupont-Roc, and G. Grynberg, *Atom-Photon Interactions*, Wiley-VCH Verlag GmbH & Co. KGaA, Weinheim, 2004.
- [34] N. Davidson, H.J. Lee, C.S. Adams, M. Kasevich, and S. Chu, “Long atomic coherence times in an optical dipole trap,” *Phys. Rev. Lett.*, **74**, 1311 (1995).
- [35] T. Kuga, Y. Torii, N. Shiokawa, and T. Hirano, “Novel optical trap of atoms with a doughnut beam,” *Phys. Rev. Lett.*, **78**, 4713 (1997).
- [36] R.J. Cook and R.K. Hill, “An electromagnetic mirror for neutral atoms,” *Opt. Commun.*, **43**, 258 (1982).
- [37] S. Chu, J.E. Bjorkholm, A. Ashkin and A. Cable, “Experimental observation of optically trapped atoms,” *Phys. Rev. Lett.*, **57**, 314 (1986).
- [38] I. Bloch, “Ultracold quantum gases in optical lattices,” *Nature Phys.*, **1**, 23 (2005).
- [39] J. Dalibard and C. Cohen-Tannoudji, “Laser cooling below the Doppler limit by polarization gradients: simple theoretical models,” *J. Opt. Soc. Am. B.*, **6**, 2023 (1989).
- [40] A. Aspect, J. Dalibard, A. Heidmann, C. Salomon, and C. Cohen-Tannoudji, “Cooling atoms with stimulated emission,” *Phys. Rev. Lett.*, **57**, 1688 (1986).
- [41] B. Bussery and M. Aubert-Frecon, “Multipolar long-range electrostatic, dispersion, and induction energy terms for the interactions between two identical alkali atoms Li, Na, K, Rb and Cs in various electronic states,” *J. Chem. Phys.*, **82**, 3224 (1985).

- [42] A. Gallagher and D. E. Pritchard, “Exoergic collisions of cold Na^*-Na ,” *Phys. Rev. Lett.*, **63**, 957 (1989).
- [43] J. Weiner, *Cold and Ultracold Collisions in Quantum Microscopic and Mesoscopic System*, Cambridge University Press, 2004.
- [44] P. S. Julienne and J. Vigue, “Cold collision of ground- and excited-state alkali-metal atoms,” *Phys. Rev. A.*, **44**, 4464 (1991).
- [45] N. Thaicharoen, *Light assisted collisions of cold atoms in an optical dipole trap*, M.S. thesis, Chiang Mai University, 2012.
- [46] *Manual of External Cavity Diode Laser models ECD-003*, MOG Laboratories Ltd, 2010.
- [47] *Manual of External Cavity Diode Laser controller models DLC-202, DLC-502*, MOG Laboratories Ltd, 2009.
- [48] A. S. Arnold, J. S. Wilson, and M. G. Boshier, “A simple extended-cavity diode laser,” *Rev. Sci. Instrum.*, **69**, 1236 (1998).
- [49] C. J. Hawthorn, K. P. Weber, and R. E. Scholten, “Littrow configuration tunable external cavity diode laser with fixed direction output beam,” *Rev. Sci. Instrum.*, **72**, 4477 (2001).
- [50] C. Tuchendler, A. M. Lance, A. Browaeys, Y. R. P. Sortais, and P. Grangier, “Energy distribution and cooling of a single atom in an optical tweezer,” *Phys. Rev. A*, **78**, 033425 (2008).
- [51] N. Schlosser, G. Reymond, and P. Grangier, “Collisional blockade in microscopic optical dipole traps,” *Phys. Rev. Lett.*, **89**, 023005(2002).

- [52] B. Bussery, and M. Aubertfrecon, “Multipolar long-range electrostatic, dispersion, and induction energy terms for the interaction between two identical alkali atoms Li, Na, K, Rb and Cs in various electronic states,” *J. Chem. Phys.*, **82**, 3224 (1985).
- [53] A. R. Gorges, N. S. Bingham, M. G. DeAngelo, M. S. Harmilton, and J. L. Roberts, “Light-assisted collisional loss in a $^{85/87}\text{Rb}$ ultracold optical trap,” *Phys. Rev. A*, **78**, 033420 (2008).
- [54] M. G. Peters, D. Hoffmann, J. D. Tobiason, and T. Walker, “Laser-induced ultracold $\text{Rb}(5S_{1/2})+\text{Rb}(5P_{1/2})$ collisions,” *Phys. Rev. A*, **50**, R906 (1994).
- [55] J. F. Sherson, C. Weitenberg, M. Endres, M. Cheneau, I. Bloch, and S. Kuhr, “Single-atom-resolved fluorescence imaging of an atomic Mott insulator,” *Nature*, **467**, 68 (2010).
- [56] D. A. Steck, “Rubidium 85 D line data,” available online at <http://steck.us/alkalidata> (revision 2.1.4, 23 December 2010).