

CONTENTS

	Page
Acknowledgement	d
Abstract in Thai	e
Abstract in English	g
List of Tables	k
List of Figures	l
List of Abbreviations	p
List of Symbols	q
Statement of Originality in Thai	r
Statement of Originality in English	s
Chapter 1 Introduction	1
Chapter 2 Literature review	4
2.1 Anaerobic digestion	4
2.2 Microbial consortium and microbiological process	5
2.3 Factors affecting anaerobic digestion	13
2.4 Co-digestion	16
2.5 Process design	19
2.6 Denaturing gradient gel electrophoresis (DGGE)	20
2.7 Application of DGGE techniques for microbial community study in biogas reactors	23
Chapter 3 Materials and methods	25
3.1 Sludge sampling	25
3.2 Microbial community analysis	27

	Page
Chapter 4 Microbial community in anaerobic co-digestion of pig manure with Napier grass in channel digester-upflow anaerobic sludge blanket (CD-UASB) and completely stirred tank reactor (CSTR)	34
4.1 Introduction	34
4.2 Materials and methods	35
4.3 Results and discussion	37
Chapter 5 Microbial community in anaerobic co-digestion of pig manure with food waste in channel digester-upflow anaerobic sludge blanket (CD-UASB) and completely stirred tank reactor (CSTR)	71
5.1 Introduction	71
5.2 Materials and methods	73
5.3 Results and discussion	75
Chapter 6 General discussion	100
Chapter 7 Conclusion	106
References	107
Appendix	121
Appendix A	122
Appendix B	128
Appendix C	130
Appendix D	132
Appendix E	133
Curriculum Vitae	141

LIST OF TABLES

		Page
Table 2.1	Examples of hydrolytic bacteria in anaerobic reactors	7
Table 2.2	Examples of methanogenic bacteria	11
Table 2.3	Chemical compositions (in % of TS) of animal slurries	17
Table 3.1	Sampling time	25
Table 3.2	Sequence of the primers used	29
Table 3.3	Reaction mixtures for amplification of bacterial 16S rDNA	30
Table 3.4	Reaction mixtures for amplification of archaeal 16S rDNA	30
Table 3.5	Preparation of DGGE gel solution for a polyacrylamide gel with a denaturant gradient between 20-45% and 30-55%	31
Table 3.6	Reaction mixtures for colony PCR amplification	33
Table 4.1	Phylogenetic affiliation of the bacterial 16S rDNA sequences from DGGE bands using BLAST search in GenBank and EzTaxon database from sludge in CD-UASB and CSTR co-digested with Napier grass under HRT 30 days	51
Table 4.2	Phylogenetic affiliation of the archaeal 16S rDNA sequences from DGGE bands using BLAST search in GenBank and EzTaxon database from sludge in CD-UASB and CSTR co-digested with Napier grass at HRT 30 days	68
Table 5.1	Phylogenetic affiliation of the bacterial 16S rDNA sequences from DGGE bands using BLAST search in GenBank and EzTaxon database from sludge in CD-UASB and CSTR co-digested with food waste under HRT 20 days	86
Table 6.1	Species richness, methane yield and COD removal efficiency in all the reactors studied	103


LIST OF FIGURES

	Page
Figure 2.1 Electron micrographs of microbial communities in anaerobic methanogenic granules	5
Figure 2.2 Schematic overview of four main steps in the anaerobic digestion process	6
Figure 2.3 Phylogenetic classification of methanogenic bacteria	12
Figure 2.4 Schematic sketches of the batch and continuous reactor	19
Figure 2.5 Migration of DNA bands by DGGE technique	21
Figure 2.6 Diagram of steps in microbial community analysis by DGGE technique	23
Figure 3.1 Sampling point of the lab scale CD-UASB reactor	26
Figure 3.2 Sampling point of the lab scale CSTR	26
Figure 4.1 DGGE profiles and dendrograms (UPGMA clustering) of bacterial 16S rDNA fragments amplified from sludge in CD-UASB reactor co-digested with Napier grass under different HRT	41
Figure 4.2 DGGE profiles and dendrograms (UPGMA clustering) of bacterial 16S rDNA fragments amplified from sludge in CSTR co-digested with Napier grass under different HRT	42
Figure 4.3 Cluster analysis of bacterial banding patterns at steady state from reactors co-digested with Napier grass under different HRT	44
Figure 4.4 DGGE profile of bacterial 16S rDNA fragments amplified from sludge in CD-UASB reactor co-digested with Napier grass at HRT 30 days	49
Figure 4.5 DGGE profile of bacterial 16S rDNA fragments amplified from sludge in CSTR co-digested with Napier grass at HRT 30 days	50

Figure 4.6	Phylogenetic tree of 16S rDNA bacterial sequences from sludge DNA in CD-UASB reactor co-digested with Napier grass at HRT 30 days as determined by the neighbor-joining method	58
Figure 4.7	Phylogenetic tree of 16S rDNA bacterial sequences from sludge DNA in CSTR co-digested with Napier grass at HRT 30 days as determined by the neighbor-joining method	59
Figure 4.8	DGGE profiles of archaeal 16S rDNA fragments amplified from sludge in CD-UASB reactor co-digested with Napier grass and under different HRT	62
Figure 4.9	DGGE profiles of archaeal 16S rDNA fragments amplified from sludge in CSTR co-digested with Napier grass under different HRT	63
Figure 4.10	DGGE profile of archaeal 16S rDNA fragments amplified from seed and sludge samples in CD-UASB and CSTR co-digested with Napier grass	64
Figure 4.11	DGGE profile of archaeal 16S rDNA fragments amplified from sludge in CD-UASB reactor co-digested with Napier grass and at HRT 30 days	66
Figure 4.12	DGGE profile of archaeal 16S rDNA fragments amplified from sludge in CSTR co-digested with Napier grass at HRT 30 days	67
Figure 4.13	Relative abundance of methanogenic archaea in the clone library	70
Figure 5.1	DGGE profiles and dendrograms (UPGMA clustering) of bacterial 16S rDNA fragments amplified from sludge in CD-UASB reactor co-digested with food waste under different HRT	78

	Page
Figure 5.2 DGGE profiles and dendrograms (UPGMA clustering) of bacterial 16S rDNA fragments amplified from sludge in CSTR co-digested with food waste under different HRT	79
Figure 5.3 Cluster analysis of bacterial banding patterns at steady state from reactors co-digested with food waste under different HRT	80
Figure 5.4 DGGE profile of bacterial 16S rDNA fragments amplified from sludge in CD-UASB reactor co-digested with food waste at HRT 20 days	84
Figure 5.5 DGGE profile of bacterial 16S rDNA fragments amplified from sludge in CSTR co-digested with food waste at HRT 20 days	85
Figure 5.6 Phylogenetic tree of 16S rDNA bacterial sequences from sludge DNA in CD-UASB reactor co-digested with food waste at HRT 20 days as determined by the neighbor-joining method	90
Figure 5.7 Phylogenetic tree of 16S rDNA bacterial sequences from sludge DNA in CSTR co-digested with food waste at HRT 20 days as determined by the neighbor-joining method.	91
Figure 5.8 DGGE profiles of archaeal 16S rDNA fragments amplified from sludge in CD-UASB reactor co-digested with food waste under different HRT	94
Figure 5.9 DGGE profiles of archaeal 16S rDNA fragments amplified from sludge in CSTR co-digested with food waste under different HRT	95
Figure 5.10 DGGE profile of archaeal 16S rDNA fragments amplified from seed and sludge samples in CD-UASB and CSTR co-digested with food waste	96
Figure 5.11 DGGE profile of archaeal 16S rDNA fragments amplified from sludge in CD-UASB reactor co-digested with food waste at HRT 20 days	98
Figure 5.12 DGGE profile of archaeal 16S rDNA fragments amplified from sludge in CSTR co-digested with food waste at HRT 20 days	99

Figure 6.1 Cluster analysis of bacterial banding pattern at steady state from reactors co-digested with Napier grass or food waste under different HRT



ลิขสิทธิ์มหาวิทยาลัยเชียงใหม่
Copyright© by Chiang Mai University
All rights reserved