

CONTENTS

	Page
Acknowledgement	iv
Abstract in Thai	v
Abstract	vi
List of Tables	ix
List of Figures	xii
Chapter 1 Introduction	1
Chapter 2 Theory	3
2.1 Binary star systems	3
2.2 Spectroscopy	9
2.3 Instrument: Spectrograph	15
Chapter 3 Observation and Data reductions	18
Chapter 4 Determination of radial velocity curve	25
4.1 The Least-Squares Deconvolution method	25
4.2 Selection of spectral range	26
4.3 Determination of Least-Squares Deconvolution profiles	26
4.4 Improvement of a radial velocity	27
4.5 Orbital analysis from radial velocities	28
Chapter 5 Determination of projected rotational velocity	36
5.1 The Fourier-transformed method	36
5.2 Reconstruction of LSD profiles	37

5.3 Determination of projected rotational velocities	38
Chapter 6 Conclusion	42
Bibliography	44
Curriculum Vitae	46



ลิขสิทธิ์มหาวิทยาลัยเชียงใหม่
Copyright© by Chiang Mai University
All rights reserved

LIST OF TABLES

	Page
Table 2.1	The catalog of RS CVn systems characteristic. 4
Table 2.2	The Balmer series of hydrogen. 12
Table 4.1	The information of radial velocity value of each spectrum before and after correction. 33
Table 4.2	Orbital parameters of HD 26337 as derived from the data collected by Washuttl (2006). 34
Table 4.3	Orbital parameters of HD 26337 analyzed from HERCULES data. 35
Table 5.1	The first minimum value and projected rotational velocity of EI Eri were calculated from Fourier transform according to phase sequence 1 to 0. 40
Table 6.1	The new orbital solution of EI Eri. 42

LIST OF FIGURES

	Page
Figure 2.1 The example spectra are showing relation between the shifted line and object motion which refer with the rest frame. (Diagram from http://www.atnf.csiro.au/outreach/education/senior/astrophysics/spectra info.html)	5
Figure 2.2 The relative orbit of a binary located in three dimensions and defined by the angle Ω , ω and i . (Hildith, 2001)	6
Figure 2.3 The example of radial-velocity curves for the two stars in a binary system with a circular orbit. The semiamplitudes are $K_1 = 100$ km/s and $K_2 = 200$ km/s, giving a mass ratio $q = m_2/m_1 = 0.5$. The systemic velocity is $\gamma = 0$ km/s. The positions of the ascending (a) and descending (d) nodes are shown and the points of zero (z) orbital radial velocity. In the upper panel, the observer views the binary orbit from the right along z-z.	7
Figure 2.4 The radial velocity of a circular orbit ($e = 0$, solid line), and elliptical orbits. The dashed line represent the curve of $e = 0.1$ and $\omega = 0^\circ$ and the dotted line represents the curve of $e = 0.5$ and $\omega = 0^\circ$. The semi-amplitude of all curves is $K = 100$ km/s and $\gamma = 0$ km/s. (Komonjinda, 2008)	8
Figure 2.5 Showing the dispertional energy compare with wavelength at difference temperatures scale according to Planck's law. (Diagram from http://www.ucolick.org)	10
Figure 2.6 The star has inclination i to the line-of-sight of an observer on the Earth and rotational velocity v at the equator. (Diagram from http://en.wikipedia.org/wiki/Stellarrotation)	11

Figure 2.7	The computed line profiles as broadened from the rotation of a star. The labels are the value of $v \sin i$ in km/s. (Komonjinda, 2008)	12
Figure 2.8	The blended line that occur from broadening line with the other. (Komonjinda, 2008)	13
Figure 2.9	The example spectrum of Vega (Henson) is showed the reference lines according to Balmer series. (Diagram from Astronomical Spectroscopy for amateurs)	13
Figure 2.10	Telluric absorption lines due to oxygen water in the Earth's atmosphere. (Diagram from http://www.threehillsobservatory.co.uk/astro/spectra 9.htm)	14
Figure 2.11	Optical design of the HERCULES spectrograph. (Komonjinda, 2008)	16
Figure 3.1	The binary system EI Eri and neighbor star HD 26294.	18
Figure 3.2	Sample spectrum from HERCULES spectrograph.	19
Figure 3.3	Example spectrum of EI Eri from 11/12/04 shows the different level continuum and gap between orders.	20
Figure 3.4	The synthetic spectrum of binary system computed assuming fundamental parameter by using VALD showing synthetic spectrum no convolve (upper) and was convolved (bottom) with rotational velocity around 50 km/s in particular wavelength 4837-4888 Å.	21
Figure 3.5	The spectrum profiles are compared to show the different observed spectrum before (solid line) and after (dotted line) normalization. We can see that the continuum level become better and useful to the LSD technique.	22
Figure 3.6	The example spectrum in order 7-13 after were reduced.	23
Figure 3.7	Both two spectra in wavelength range 4848-4880 Angstrom are compared each other to see the variation of spectral which both of them are observed at a different time.	23
Figure 3.8	The reduction method of HERCULES spectra.	24

Figure 4.1	The example original LSD profile represent all individual spectral line.	27
Figure 4.2	RV curve was obtained from each LSD profiles.	28
Figure 4.3	The expample LSD profile was adjusted Barycentric correction and substracted 81.35 km/s.	29
Figure 4.4	The Least-Squares Deconvolution method and improved of a radial velocities curve.	32
Figure 4.5	The RV curve was indicated total adjusted radial velocity of spectrum.	34
Figure 4.6	Phase plot of EI Eri radial velocities.	35
Figure 5.1	Reconstruction of LSD profile by inverse and trimed both end of this spectra.	37
Figure 5.2	A fourier transformed rotational broadening function $g(\sigma)$ was indicated the first zero minimum.	38
Figure 5.3	LSD profile with phase shows the variation in line profiles.	39
Figure 6.1	The variation LSD profiles according to phase of EI Eri.	43