

REFERENCES

- [1] Sambuy Y. Sideways Glance: Genome wide association studies for type 2 diabetes mellitus. *Genes & nutrition.* 2007; 2(3):245-8.
- [2] Ahima RS. Adipose tissue as an endocrine organ. *Obesity (Silver Spring, Md).* 2006; 14 Suppl 5:242s-9s.
- [3] Parichart Kumcheen, “Thailand under obesity crisis”, 2009, http://www.siam.edu/siamedu_thai_mainpage/images/stories/article/54/siam_article_7_54.pdf, 17June2014
- [4] Karelis AD, Faraj M, Bastard JP, St-Pierre DH, Brochu M, et al. The metabolically healthy but obese individual presents a favorable inflammation profile. *J ClinEndocrinolMetab.* 2005; 90(7):4145-50.
- [5] Unger RH (1), Clark GO, Scherer PE, Orci L. Lipid homeostasis, lipotoxicity and the metabolic syndrome. *BiochimBiophysActa.* 2010; 1801(3):209-14. Review.
- [6] Lee WJ, Koh EH, Won JC, Kim MS, Park JY, Lee KU. Obesity: the role of hypothalamic AMPactivated protein kinase in body weight regulation. *Int J Biochem Cell Biol.* 2005 Nov; 37(11):2254-9.
- [7] Kelley DE, Goodpaster BH. Skeletal muscle triglyceride.An aspect of regional adiposity and insulin resistance.*Diabetes Care.* 2001 May;24(5):933-41.
- [8] Fruhbeck G, Gomez-Ambrosi J, Muruzabal FJ, Burrel MA. The adipocyte: A model for integration of endocrine and metabolic signaling in energy metabolism regulation.*Am J PhysiolEndocrinolMetab.* 2001 June; 280(6):E827–47.
- [9] Galic S, Oakhill JS, Steinberg GR. Adipose tissue as an endocrine organ. *Mol Cell Endocrinol.* 2010 Mar 25; 316(2):129-39.

- [10] Arner P, Spalding KL. Fat cell turnover in humans. *Biochem Biophys Res Commun.* 2010 May 21; 396(1):101-4.
- [11] Spalding KL, Arner E, Westermark PO, Bernard S, Buchholz BA, Bergmann O, Blomqvist L, Hoffstedt J, Näslund E, Britton T, Concha H, Hassan M, Rydén M, Frisén J, Arner P. Dynamics of fat cell turnover in humans. *Nature.* 2008 Jun 5;453(7196):783-7.
- [12] Salans LB, Horton ES, Sims EA. Experimental obesity in man: cellular character of the adipose tissue. *J Clin Invest.* 1971 May; 50(5):1005-11.
- [13] Löfgren P, Andersson I, Adolfsson B, Leijonhufvud BM, Hertel K, Hoffstedt J, Arner P. Longterm prospective and controlled studies demonstrate adipose tissue hypercellularity and relative leptin deficiency in the postobese state. *J Clin Endocrinol Metab.* 2005 Nov; 90(11):6207-13.
- [14] Rayalam S, Della-Fera MA, Baile CA. Phytochemicals and regulation of the adipocyte life cycle. *J Nutr Biochem.* 2008 Nov; 19(11):717-26.
- [15] Gregoire FM. Adipocyte differentiation: from fibroblast to endocrine cell. *Exp Biol Med (Maywood).* 2001 Dec; 226(11):997-1002.
- [16] Moreno-Navarrete J, Fernández-Real J. Adipocyte Differentiation. In: Symonds ME, editor. *Adipose Tissue Biology:* Springer New York; 2012. p. 17-38.
- [17] Green H, Kehinde O. An established preadipose cell line and its differentiation in culture II. Factors affecting the adipose conversion. *Cell.* 1975 May;5(1):19-27.
- [18] Cornelius P, MacDougald OA, Lane MD. Regulation of adipocyte development. *Annu Rev Nutr.* 1994; 14:99-129.
- [19] Hwang C-S, Loftus TM, Mandrup S, Lane MD. ADIPOCYTE DIFFERENTIATION AND LEPTIN EXPRESSION. *Annual Review of Cell and Developmental Biology.* 1997;13(1):231-59.

- [20] Rosen ED, Spiegelman BM. Molecular regulation of adipogenesis. *Annu Rev Cell Dev Biol.* 2000;16:145-71.
- [21] Ashammakhi N, Reis R, Chiellini F. Topics in Tissue Engineering, Vol. 4. Eds. 2008
- [22] Rosen ED, Walkey CJ, Puigserver P, Spiegelman BM. Transcriptional regulation of adipogenesis. *Genes Dev.* 2000 Jun 1;14(11):1293-307.
- [23] Darlington GJ, Ross SE, MacDougald OA. The role of C/EBP genes in adipocyte differentiation. *The Journal of biological chemistry.* 1998;273(46):30057-60.
- [24] Tang QQ, Lane MD. Activation and centromeric localization of CCAAT/enhancer-binding proteins during the mitotic clonal expansion of adipocyte differentiation. *Genes & development.* 1999;13(17):2231-41.
- [25] Zhang JW, Klemm DJ, Vinson C, Lane MD. Role of CREB in transcriptional regulation of CCAAT/enhancer-binding protein beta gene during adipogenesis. *The Journal of biological chemistry.* 2004; 279(6):4471-8.
- [26] Rosen ED, MacDougald OA. Adipocyte differentiation from the inside out. *Nature reviews Molecular cell biology.* 2006; 7(12):885-96.
- [27] Andersen C, Rayalam S, Della-Fera MA, Baile CA. Phytochemicals and adipogenesis. *Biofactors.* 2010 Nov-Dec; 36(6):415-22.
- [28] Huang C, Zhang Y, Gong Z, Sheng X, Li Z, Zhang W, Qin Y. Berberine inhibits 3T3-L1 adipocyte differentiation through the PPARgamma pathway. *BiochemBiophys Res Commun.* 2006 Sep 22; 348(2):571-8.
- [29] Moon HS, Lee HG, Choi YJ, Kim TG, Cho CS. Proposed mechanisms of (-)-epigallocatechin-3-gallate for anti-obesity. *ChemBiol Interact.* 2007 Apr 25; 167(2):85-98.

- [30] Kim D, Han GD. Ameliorating Effects of Fermented Rice Bran Extract on Oxidative Stress Induced by High Glucose and Hydrogen Peroxide in 3T3-L1 Adipocytes. *Plant Foods Hum Nutr.* 2011 Jul 12.
- [31] Xu H, Barnes GT, Yang Q, Tan G, Yang D, et al. Chronic inflammation in fat plays a crucial role in the development of obesity-related insulin resistance. *J Clin Invest.* 2003; 112(12):1821-30.
- [32] Weisberg SP, McCann D, Desai M, Rosenbaum M, Leibel RL, Ferrante AW Jr. Obesity is associated with macrophage accumulation in adipose tissue. *J Clin Invest.* 2003 Dec; 112(12):1796-808.
- [33] De Souza CT, Araujo EP, Bordin S, Ashimine R, Zollner RL, et al. Consumption of a fat-rich diet activates a proinflammatory response and induces insulin resistance in the hypothalamus. *Endocrinology.* 2005; 146:4192–99.
- [34] Hotamisligil GS, Shargill NS, Spiegelman BM. Adipose expression of tumor necrosis factor- α : direct role in obesity-linked insulin resistance. *Science.* 1993; 259:87–91.
- [35] Cai D, Yuan M, Frantz DF, Melendez PA, Hansen L, et al. Local and systemic insulin resistance resulting from hepatic activation of IKK- β and NF- κ B. *Nat Med.* 2005; 11:183–90.
- [36] Ehses JA, Perren A, Eppler E, Ribaux P, Pospisilik JA, et al. Increased number of islet-associated macrophages in type 2 diabetes. *Diabetes.* 2007; 56:2356–70.
- [37] Saghizadeh M, Ong JM, Garvey WT, Henry RR, Kern PA. The expression of TNF α by human muscle. Relationship to insulin resistance. *J Clin Investig.* 1996; 97:1111–16.
- [38] Gregor MF, Hotamisligil GS. Inflammatory mechanisms in obesity. *Annu Rev Immunol.* 2011; 29:415-45. Review.

- [39] Solinas G, Karin M. JNK1 and IKK β : molecular links between obesity and metabolic dysfunction. *FASEB J.* 2010; 24:2596–611.
- [40] Aljada A, Mohanty P, Ghanim H, Abdo T, Tripathy D, et al. Increase in intranuclear nuclear factor κ B and decrease in inhibitor κ B in mononuclear cells after a mixed meal: evidence for a proinflammatory effect. *Am J Clin Nutr.* 2004; 79:682–90.
- [41] Bashan N, Dorfman K, Tarnovscki T, Harman-Boehm I, Liberty IF, et al. Mitogen-activated protein kinases, inhibitory- κ B kinase, and insulin signaling in human omental versus subcutaneous adipose tissue in obesity. *Endocrinology.* 2007; 148:2955–62.
- [42] Olefsky JM, Glass CK. Macrophages, inflammation, and insulin resistance. *Annu Rev Physiol.* 2010; 72:219–46.
- [43] Liu J, Divoux A, Sun J, Zhang J, Clement K, et al. Genetic deficiency and pharmacological stabilization of mast cells reduce diet-induced obesity and diabetes in mice. *Nat Med.* 2009; 15:940–45.
- [44] Ohmura K, Ishimori N, Ohmura Y, Tokuhara S, Nozawa A, et al. Natural killer T cells are involved in adipose tissue inflammation and glucose intolerance in diet-induced obese mice. *Arterioscler Thromb Vasc Biol.* 2010; 30:193–99.
- [45] Feuerer M, Herrero L, Cipolletta D, Naaz A, Wong J, et al. Lean, but not obese, fat is enriched for a unique population of regulatory T cells that affect metabolic parameters. *Nat Med.* 2009; 15:930–39.
- [46] Nishimura S, Manabe I, Nagasaki M, Eto K, Yamashita H, et al. CD8+ effector T cells contribute to macrophage recruitment and adipose tissue inflammation in obesity. *Nat Med.* 2009; 15:914–20.
- [47] Pradhan AD, Manson JE, Rifai N, Buring JE, Ridker PM. C-reactive protein, interleukin 6, and risk of developing type 2 diabetes mellitus. *JAMA.* 2001; 286:327–34.

- [48] Vozarova B, Weyer C, Lindsay RS, Pratley RE, Bogardus C, Tataranni PA. High white blood cell count is associated with a worsening of insulin sensitivity and predicts the development of type 2 diabetes. *Diabetes*. 2002; 51:455–61.
- [49] Emilsson V, Thorleifsson G, Zhang B, Leonardson AS, Zink F, et al. Genetics of gene expression and its effect on disease. *Nature*. 2008; 452:423–28.
- [50] Stephens JM, Lee J, Pilch PF. Tumor necrosis factor- α -induced insulin resistance in 3T3-L1 adipocytes is accompanied by a loss of insulin receptor substrate-1 and GLUT4 expression without a loss of insulin receptor-mediated signal transduction. *J Biol Chem*. 1997; 272:971–76.
- [51] Engelman JA, Berg AH, Lewis RY, Lisanti MP, Scherer PE. Tumor necrosis factor α -mediated insulin resistance, but not dedifferentiation, is abrogated by MEK1/2 inhibitors in 3T3-L1 adipocytes. *Mol Endocrinol*. 2000; 14:1557–69.
- [52] Boura-Halfon S, Zick Y. Phosphorylation of IRS proteins, insulin action, and insulin resistance. *Am J Physiol Endocrinol Metab*. 2009; 296:E581–91.
- [53] Hotamisligil GS. Endoplasmic reticulum stress and the inflammatory basis of metabolic disease. *Cell*. 2010; 140:900–17.
- [54] Feingold KR, Doerrler W, Dinarello CA, Fiers W, Grunfeld C. 1992. Stimulation of lipolysis in cultured fat cells by tumor necrosis factor, interleukin-1, and the interferons is blocked by inhibition of prostaglandin synthesis. *Endocrinology*. 130:10–16.
- [55] Wall R, Ross RP, Fitzgerald GF, Stanton C. Fatty acids from fish: the anti-inflammatory potential of long-chain omega-3 fatty acids. *Nutr Rev*. 2010; 68:280–89.
- [56] Todoric J, Loffler M, Huber J, Bilban M, Reimers M, et al. Adipose tissue inflammation induced by high-fat diet in obese diabetic mice is prevented by n-3 polyunsaturated fatty acids. *Diabetologia* 2006; 49:2109–19.

- [57] Hu FB, Cho E, Rexrode KM, Albert CM, Manson JE. Fish and long-chain omega-3 fatty acid intake and risk of coronary heart disease and total mortality in diabetic women. *Circulation* 2003; 107:1852–57.
- [58] Tanaphat Rungwattanapong, Kanokwan Sringarm and Dumnern Karladee. Total Phenolic Compound in Glutinous Purple Rice Grains and Relation to Phenotype Characteristics, *Agricultural Science Journal* 42: 1 (Suppl.): 417-420 (2011).
- [59] Dumnern Karladee and Sunsanee Jamjod. 2007. Genotypic Variation in Grain Nutrition in Local Purple Rice Genotypes, The 2nd International Conference on Rice for the Future 2007, 5 - 9 November 2007, Queen Sirikit National Convention Center Bangkok, Thailand, Page 138-140.
- [60] Natsima Tokhun, RD6, <http://www.natsima.com/agriculture-environment/rd-6-rice/>, 17June2014
- [61] Fung TT, Hu FB, Pereira MA, Liu S, Stampfer MJ, Colditz GA, et al. Whole-grain intake and the risk of type 2 diabetes: a prospective study in men. *The American journal of clinical nutrition*. 2002; 76(3):535-40.
- [62] Zhang G, Pan A, Zong G, Yu Z, Wu H, Chen X, et al. Substituting white rice with brown rice for 16 weeks does not substantially affect metabolic risk factors in middle-aged Chinese men and women with diabetes or a high risk for diabetes. *The Journal of nutrition*. 2011; 141(9):1685-90.
- [63] Qureshi AA, Sami SA, Khan FA. Effects of stabilized rice bran, its soluble and fiber fractions on blood glucose levels and serum lipid parameters in humans with diabetes mellitus Types I and II. *The Journal of nutritional biochemistry*. 2002; 13(3):175-87.
- [64] Juliana, B.O, Rice in human nutrition, 1993, 35-59, 92-5-103149-5
- [65] Kozuka C, Yabiku K, Sunagawa S, Ueda R, Taira S, Ohshiro H, et al. Brown rice and its component, gamma-oryzanol, attenuate the preference for high-fat diet by decreasing hypothalamic endoplasmic reticulum stress in mice. *Diabetes*. 2012; 61(12):3084-93.

- [66] Lerma-García MJ, Herrero-Martínez JM, Simó-Alfonso EF, Mendonça CRB, Ramis-Ramos G. Composition, industrial processing and applications of rice bran γ -oryzanol. *Food Chemistry*. 2009; 115(2):389-404.
- [67] Jin Son M, W. Rico C, Hyun Nam S, Young Kang M. Influence of Oryzanol and Ferulic Acid on the Lipid Metabolism and Antioxidative Status in High Fat-Fed Mice. *Journal of Clinical Biochemistry and Nutrition*. 2010; 46(2):150-6.
- [68] Imam M, Ismail M, Ithnin H, Tubesha Z, Omar A. Effects of Germinated Brown Rice and Its Bioactive Compounds on the Expression of the Peroxisome Proliferator-Activated Receptor Gamma Gene. *Nutrients*. 2013; 5(2):468-77.
- [69] Ghosh D, Konishi T. Anthocyanins and anthocyanin-rich extracts: role in diabetes and eye function. *Asia Pacific journal of clinical nutrition*. 2007; 16(2):200-8.
- [70] Kowalczyk E, Krzesiński P, Kura M, Szmigiel B, Błaszczyk J. Anthocyanins in medicine. *Pol J Pharmacol*. 2003; 55: 699–702.
- [71] Ghosh D and Konishi T. Anthocyanins and anthocyanin-rich extracts: role in diabetes and eye function. *Asia Pac J Clin Nutr* 2007; 16 (2):200-208.
- [72] Tsuda T, Horio F, Uchida K, Aoki H, Osawa T. Dietary cyanidin 3-O-beta-D-glucoside-rich purple corn color prevents obesity and ameliorates hyperglycemia in mice. *J Nutr* 2003; 133: 2125-30.
- [73] Tsuda T, Ueno Y, Aoki H, Koda T, Horio F, et al. Anthocyanin enhances adipocytokine secretion and adipocyte-specific gene expression in isolated rat adipocytes. *Biochem Biophys Res Commun*. 2004; 316:149-57.
- [74] Min SW, Ryu SN, Kim DH. Anti-inflammatory effects of black rice, cyanidin-3-O-beta-Dglycoside, and its metabolites, cyanidin and protocatechuic acid. *Int Immunopharmacol*. 2010 Aug; 10(8):959-66.
- [75] Murakami A, Nishizawa T, Egawa K, Kawada T, Nishikawa Y, et al. New class of linoleic acid metabolites biosynthesized by corn and rice lipoxygenases:


- suppression of proinflammatory mediator expression via attenuation of MAPK- and Akt-, but not PPARgamma-, dependent pathways in stimulated macrophages. *Biochem Pharmacol.* 2005 Nov 1; 70 (9):1330-42.
- [76] Lee SH, Sohn YS, Kang KK, Kwon JW, Yoo M. Inhibitory effect of DA-9201, an extract of *Oryza sativa* L., on airway inflammation and bronchial hyperresponsiveness in mouse asthma model. *Biol Pharm Bull.* 2006 Jun; 29(6):1148-53.
- [77] Chacón MR, Ceperuelo-Mallafré V, Maymó-Masip E, Mateo-Sanz JM, Arola L, et al. Grapeseed procyanidins modulate inflammation on human differentiated adipocytes in vitro. *Cytokine.* 2009; 47:137–142.
- [78] Yin LM, Wei Y, Wang Y, Xu YD, Yang YQ. Long term and standard incubations of WST-1 reagent reflect the same inhibitory trend of cell viability in rat airway smooth muscle cells. *International journal of medical sciences.* 2013; 10(1):68-72.
- [79] Livak KJ, Schmittgen TD. Analysis of relative gene expression data using real-time quantitative PCR and the 2(-Delta Delta C (T)) Method. *Methods (San Diego, Calif).* 2001; 25(4):402-8.
- [80] Hotamisligil GS, Arner P, Caro JF, Atkinson RL, Spiegelman BM. Increased adipose tissue expression of tumor necrosis factor-alpha in human obesity and insulin resistance. *The Journal of clinical investigation.* 1995; 95(5):2409-15.
- [81] Zhang G, Pan A, Zong G, Yu Z, Wu H, Chen X, et al. Substituting white rice with brown rice for 16 weeks does not substantially affect metabolic risk factors in middle-aged Chinese men and women with diabetes or a high risk for diabetes. *The Journal of nutrition.* 2011; 141(9):1685-90.
- [82] Fung TT, Hu FB, Pereira MA, Liu S, Stampfer MJ, Colditz GA, et al. Whole-grain intake and the risk of type 2 diabetes: a prospective study in men. *The American journal of clinical nutrition.* 2002;76(3):535-40.

- [83] Leardkamolkarn V, Thongthe W, Suttiarporn P, Kongkachuchai R, Wongpornchai S, Wanavijitr A. Chemopreventive properties of the bran extracted from a newly-developed Thai rice: The Riceberry. *Food Chemistry*. 2011; 125(3):978-85.
- [84] Lee B, Lee M, Lefevre M, Kim HR. Anthocyanins inhibit lipogenesis during adipocyte differentiation of 3T3-L1 preadipocytes. *Plant foods for human nutrition* (Dordrecht, Netherlands). 2014; 69(2):137-41.
- [85] The USDA national nutrient database for standard reference 2012
- [86] Sangkitikomol W, Tencomnao T, Rocejanasaroj A. Effects of Thai black sticky rice extract on oxidative stress and lipid metabolism gene expression in HepG2 cells. *Genetics and molecular research: GMR*. 2010; 9 (4):2086-95.
- [87] Ryan EP. Bioactive food components and health properties of rice bran. *Journal of the American Veterinary Medical Association*. 2011; 238 (5):593-600.
- [88] Matsunaga T, Shoji A, Gu N, Joo E, Li S, Adachi T, et al. gamma-Tocotrienol attenuates TNF-alpha-induced changes in secretion and gene expression of MCP-1, IL-6 and adiponectin in 3T3-L1 adipocytes. *Molecular medicine reports*. 2012; 5 (4):905-9.
- [89] Kozuka C, Yabiku K, Takayama C, Matsushita M, Shimabukuro M. Natural food science based novel approach toward prevention and treatment of obesity and type 2 diabetes: recent studies on brown rice and gamma-oryzanol. *Obesity research & clinical practice*. 2013; 7 (3):e165-72.
- [90] Madsen L, Petersen RK, Kristiansen K. Regulation of adipocyte differentiation and function by polyunsaturated fatty acids. *Biochimica et Biophysica Acta (BBA) - Molecular Basis of Disease*. 2005; 1740 (2):266-86.
- [91] Jang S, Xu Z. Lipophilic and hydrophilic antioxidants and their antioxidant activities in purple rice bran. *Journal of agricultural and food chemistry*. 2009; 57(3):858-62.

- [92] Eitenmiller, R. R. Vitamin E Content of Fats and Oils: Nutritional Implications. *Food Technol.* 1997; 51(5): 78-81.
- [93] Yu L, Tsao R, Shahidi F. Cereals and Pulses: Nutraceutical Properties and Health Benefits. Wiley-Blackwell. 2012; 328.
- [94] Tanaka T, Kono T, Terasaki F, Yasui K, Soyama A, Otsuka K, et al. Thiamine prevents obesity and obesity-associated metabolic disorders in OLETF rats. *Journal of nutritional science and vitaminology.* 2010; 56(6):335-46.
- [95] Jin Son M, C WR, Hyun Nam S, Young Kang M. Influence of oryzanol and ferulic Acid on the lipid metabolism and antioxidative status in high fat-fed mice. *Journal of clinical biochemistry and nutrition.* 2010; 46(2):150-6.
- [96] Ho JN, Son ME, Lim WC, Lim ST, Cho HY. Anti-obesity effects of germinated brown rice extract through down-regulation of lipogenic genes in high fat diet-induced obese mice. *Bioscience, biotechnology, and biochemistry.* 2012; 76(6):1068-74.
- [97] Kang SW, Kang SI, Shin HS, Yoon SA, Kim JH, Ko HC, et al. Sasa quelpaertensis Nakai extract and its constituent p-coumaric acid inhibit adipogenesis in 3T3-L1 cells through activation of the AMPK pathway. *Food and chemical toxicology: an international journal published for the British Industrial Biological Research Association.* 2013; 59:380-5.
- [98] Moon J, Do HJ, Kim OY, Shin MJ. Antiobesity effects of quercetin-rich onion peel extract on the differentiation of 3T3-L1 preadipocytes and the adipogenesis in high fat-fed rats. *Food and chemical toxicology: an international journal published for the British Industrial Biological Research Association.* 2013; 58:347-54.
- [99] Lee B, Lee M, Lefevre M, Kim HR. Anthocyanins inhibit lipogenesis during adipocyte differentiation of 3T3-L1 preadipocytes. *Plant foods for human nutrition (Dordrecht, Netherlands).* 2014; 69(2):137-41.

- [100] Kim HK, Kim JN, Han SN, Nam JH, Na HN, Ha TJ. Black soybean anthocyanins inhibit adipocyte differentiation in 3T3-L1 cells. *Nutrition research* (New York, NY). 2012; 32(10):770-7.
- [101] Song Y, Park HJ, Kang SN, Jang SH, Lee SJ, Ko YG, et al. Blueberry peel extracts inhibit adipogenesis in 3T3-L1 cells and reduce high-fat diet-induced obesity. *PLoS One*. 2013; 8(7):e69925.
- [102] Uysal KT, Wiesbrock SM, Marino MW, Hotamisligil GS. Protection from obesity-induced insulin resistance in mice lacking TNF-alpha function. *Nature*. 1997; 389(6651):610-4.
- [103] Kozuka C, Yabiku K, Sunagawa S, Ueda R, Taira S, Ohshiro H, et al. Brown rice and its component, gamma-oryzanol, attenuate the preference for high-fat diet by decreasing hypothalamic endoplasmic reticulum stress in mice. *Diabetes*. 2012; 61(12):3084-93.
- [104] Zhao L, Kang I, Fang X, Wang W, Lee MA, Hollins RR, et al. Gamma-tocotrienol attenuates high-fat diet-induced obesity and insulin resistance by inhibiting adipose inflammation and M1 macrophage recruitment. *International journal of obesity* (2005). 2014.
- [105] Lee S-H, Park H-J, Chun H-K, Cho S-Y, Cho S-M, Lillehoj HS. Dietary phytic acid lowers the blood glucose level in diabetic KK mice. *Nutrition Research*. 2006;26(9):474-9.
- [106] Kozuka C, Yabiku K, Sunagawa S, Ueda R, Taira S, Ohshiro H, et al. Brown rice and its component, gamma-oryzanol, attenuate the preference for high-fat diet by decreasing hypothalamic endoplasmic reticulum stress in mice. *Diabetes*. 2012; 61(12):3084-93.
- [107] Choi R, Kim BH, Naowaboot J, Lee MY, Hyun MR, Cho EJ, et al. Effects of ferulic acid on diabetic nephropathy in a rat model of type 2 diabetes. *Experimental & molecular medicine*. 2011; 43(12):676-83.

- [108] Tsuda T, Horio F, Uchida K, Aoki H, Osawa T. Dietary cyanidin 3-O-beta-D-glucoside-rich purple corn color prevents obesity and ameliorates hyperglycemia in mice. *The Journal of nutrition*. 2003; 133(7):2125-30.
- [109] Sasaki R, Nishimura N, Hoshino H, Isa Y, Kadokawa M, Ichi T, et al. Cyanidin 3-glucoside ameliorates hyperglycemia and insulin sensitivity due to downregulation of retinol binding protein 4 expression in diabetic mice. *Biochemical pharmacology*. 2007;74(11):1619-27.



ลิขสิทธิ์มหาวิทยาลัยเชียงใหม่
Copyright© by Chiang Mai University
All rights reserved