REFERENCES

- Acosta, R.M., Neito, A.D., Mena, N.G.V., Vaquera, H.H., Teliz, O.D., Nieto, A.R. and Pichakum, A. 2000. Effect of post-harvest temperatures on the development of internal darkening in mango fruits (*Mangifera indica* L.) cv. Haden and their quality. Acta Horticulturae 509: 401-412.
- Aghdam, M.S., Asghari, M., Moradbeygi, H., Mohammadkhani, N., Mohayeji, M. and Rezapour-Fard, J. 2012. Effect of postharvest salicylic acid treatment on reducing chilling injury in tomato fruit. Romanian Biotechnological Letters 17: 7466-7473.
- Aghdam, M.S. and Bodbodak, S. 2013. Physiological and biochemical mechanisms regulating chilling tolerance in fruits and vegetables under postharvest salicylates and jasmonates treatments. Scientia Horticulturae 156: 73-85.
- Ajila, C.M., Naidu, K.A., Bhat, S.G. and Rao, U.J.S.P. 2007. Bioactive compounds and antioxidant potential of mango peel extract. Food Chemistry 105: 982-988.
- Alférez, F. and Burns, J.K. 2004. Postharvest peel pitting at non-chilling temperatures in grapefruit is promoted by changes from low to high relative humidity during storage. Postharvest Biology and Technology 32: 79-87.
- Ali, M.B., Hahn, E.J. and Paek, K.Y. 2007. Methyl jasmonate and salicylic acid induced oxidative stress and accumulation of phenolics in *Panax ginseng* bioreactor root suspension cultures. Molecules 12: 607-621.
- Al-Qurashi, A.D. 2012. Effect of pre-storage salicylic acid, calcium chloride and 2,4dichlorophenoxyacetic acid dipping on chilling injury and quality of 'Taify' cactus pear fruit during cold storage. African Journal of Biotechnology 11: 6501-6509.

- Al-Qurashi, A.D. and Awad, M.A. 2012. Postharvest salicylic acid treatment reduces chilling injury of 'Taify' cactus pear fruit during cold storage. Journal of Food Agriculture and Environment 10: 120-124.
- Ananieva, E.A., Christov, K.N. and Popova, L.P. 2004. Exogenous treatment with salicylic acid leads to increased antioxidant capacity in leaves of barley plants exposed to paraquat. Journal of Plant Physiology 161: 319-328.
- AOAC. 1990. Official Methods of Analysis of AOAC International: Current Through March 1990 Supplement. Washington, D.C.
- Apel, K. and Hirt, H. 2004. Reactive oxygen species: metabolism, oxidative stress, and signal transduction. Annual Review of Plant Biology 55: 373-399.
- Arora, A., Sairam, R.K. and Srivastava, G.C. 2002. Oxidative stress and antioxidative system in plant. Current Science 82: 1227-1238.
- Asada, K. 1992. Ascorbate peroxidase: a hydrogen peroxide-scavenging enzyme in plants. Physiologia Plantarum 85: 235-241.
- Baiyewu, R.A. and Amusa, N.A. 2005. The effect of temperature and relative humidity on pawpaw fruit rot in south-western Nigeria. World Journal of Agricultural Sciences 1: 80-83.
- Bhattacharjee, S. 2012. The language of reactive oxygen species signaling in plants. Journal of Botany 2012: 1-22.
- Bloknina, O., Virolainen, E. and Fagerstedt, K.V. 2003. Antioxidant, oxidative damage and oxygen deprivation stress: a review. Annals of Botany 91: 179-194.
- Cai, C., Li, X. and Chen, K.S. 2006. Acetylsalicylic acid alleviates chilling injury of postharvest loquat (*Eriobotrya japonica* Lindl.) fruit. European Food Research and Technology 223: 533-539.

- Cai, Y., Cao, S., Yang, Z. and Zheng, Y. 2011. MeJA regulates enzymes involved in ascorbic acid and glutathione metabolism and improves chilling tolerance in loquat fruit. Postharvest Biology and Technology 59: 324-326.
- Camelo, A.F.L. 2004. Manual for the preparation and sale of fruits and vegetables. From field to market. FAO Agricultural Services Bulletin 151: 47-51.
- Cao, S., Yang, Z., Cai, Y. and Zheng, Y. 2011. Fatty acid composition and antioxidant system in relation to susceptibility of loquat fruit to chilling injury. Food Chemistry 127: 1777-1783.
- Cao, S., Zheng, Y., Wang, K., Jin, P. and Rui, H. 2009. Methyl jasmonate reduces chilling injury and enhances antioxidant enzyme activity in postharvest loquat fruit. Food Chemistry 115: 1458-1463.
- Cao, H., Zheng, Y., Wang, K., Rui, H. and Tang, S. 2010. Effect of methyl jasmonate on cell wall modification of loquat fruit in relation to chilling injury after harvest. Food Chemistry 118: 641-647.
- Chan, H.T., Sanxter, T.S. and Couey, H.M. 1985. Electrolyte leakage and ethylene production induced chilling injury of papayas. HortScience 20: 1070-1072.
- Chen, S., Zimei, L., Cui, J., Jiangang, D., Xia, X., Liu, D. and Yu, J. 2011. Alleviation of chilling-induced oxidative damage by salicylic acid pretreatment and related gene expression in eggplant seedlings. Plant Growth Regulation 65: 101-108.
- Chen, Z., Silva, H. and Klessig, D.F. 1993. Active oxygen species in the induction of plant systemic acquired resistance by salicylic acid. Science 262: 1883-1886.
- Cheong, J.J. and Choi, Y.D. 2003. Methyl jasmonate as a vital substance in plants. Trends in Genetics 19: 409-413

- Chidtragool, S., Ketsa, S., Bowen, J., Ferguson, I.B. and van Doorn, W.G. 2011. Chilling injury in mango fruit peel: cultivar differences are related to the activity of phenylalanine ammonia lyase. Postharvest Biology and Technology 62: 59-63.
- Chongchatuporn, U., Ketsa, S. and van Doorn, W.G. 2013. Chilling injury in mango (*Mangifera indica*) fruit peel: relationship with ascorbic acid concentrations and antioxidant enzyme activities. Postharvest Biology and Technology 86: 409-417.
- Conrath, U., Chen, Z., Ricigliano, J.R. and Klessig, D.F. 1995. Two inducers of plant defense responses, 2,6-dichloroisonicotinec acid and salicylic acid, inhibit catalase activity in tobacco. Proceedings of the National Academy of Sciences of the United States of America 92: 7143-7147.
- Côté, F., Thompson, J.E. and Willemot, C. 1993. Limitation to the use of electrolyte leakage for the measurement of chilling injury in tomato fruit. Postharvest Biology and Technology 3: 103-110.
- Creelman, R.A. and Mullet, J.E. 1997. Oligosaccharins, brassinolides, and jasmonates: nontraditional regulators of plant growth, development, and gene expression. The Plant Cell 9: 1211-1223.
- Cruz de Carvalho, M.H. 2008. Drought stress and reactive oxygen species. Plant Signaling and Behavior 3: 156-165.
- Dat, J., Vandenabeele, S., Montagu, M.V. and Breusegem, F.V. 2000. Dual action of the active oxygen species during plant stress responses. Cellular and Molecular Life Sciences 57: 779-795.
- Deepa, N., Kaur, C., Singh, B. and Kapoor, H.C. 2006. Antioxidant activity in some red sweet pepper cultivars. Journal of Food Composition and Analysis 19: 572-578.
- Department of Agriculture. 2011. Storage of fruits and vegetables. The Agriculture Cooperative Federation of Thailand, Limited. Bangkok, 13-14, 40 (in Thai).

- Ding, C.K., Wang, C.Y., Gross, K.C. and Smith, D.L. 2001. Reduction of chilling injury and transcript accumulation of heat shock proteins in tomato fruit by methyl jasmonate and methyl salicylate. Plant Science 161: 1153-1159.
- Ding, C.K., Wang, C.Y., Gross, K.C. and Smith, D.L. 2002. Jasmonate and salicylate induce the expression of pathogenesis-related-protein genes and increase resistance to chilling injury in tomato fruit. Planta 214: 895-901.
- Ding, Z.S., Tian, S.P., Zheng, X.L., Zhou, Z.W. and Xu, Y. 2007. Responses of reactive oxygen metabolism and quality in mango fruit to exogenous oxalic acid or salicylic acid under chilling temperature stress. Physiologia Plantarum 130: 112-121.
- Dokhanieh, A.Y., Aghdam, M.S., Fard, J.R. and Hassanpour, H. 2013. Postharvest salicylic acid treatment enhances antioxidant potential of cornelian cherry fruit. Scientia Horticulturae 154: 31-36.
- Duan, X., Su, X., You, Y., Qu, H., Li, Y. and Jiang, Y. 2007. Effect of nitric oxide on pericarp browning of harvested longan fruit in relation to phenolic metabolism. Food Chemistry 104: 571-576.
- Durner, J. and Klessig, D.F. 1995. Inhibition of ascorbate peroxidase by salicylic acid and 2,6-dichloroisonicotinic acid, two inducers of plant defense responses. Proceedings of the National Academy of Sciences of the United States of America 92: 11312-11316.
- Evans, E.A. 2008. Recent trends in world and U.S. mango production, trade, and consumption. University of Florida, Gainesville. IFAS Extension, Document FE718. http://edis.ifas.ufl.edu.
- Foyer, C.H. and Noctor, G. 2003. Redox sensing and signalling associated with reactive oxygen in chloroplasts, peroxisomes and mitochondria. Physiologia Plantarum 119: 355-364.

- Foyer, C.H. and Noctor, G. 2005. Oxidant and antioxidant signaling in plants: a reevaluation of the concept of oxidative stress in a physiological context. Plant, Cell and Environment 28: 1056-1071.
- Freitas, S.T. and Mitcham, E.J. 2013. Quality of pitaya fruit (*Hylocereus undatus*) as influenced by storage temperature and packaging. Scientia Agricola 70: 257-262.
- Fung, R.W.M., Wang, C.Y., Smith, D.L., Gross, K.C. and Tian, M. 2004. MeSA and MeJA increase steady-state transcript levels of alternative oxidase and resistance against chilling injury in sweet peppers (*Capsicum annuum* L.). Plant Science 166: 711-719.
- Fung, R.W.M., Wang, C.Y., Smith, D.L., Gross, K.C., Tao, Y. and Tian, M. 2006. Characterization of alternative oxidase (AOX) gene expression in response to methyl salicylate and methyl jasmonate pre-treatment and low temperature in tomatoes. Journal of Plant Physiology 163: 1049-1060.
- Gallie, D.R. 2013. L-ascorbic acid: a multifunctional molecule supporting plant growth and development. Scientifica 2013: article ID 795964, 24 pp.
- Gill, S.S. and Tuteja, N. 2010. Reactive oxygen species and antioxidant machinery in abiotic stress tolerance in crop plants. Plant Physiology and Biochemistry 48: 909-930.
- González-Aguilar, G.A., Buta, J.G. and Wang, C.Y. 2001. Methyl jasmonate reduces chilling injury symptoms and enhances colour development of 'Kent' mangoes. Journal of the Science of Food and Agriculture 81: 1244-1249.
- González-Aguilar, G.A., Fortiz, J., Cruz, R., Baez, R. and Wang, C.Y. 2000. Methyl jasmonate reduces chilling injury and maintains postharvest quality of mango fruit. Journal of Agricultural and Food Chemistry 48: 515-519.

- González-Aguilar, G.A., Tiznado-Hernández, M.E., Zavaleta-Gatica, R. and Martínez-Téllez, M.A. 2004. Methyl jasmonate treatments reduce chilling injury and activate the defense response of guava fruits. Biochemical and Biophysical Research Communications 313: 694-701.
- Graham, D. and Paterson, B.D. 1982. Responses plants to low, nonfreezing temperatures: proteins, metabolism and acclimation. Annual Review of Plant Physiology 33: 347-372.
- Gronwald, J.W., Fuerst, E.P., Eberlein, C.V. and Egli, M.A. 1987. Effect of herbicide antidotes on glutathione content and glutathione S-transferase activity of sorghum shoots. Pesticide Biochemistry and Physiology 29: 66-76.
- Han, J., Tian, S.P., Meng, X.H. and Ding, Z.S. 2006. Response of physiologic metabolism and cell structures in mango fruit to exogenous methyl salicylate under low-temperature stress. Physiologia Plantarum 128: 125-133.
- Harfouche, A.L., Rugini, E., Mencarelli, F., Botondi, R. and Muleo, R. 2008. Salicylic acid induces H₂O₂ production and endochitinase gene expression but not ethylene biosynthesis in *Castanea sativa in vitro* model system. Journal of Plant Physiology 165: 734-744.
- Hayat, S. and Ahmad, A. 2007. Salicylic Acid: A Plant Hormone. Springer. Dordrecht. 401 pp.
- Hayat, Q., Hayat, S., Irfan, M. and Ahmad, A. 2010. Effect of exogenous salicylic acid under changing environment: a review. Environmental and Experimental Botany 68: 14-25.
- He, Y., Lui, Y., Cao, W., Huai, M., Xu, B. and Huang, B. 2005. Effects of salicylic acid on heat tolerance associated with antioxidant metabolism in kentucky bluegrass. Crop Science 45: 988-995.

- Hodges, D.M., DeLong, J.M., Forney, C.F. and Prange, R.K. 1999. Improving the thiobarbituric acid-reactive-substances assay for estimating lipid peroxidation in plant tissues containing anthocyanin and other interfering compounds. Planta 207: 604 -611.
- Hodges, D.M., Lester, G.E., Munro, K.D. and Toivonen, P.M.A. 2004. Oxidative stress: importance for postharvest quality. HortScience 39: 924-929.
- Hu, X., Li, W., Chen, Q. and Yang, Y. 2009. Early signal transduction linking the synthesis of jasmonic acid in plant. Plant Signaling and Behavior 4: 696-697.
- Huang, D., Ou, B. and Prior, R.L. 2005. The Chemistry behind antioxidant capacity assays. Journal of Agricultural and Food Chemistry 53: 1841-1856.
- Imahori, Y., Takemura, M. and Bai, J. 2008. Chilling-induced oxidative stress and antioxidant responses in mume (*Prunus mume*) fruit during low temperature storage. Postharvest Biology and Technology 49: 54-60.
- Imsabai, W., Ketsa, S. and van Doorn, W.G. 2002. Effect of temperature on softening and the activities of polygalacturonase and pectinesterase in durian fruit. Postharvest Biology and Technology 26: 347-351.
- Janda, T., Szalai, G., Tari, I. and Páldi, E. 1999. Hydroponic treatment with salicylic acid decreases the effects of chilling injury in maize (*Zea mays L.*) plants. Planta 208: 175-180.
- Jiang, L., Hou, T., Yuan, X., Jiang, J. and Yu, Z. 2010. Effect of storage temperature and packaging method on the decay and physiology of fresh leaves of *Gynura bicolor* DC. Journal of Food Processing and Preservation 34: 858-871.
- Jin, P., Wang, K., Shang, H., Tong, J. and Zheng, Y. 2009a. Low-temperature conditioning combined with methyl jasmonate treatment reduces chilling injury of peach fruit. Journal of the Science of Food and Agriculture 89: 1690-1696.

- Jin, P., Zheng, Y., Tang, S., Rui, H. and Wang, C.Y. 2009b. A combination of hot air and methyl jasmonate vapor treatment alleviates chilling injury of peach fruit. Postharvest Biology and Technology 52: 24-29.
- Jin, P., Zhu, H., Wang, J., Chen, J., Wang, X. and Zheng, Y., 2013. Effect of methyl jasmonate on energy metabolism in peach fruit during chilling stress. Journal of the Science of Food and Agriculture 93: 1827-1832.
- Johnson, G.I. and Coates, L.M. 1993. Postharvest diseases of mango. Postharvest News and Information 4: 27-34.
- Kami, D., Muro, T. and Sugiyama, K. 2011. Changes in starch and soluble sugar concentrations in winter squash mesocarp during storage at different temperatures. Scientia Horticulturae 127: 444-446.
- Kane, O., Boulet, M. and Castaigne, F. 1982. Effect of chilling injury on texture and fungal rot of mangoes (*Mangifera indica* L.). Journal of Food Science 47: 992-995.
- Kang, G., Wang, C., Sun, G. and Wang, Z. 2003. Salicylic acid changes activities of H₂O₂-metabolizing enzymes and increases the chilling tolerance of banana seedlings. Environmental and Experimental Botany 50: 9-15.
- Kang, H.M. and Saltveit, M.E. 2002. Chilling tolerance of maize, cucumber and rice seedling leaves and roots are differentially affected by salicylic acid. Physiologia Plantarum 115: 571-576.
- Kaniuga, Z. 2008. Chilling response of plants: importance of galactolipase, free fatty acids and free radicals. Plant Biology 10: 171-184.
- Ketsa, S. and Chidtragool, S. 2005. Changes of phenolic content and activities of phenylalanine ammonia lyase and polyphenol oxidase in relation to discoloration of mango fruit during low temperature storage. Acta Horticulturae 687: 343-345.

- Ketsa, S., Chidtragool, S. and Lurie, S. 2000. Prestorage heat treatment and poststorage quality of mango fruit. HortScience 35: 247-249.
- Ketsa, S., Chidtragool, S., Klein, J.D. and Lurie, S. 1998. Effect of heat treatment on changes in softening, pectic substances and activities of polygalacturonase, pectinesterase and β-galactosidase of ripening mango. Journal of Plant Physiology 153: 457-461.
- Khokon, A.R., Okuma, E., Hossain, M.A., Munemasa, S., Uraji, M., Nakamura, Y., Mori, I.C. and Murata, Y. 2011. Involvement of extracellular oxidative burst in salicylic acid induced stomatal closure in Arabidopsis. Plant, Cell and Environment 34: 434-443.
- Kumpoun, W. and Uthaibutra, J. 2010. Storage life extension of exported 'Nam Dok Mai' mango by refrigerated modified atmosphere packing. Acta Horticulturae 876: 221-226.
- Lei, T., Feng, H., Sun, X. and Dai, Q.L. 2010. The alternative pathway in cucumber seedlings under low temperature stress was enhanced by salicylic acid. Plant Growth Regulation 60: 35-42.
- Lederman, I.E., Zauberman, G., Weksler, A., Rot, I. and Fuchs, Y. 1997. Ethyleneforming capacity during cold storage and chilling injury development in 'Keitt' mango fruit. Postharvest Biology and Technology 10: 107-112.
- Lim, C.S., Kang, S.M., Cho, J.L. and Gross, K.C. 2009. Antioxidizing enzyme activities in chilling-sensitive and chilling-tolerant pepper fruit as affected by stage of ripeness and storage temperature. Journal of the American Society for Horticultural Science 134: 156-163.
- Lim, C.S., Kang, S.M., Cho, J.L., Gross, K.C. and Woolf, A.B. 2007. Bell pepper (*Capsicum annuum* L.) fruits are susceptible to chilling injury at the breaker stage of ripeness. HortScience 42: 1659-1664.

- Lowry, O.H., Rosebrough, N.J., Farr, A.L. and Randall, R.J. 1951. Protein measurement with the folin phenol reagent. The Journal of Biological Chemistry 193: 265-275.
- Luengwilai, K. and Beckles, D.M. 2013. Effect of low temperature storage on fruit physiology and carbohydrate accumulation in tomato ripening-inhibited mutants. Journal of Stored Products and Postharvest Research 4: 35-43.
- Luo, Z., Chen, C. and Xie, J. 2011. Effect of salicylic acid treatment on alleviating postharvest chilling injury of 'Qingnai' plum fruit. Postharvest Biology and Technology 62: 115-120.
- Luo, Z., Wu, X., Xie, Y. and Chen, C. 2012. Alleviation of chilling injury and browning of postharvest bamboo shoot by salicylic acid treatment. Food Chemistry 131: 456-461.
- Lynch, D. and Thompson, J.E. 1984. Lipoxygenase-mediated production of superoxide anion in senescing plant tissue. Federation of European Biochemical Societies Letters 173: 251-254.
- Lyons, J.M. 1973. Chilling injury in plants. Annual Review of Plant Physiology 24: 445-466.
- Mahajan, S. and Tuteja, N. 2005. Cold, salinity and drought stresses: an overview. Archives of Biochemistry and Biophysics 444: 139-158.
- Manthe, B., Schulz, M. and Schnabl, H. 1992. Effects of salicylic acid on growth and stomatal movements of *Vicia faba* L.: evidence for salicylic acid metabolization. Journal of Chemical Ecology 18: 1525-1539.
- Marangoni, A.G., Palma, T. and Stanley, D.W. 1996. Membrane effects in postharvest physiology. Postharvest Biology and Technology 7: 193-217.
- Martinez, M.V. and Whitaker, J.R. 1995. The biochemistry and control of enzymatic browning. Trends in Food Science Technology 6: 195-200.

- Martysiak-Żurowska, D. and Wenta, W. 2012. A comparison of ABTS and DPPH methods for assessing the total antioxidant capacity of human milk. Acta Scientiarum Polonorum Technologia Alimentaria 11: 83-89.
- Mayer, A.M. 1987. Polyphenoloxidases in plants-recent progress. Phytochemistry 26: 11-20.
- Meng, X., Han, J., Wang, Q. and Tian, S. 2009. Changes in physiology and quality of peach fruits treated by methyl jasmonate under low temperature stress. Food Chemistry 114: 1028-1035.
- Millar, A.H., Mittova, V., Kiddle, G., Heazlewood, J.L., Bartoli, C.G., Theodoulou, F.L. and Foyer, C.H. 2003. Control of ascorbate synthesis by respiration and its implications for stress responses. Plant Physiology 133: 443-447.
- Minolta Camera Company Limited. 1990. Precise Color Communication: Color Control from Feeling to Instrumentation. Minolta Camera Company Limited, Osaka. 21 pp.
- Mittler, R. 2002. Oxidative stress, antioxidants and stress tolerance. Trends in Plant Science 7: 1360-1385.
- Miura, K. and Tada, Y. 2014. Regulation of water, salinity, and cold stress responses by salicylic acid. Frontiers in Plant Science 5: 1-12.
- Moreira, X., Sampedro, L. and Zas, R. 2009. Defensive responses of *Pinus pinaster* seedlings to exogenous application of methyl jasmonate: concentration effect and systemic response. Environmental and Experimental Botany 67: 94-100.
- Mori, I.C., Pinontoan, R., Kawano, T. and Muto, S. 2001. Involvement of superoxide generation in salicylic acid-induced stomatal closure in *Vicia faba*. Plant and Cell Physiology 42: 1383-1388.
- Morrelli, K.L., Hess-Pierce, B.M. and Kader, A.A. 2003. Genotypic variation in chilling sensitivity of mature-green bananas and plantains. HortTechnology 13: 328-332.

- Møller, I.M. 2001. Plant mitochondria and oxidative stress: electron transport, NADPH turnover, and metabolism of reactive oxygen species. Annual Review of Plant Physiology and Plant Molecular Biology 52: 561-591.
- Møller, I.M., Jensen, P.E. and Hansson, A. 2007. Oxidative modifications to cellular components in plants. The Annual Review of Plant Biology 58: 459-481.
- Mun'im, A., Negishi, O. and Ozawa, T. 2003. Antioxidant compounds from Crotalaria sessiliflora. Bioscience Biotechnology and Biochemistry 67: 410-414.
- Nair, S., Singh, Z. and Tan, S.C. 2003. Aroma volatiles emission in relation to chilling injury in 'Kensington Pride' mango fruits. The Journal of Horticultural Science and Biotechnology 78: 866-873.
- Nguyen, T.B.T., Ketsa, S. and van Doorn, W.G. 2003. Relationship between browning and the activities of polyphenol oxidase and phenylalanine ammonia lyase in banana peel during low temperature storage. Postharvest Biology and Technology 30: 187-193.
- Nilprapruck, P., Pradisthakarn, N., Authanithee, F. and Keebjan, P. 2008. Effect of exogenous methyl jasmonate on chilling injury and quality of pineapple (*Ananas comosus* L.) cv. Pattavia. Silpakorn University Science and Technology Journal 2: 33-44.
- Noctor, G. and Foyer, C.H. 1998. Ascorbate and glutathione: keeping active oxygen under control. Annual Review of Plant Physiology and Plant Molecular Biology 49: 249-279.
- Office of Agricultural Economics. 2009. Agricultural economics fundamental data 2009. [online]. Available http://www.oae.go.th/download/download_journal/fundamation-2552.pdf (27 August 2010).
- Pala, F.S. and Tabakçioğlu, K. 2007. Free radicals: our enemies or friends? Advances in Molecular Biology 1: 63-69.

- Pantastico, E.B., Lam, P.F., Ketsa, S., Yuniarti and Kosittrakul, M. 1984. Postharvest physiology and storage of mango. In: Mendoza, D.B.Jr. and Wills, R.B.H. (Eds.), Mango: Fruit Development, Postharvest Physiology and Marketing in ASEAN. ASEAN Food Handling Bureau, Kuala Lumpur. p. 39-52.
- Parkin, K.L., Marangoni, A., Jackman, R., Yada, R. and Stanley, D.W. 1989. Chilling injury: a review of possible mechanisms. Journal of Food Biochemistry 13: 127-153.
- Pattanapo, W., Sardsud, V., Noimanee, P., Theanjumpon, P., Pornpasit, R. and Karaboon, S. 2010. Prevention of chilling injury of mango cv. Nam Dok Mai Si Thong by packaging. Agricultural Science Journal 41: 141-144. (in Thai).
- Pérez, A.G., Sanz, C., Richardson, D.G. and Olías, J.M. 1993. Methyl jasmonate vapor promotes β-carotene synthesis and chlorophyll degradation in Golden Delicious apple peel. Journal of Plant Growth Regulation 12: 163-167.
- Pesis, E., Aharoni, D., Aharon, Z., Ben-Arie, R., Aharoni, N. and Fuchs, Y. 2000.Modified atmosphere and modified humidity packaging alleviates chilling injury symptoms in mango fruit. Postharvest Biology and Technology 19: 93-101.
- Petracek, P.D., Wardowski, W.F. and Brown, G.E. 1995. Pitting of grapefruit that resembles chilling injury. HortScience 30: 1422-1426.
- Phakawatmongkol, W., Ketsa, S. and van Doorn, W.G. 2004. Variation in fruit chilling injury among mango cultivars. Postharvest Biology and Technology 32: 115-118.

Pinhero, R.G., Paliyath, G., Yada, R.Y. and Murr, D.P. 1998. Modulation of phospholipase D and lipoxygenase activities during chilling. Relation to chilling tolerance of maize seedlings. Plant Physiology and Biochemistry 36: 213-224.

- Prasad, T.K., Anderson, M.D. and Stewart, C.R. 1994. Acclimation, hydrogen peroxide, and abscisic acid protect mitochondria against irreversible chilling injury in maize seedlings. Plant Physiology 105: 619-627.
- Prior, R.L., Wu, X. and Schaich, K. 2005. Standardized methods for determination of antioxidant capacity and phenolics in foods and dietary supplements. Journal of Agricultural and Food Chemistry 53: 4290-4302.
- Promyou, S., Ketsa, S. and van Doorn, W.G. 2012. Salicylic acid alleviates chilling injury in anthurium (*Anthurium andraeanum* L.) flowers. Postharvest Biology and Technology 64: 104-110.
- Purvis, A.C. and Shewfelt, R.L. 1993. Does the alternative pathway ameliorate chilling injury in sensitive plant tissues? Physiologia Plantarum 88: 712-718.
- Rab, A., Sajid, M., Khan, N.U., Nawab, K., Arif, M. and Khattak, M.K. 2012. Influence of storage temperature on fungal prevalence and quality of citrus fruit (cv. Blood Red). Pakistan Journal of Botany 44: 831-836.
- Rao, T.V.R., Gol, N.B. and Shah, K.K. 2011. Effect of postharvest treatments and storage temperatures on the quality and shelf life of sweet pepper (*Capsicum annum* L.). Scientia Horticulturae 132: 18-26.
- Raskin, I. 1992. Salicylate, a new plant hormone. Plant Physiology 99: 799-803.
- Ribeiro, S.M.R., Barbosa, L.C.A., Queiroz, J.H., Knödler, M. and Schieber, A. 2008. Phenolic compounds and antioxidant capacity of Brazilian mango (*Mangifera indica* L.) varieties. Food Chemistry 110: 620-626.
- Richard, F.F. and Gauillard, F.A. 1997. Oxidation of chlorogenic acid, catechins, and 4-methylcatechol in model solutions by combinations of pear (*Pyrus communis* cv. Williams) polyphenol oxidase and peroxidase: a possible involvement of peroxidase in enzymatic browning. Journal of Agriculture and Food Chemistry 45: 2472-2476.

- Safizadeh, M.R., Rahemi, M. and Aminlari, M. 2007. Effect of postharvest calcium and hot-water dip treatments on catalase, peroxidase and superoxide dismutase in chilled Lisbon lemon fruit. International Journal of Agricultural Research 2: 440-449.
- Sayyari, M., Babalar, M., Kalantari, S., Martínez-Romero, D., Guillén, F., Serrano, M. and Valero, D. 2011a. Vapour treatments with methyl salicylate or methyl jasmonate alleviated chilling injury and enhanced antioxidant potential during postharvest storage of pomegranates. Food Chemistry 124: 964-970.
- Sayyari, M., Babalar, M., Kalantari, S., Serrano, M. and Valero, D. 2009. Effect of salicylic acid treatment on reducing chilling injury in stored pomegranates. Postharvest Biology and Technology 53: 152-154.
- Sayyari, M., Castillo, S., Valero, D., Díaz-Mula, H.M. and Serrano, M. 2011b. Acetyl salicylic acid alleviates chilling injury and maintains nutritive and bioactive compounds and antioxidant activity during postharvest storage of pomegranates.
 Postharvest Biology and Technology 60: 136-142.
- Sevillano, L., Sanchez-Ballesta, M.T., Romojaro, F. and Flores, F.B. 2009.
 Physiological, hormonal and molecular mechanisms regulating chilling injury in horticultural species. Postharvest technologies applied to reduce its impact. Journal of the Science of Food and Agriculture 89: 555-573.
- Shao, H.B., Chu, L.Y., Lu, Z.H. and Kang, C.M. 2008. Primary antioxidant free radical scavenging and redox signaling pathways in higher plant cells. International Journal of Biological Sciences 4: 8-14.
- Sharma, P., Jha, A.B., Dubey, R.S. and Pessarakli, M. 2012. Reactive oxygen species, oxidative damage, and antioxidative defense mechanism in plants under stressful conditions. Journal of Botany 2012: 1-26.
- Shewfelt, R.L. and del Rosario, B.A. 2000. The role of lipid peroxidation in storage disorders of fresh fruits and vegetable. HortScience 35: 575-579.

- Shi, Q., Bao, Z., Zhu, Z., Ying, Q. and Qian, Q. 2006. Effects of different treatments of salicylic acid on heat tolerance, chlorophyll fluorescence, and antioxidant enzyme activity in seedlings of *Cucumis sativa* L. Plant Growth Regulation 48: 127-135.
- Siboza, X.I. and Bertling, I. 2013. The effects of methyl jasmonate and salicylic acid on suppressing the production of reactive oxygen species and increasing chilling tolerance in 'Eureka' lemon [*Citrus limon* (L.) Burm. F.]. Journal of Horticultural Science and Biotechnology 88: 269-276.
- Singleton, V.L. and Rossi, J.R. 1965. Colorimetry of total phenolics with phosphomolybdic-phosphotungstic acid reagents. American Journal of Enology and Viticulture 16: 144-157.
- Slaymaker, D.H., Navarre, D.A., Clark, D., del Pozo, O., Martin, G.B. and Klessig, D.F. 2002. The tobacco salicylic acid-binding protein 3 (SABP3) is the chloroplast carbonic anhydrase, which exhibits antioxidant activity and plays a role in the hypersensitive defense response. Proceedings of the National Academy of Sciences of the United States of America 99: 11640-11645.
- Smirnoff, N. 2000. Ascorbic acid: metabolism and functions of a multifaceted molecule. Current Opinion in Plant Biology 3: 229-235.
- Smirnoff, N. 2005. Antioxidant and Reactive Oxygen Species in Plants. Blackwell Publishing Ltd., Oxford. 302 pp.
- Sofo, A., Rajabpoor, S., Yaghini, H., Shirani, M., Archangi, A., Sangi, S.E., Tavakoli, F., Khodai, M. and Sorkheh, K. 2014. Cold-induced changes in antioxidant defenses and reactive oxygen species in eight wild almond species. Free Radicals and Antioxidants 4: 70-74.
- Sovrano, S., Buiatti, S. and Anese, M. 2006. Influence of malt browning degree on lipoxygenase activity. Food Chemistry 99: 711-717.

- Sunohara, Y. and Matsumoto, H. 2004. Oxidative injury induced by the herbicide quinclorac on *Echinochloa oryzicola* Vasing. and the involvement of antioxidative ability in its highly selective action in grass species. Plant Science 167: 597-606.
- Tareen, M.J., Abbasi, N.A. and Hafiz, I.A. 2012. Postharvest application of salicylic acid enhanced antioxidant enzyme activity and maintained quality of peach cv. 'Flordaking' fruit during storage. Scientia Horticulturae 142: 221–228.
- Taşgín, E., Atící, Ö. And Nalbantoğlu, B. 2003. Effects of salicylic acid and cold on freezing tolerance in winter wheat leaves. Plant Growth Regulation 41: 231-236.
- Thai Customs Department. 2 0 1 4. Import and export statistics. [online]. Available http://www.customs.go.th/Statistic/StatisticResult2550.jsp (2 May 2014).
- Turrens, J.F. 2003. Mitochondrial formation of reactive oxygen species. Journal of Physiology 552: 335-344.
- Vajragupta, O., Boonchoong, P., Boonyasat, J. and Utsintong, M. 2006. Radical Scavenging Agents. P.S Print, Nonthaburi. 200 pp. (in Thai).
- Valpuesta, V. and Botella, M.A. 2004. Biosynthesis of L-ascorbic acid in plants: new pathways for an old antioxidant. Trends in Plant Science 9: 573-577.
- van Rooyen, Z. and Bower, J.P. 2003. The role of fruit mineral composition, phenolic concentration and polyphenol oxidase activity on mesocarp discolouration in 'Pinkerton'. South African Avocado Growers' Association Yearbook 26: 72-75, 77-79, 81-82.
- Velikova, V., Yordanov, I. and Edreva, A. 2000. Oxidation stress and some antioxidant systems in acid rain-treated bean plants: protective role of exogenous polyamines. Plant Science 151: 59-66.

- Villalobos-Acuña, M.G., Biasi, W.V., Mitcham, W.J. and Holcroft, D. 2011. Fruit temperature and ethylene modulate 1-MCP response in 'Bartlett' pears. Postharvest Biology and Technology 60: 17-23.
- Walker, M.A., McKersie, B.O. and Pauls, K.P. 1990. Effect of chilling on the biochemical and functional properties of thylakoid membranes. Plant Physiology 97: 663-669.
- Wang, B., Wang, J., Liang, H., Yi, J., Zhang, J., Lin, L., Wu, Y., Feng, X., Cao, J. and Jiang, W. 2008. Reduced chilling injury in mango fruit by 2,4dichlorophenoxyacetic acid and the antioxidant response. Postharvest Biology and Technology 48: 172-181.
- Wang, C.Y. 1994. Chilling injury of tropical horticultural commodities. Horticultural Science 29: 986-988.
- Wang, C.Y. 1995 Effect of temperature preconditioning on catalase, peroxidase, and superoxide dismutase in chilled zucchini squash. Postharvest Biology and Technology 5: 67-76.
- Wang, C.Y. 1998. Methyl jasmonate inhibits sprouting and improves storage quality of radishes. Postharvest Biology and Technology 14: 179-183.
- Wang, C.Y. 2010. Alleviation of chilling injury in tropical and subtropical fruits. Acta Horticulturae 864: 267-273.
- Wang, C.Y. and Wang, S.Y. 2009. Effect of storage temperatures on fruit quality of various cranberry cultivars. Acta Horticulturae 810: 853-861.
- Wang, K., Jin, P., Cao, S., Shang, H., Yang, Z. and Zheng, Y. 2009. Methyl jasmonate reduces decay and enhances antioxidant capacity in Chinese bayberries. Journal of Agricultural and Food Chemistry 57: 5809-5815.

- Wang, L., Chen, S., Kong, W., Li, S. and Archbold, D.D. 2006. Salicylic acid pretreatment alleviates chilling injury and affects the antioxidant system and heat shock proteins of peaches during cold storage. Postharvest Biology and Technology 41: 244-251.
- Wang, L.J. and Li, S.H. 2006. Salicylic acid-induced heat or cold tolerance in relation to Ca²⁺ homeostasis and antioxidant systems in young grape plants. Plant Science 170: 685-694.
- Wang, Q., Ding, T., Gao, L., Pang, J. and Yang, N. 2012. Effect of brassinolide on chilling injury of green bell pepper in storage. Scientia Horticulturae 144: 195-200.
- Wen, P.F., Chen, J.Y., Kong, W.F., Pan, Q.H., Wan, S.B. and Huang, W.D. 2005. Salicylic acid induced the expression of phenylalanine ammonia-lyase gene in grape berry. Plant Science 169: 928-934.
- Wongsheree, T., Ketsa, S. and van Doorn, E.G. 2009. The relationship between chilling injury and membrane damage in lemon basil (*Ocimum x citriodourum*) leaves.Postharvest Biology and Technology 51: 91-96.
- Wu, M.C., Chen, C.H. and Chen, C.S. 1999. Changes in sugars, starch and amylase activity during ripening of sugar apples at different storage temperatures. Food Preservation Science 25: 57-61.
- Xiong, L., Schumaker, K.S. and Zhu, J.K. 2002. Cell signaling during cold, drought, and salt stress. The Plant Cell 14: 165-183.
- Xu, X. and Tian, S. 2008. Salicylic acid alleviated pathogen-induced oxidative stress in harvested sweet cherry fruit. Postharvest Biology and Technology 49: 379-385.
- Yadav, S.K. 2010. Cold stress tolerance mechanisms in plants: a review. Agronomy for Sustainable Development 30: 515-527.

- Yang, E., Lu, W., Qu, H., Lin, H., Wu, F., Yang, S., Chen, Y. and Jiang, Y. 2009. Altered energy status in pericarp browning of lychee fruit during storage. Pakistan Journal of Botany 41: 2271-2279.
- Yang, H., Wu, F. and Cheng, J. 2011. Reduced chilling injury in cucumber by nitric oxide and the antioxidant response. Food Chemistry 127: 1237-1242.
- Yang, Q., Rao, J., Yi, S., Meng, K., Wu, J. and Hou, Y. 2012. Antioxidant enzyme activity and chilling injury during low-temperature storage of kiwifruit cv. Hongyang exposed to gradual postharvest cooling. Horticulture, Environment and Biotechnology 53: 505-512.
- Yang, Q., Zhang, Z., Rao, J., Wang, Y., Sun, Z., Ma, Q. and Dong, X. 2013. Lowtemperature conditioning induces chilling tolerance in 'Hayward' kiwifruit by enhancing antioxidant enzyme activity and regulating en-dogenous hormones levels. Journal of the Science of Food and Agriculture 93: 3691-3699.
- Yang, S., Su, X., Prasad, K.N., Yang, B., Cheng, G., Chen, Y., Yang, E. and Jiang, Y.2008. Oxidation and peroxidation of postharvest banana fruit during softening.Pakistan Journal of Botany 40: 2023-2029.
- Yu, H. and Goh, C.J. 2000. Identification and characterization of three orchid MADSbox genes of the AP1/AGL9 subfamily during floral transition. Plant Physiology 123: 1325-1336.
- Yuan, S. and Lin, H.H. 2008. Role of salicylic acid in plant abiotic stress. Zeitschrift f
 ür Naturforschung 63: 313-320.
- Zaharah, S.S. and Singh, Z. 2011. Postharvest nitric oxide fumigation alleviates chilling injury, delays fruit ripening and maintains quality in cold-stored 'Kensington Pride' mango. Postharvest Biology and Technology 60: 202-210.

- Zauberman, G. and Jobin-Decor, M.P. 1995. Avocado (*Persea americana* Mill.) quality changes in response to low-temperature storage. Postharvest Biology and Technology 5: 235-243.
- Zeng, K., Cao, J. and Jiang, W. 2006. Enhancing disease resistance in harvested mango (*Mangifera indica* L. cv. 'Matisu') fruit by salicylic acid. Journal of the Science of Food and Agriculture 86: 694-698.
- Zhao, Z., Chen, G. and Zhang, C. 2001. Interaction between reactive oxygen species and nitric oxide in drought-induced abscisic acid synthesis in root tips of wheat seedlings. Australian Journal of Plant Physiology 28: 1055-1061.
- Zhao, Z., Jiang, W., Cao, J., Zhao, Y. and Gu, Y. 2006. Effect of cold-shock treatment on chilling injury in mango (*Mangifera indica* L. cv. 'Wacheng') fruit. Journal of the Science of Food and Agriculture 86: 2458-2462.
- Zheng, Y. and Zhang, Q. 2004. Effects of polyamines and salicylic acid on postharvest storage of 'Ponkan' mandarin. Acta Horticulturae 632: 317-320.
- Zulkifli, K.S., Abdullah, N., Abdullah, A., Aziman, N. and Kamarudin, W.S.S.W. 2012.
 Bioactive phenolic compounds and antioxidant activity of selected fruit peels.
 International Proceedings of Chemical, Biological and Environmental Engineering 49: 66-70.

ลิ<mark>ปสิทธิ์มหาวิทยาลัยเชียงใหม่</mark> Copyright[©] by Chiang Mai University All rights reserved

LIST OF PUBLICATIONS

- Junmatong, C., Uthaibutra, J., Boonyakiat, D., Faiyue, B. and Saengnil, K. 2012. Reduction of chilling injury of 'Nam Dok Mai No. 4' mango fruit by treatments with salicylic acid and methyl jasmonate. Journal of Agricultural Science 4: 126-136.
- Junmatong, C., Chomkitichai, W., Boonyakiat, D., Uthaibutra, J. and Saengnil, K. 2014. Reduction of free radical content and chilling injury in Nam Dok Mai No. 4 mango fruit with methyl jasmonate during low temperature storage. Acta Horticulturae (in press).
- Junmatong, C., Faiyue, B., Rotarayanont, S., Uthaibutra, J., Boonyakiat, D. and Saengnil, K. 2014. Enhancement of enzymatic and non-enzymatic antioxidants of Nam Dok Mai No. 4 mango fruit during cold storage by salicylic acid. Science Asia (in press).

ลิขสิทธิ์มหาวิทยาลัยเชียงใหม่ Copyright[©] by Chiang Mai University All rights reserved