

REFERENCES

Aderibigbe, A.O., Emudianughe, T.S., Lawal, B.S., Evaluation of the antidiabetic action of *Mangifera indica* in mice, *Phytotherapy Research*, 2001, 15(5), 456–458.

Adisakwattana, S., Chantarasinlapin, P., Thammarat, H. and Yibchok-Anun, S., A series of cinnamic acid derivatives and their inhibitory activity on intestinal α -glucosidase, *Journal of Enzyme Inhibition and Medicinal Chemistry*, 2009, 24(5), 1194-1200.

Ahmad, M.I., Ahmad, S. and Moinuddin., Preferential recognition of methylglyoxal-modified calf thymus DNA by circulating antibodies in cancer patients, *Indian Journal of Biochemistry and Biophysics*, 2011, 48(4), 290-296.

Ahmed, N., Advanced glycation endproducts—role in pathology of diabetic complications, *Diabetes Research and Clinical Practice*, 2005, 67(1), 3-21.

Akbar, S., *Andrographis paniculata*: a review of pharmacological activities and clinical effects. *Alternative Medicine Review*, 2011, 16(1), 66-77.

Akhter, F., Khan, M.S., Shahab, U. and Ahmad, S., Bio-Physical characterization of ribose induced glycation: A mechanistic study on DNA perturbations, *International Journal of Biological Macromolecules*, 2013, 58, 206-210.

Al-Amin, Z.M., Thomson M., Al-Qattan, K.K., Peltonen-Shalaby, R. and Ali, M., Anti-diabetic and hypolipidaemic properties of ginger (*Zingiber officinale*) in streptozotocin-induced diabetic rats, *British Journal of Nutrition*, 2006, 96, 660-666.

Ardestani, A. and Yazdanparast, R., Inhibitory effects of ethyl acetate extract of *Teucrium polium* on *in vitro* protein glycoxidation, *Food and Chemical Toxicology*, 2007, 45(12), 2402-2411.

Asolkar, L., Kakkar, K. and Chakre, O., Second supplement to glossary of Indian medicinal plants with active principles Part-I (AK)(1965-1981), Publications and Information directorate (CSIR), Dr. KS Krishan Mar, New Delhi, 1992.

Babizhayev, M.A., Deyev, A.I., Yermakova, V.N., Brikman, I.V. and Bours, J., Lipid peroxidation and cataracts, *Drugs in R & D*, 2004, 5(3), 125-139.

Babu, P.V.A., Sabitha, K.E., Srinivasan, P. and Shyamaladevi, C.S., Green tea attenuates diabetes induced Maillard-type fluorescence and collagen cross-linking in the heart of streptozotocin diabetic rats, *Pharmacological Research*, 2007, 55(5), 433-440.

Barreto, J.C., Trevisan, M.T., Hull, W.E., Erben, G., de Brito, E.S., Pfundstein, B., Würtele, G., Spiegelhalder, B. and Owen, R.W., Characterization and quantitation of polyphenolic compounds in bark, kernel, leaves, and peel of mango (*Mangifera indica* L.), *Journal of Agricultural and Food Chemistry*, 2008, 56(14), 5599-5610.

Beswick, H.T. and Harding, J.J., Conformational changes induced in lens alpha- and gamma-crystallins by modification with glucose 6-phosphate. Implications for cataract, *Biochemical Journal*, 1986, 246, 761-769.

Bischcoff H., Pharmacology of alpha – glucosidase inhibition, *European Journal of Clinical Investigation*, 1994, 24(3), 3-10.

Bolton, W.K., Cattran, D.C., Williams, M.E., Adler, S.G., Appel, G.B., Cartwright, K., Foiles, P.G., Freedman, B.I., Raskin, P. and Ratner, R.E., Randomized trial of an inhibitor of formation of advanced glycation end products in diabetic nephropathy, *American Journal of Nephrology*, 2004, 24(1), 32-40.

Booth, A.A., Khalifah, R.G. and Hudson, B.G., Thiamine pyrophosphate and pyridoxamine inhibit the formation of antigenic advanced glycation end-products: comparison with aminoguanidine, *Biochemical and Biophysical Research Communications*, 1996, 220(1), 113-119.

Booth, A.A., Khalifah, R.G., Todd, P. and Hudson, B.G., *In vitro* kinetic studies of formation of antigenic advanced glycation end products (AGEs) novel inhibition of post-Amadori glycation pathways, *Journal of Biological Chemistry*, 1997, 272(9), 5430-5437.

Borhanuddin M, Shamsuzzoha M. and Hussain A.H., Hypoglycemia effects of *Andrographis paniculata* Nees on non-diabetic rabbits, *Bangladesh Medical Research Council Bulletin*, 1994,20, 24-26.

Brownlee, M., Biochemistry and molecular cell biology of diabetic complications, *Nature*, 2001, 414(6865), 813-820.

Cameron, N.E., Gibson, T.M., Nangle, M.R. and Cotter, M.A., Inhibitors of advanced glycation endproduct formation and neurovascular dysfunction in experimental diabetes, *Annals of the New York Academy of Sciences*, 2005, 1043(1), 784-792.

Carini, M., Aldini, G., Beretta, G., Arlandini, E. and Facino, R.M., Acrolein-sequestering ability of endogenous dipeptides: characterization of carnosine and homocarnosine/acrolein adducts by electrospray ionization tandem mass spectrometry, *Journal of Mass Spectrometry*, 2003, 38(9), 996-1006.

Casazza, J.P., Felver, M.E. and Veech, R.L., The metabolism of acetone in rat, *Journal of Biological Chemistry*, 1984, 259(1), 231-236.

Cavalcanti, R.N., Navarro-Díaz, H.J., Santos, D.T., Rostagno, M.A. and Meireles, M.A.A., Supercritical carbon dioxide extraction of polyphenols from pomegranate (*Punica granatum* L.) leaves: Chemical composition, economic evaluation and chemometric approach, *Journal of Food Research*, 2012, 1(3), 282-294.

Cervantes-Laurean, D., Schramm, D.D., Jacobson, E.L., Halaweish, I., Bruckner, G.G. and Boissonneault, G.A., Inhibition of advanced glycation end product formation on collagen by rutin and its metabolites, *The Journal of Nutritional Biochemistry*, 2006, 17(8), 531-540.

Chakrabarti R., and Rajagopalan R., Diabetes and insulin resistance associated disorders: Disease and the therapy, *Current Science*, 2002, 83 (12),1533-1538.

Chenwitheesuk, A., Teerawutgulrag, A. and Rakariyatham, N., Screening of antioxidant activity and antioxidant compounds of some edible plants of Thailand, *Food Chemistry*, 2005, 92(3), 491-497.

Charles, D.J., Simon, J.E. and Singh, N.K., The essential oil of *Alpinia galanga* Willd. *Journal of Essential Oil Research*, 1992, 4(1), 81-82.

Chattopadhyay R.R., Chattopadhyay R.N., Nandy A.K., Poddar G. and Maitra S.K., The effect of fresh leaves of *Azadiracta indica* on glucose uptake and glycogen content in the isolated rat hemidiaphragm, *Bulletin of the Calcutta School of Tropical Medicine*, 1987, 35, 8-12.

Cheng, J. and Yang, R., Hypoglycemic effect of guava juice in mice and human subjects, *American Journal of Chinese Medicine*, 1983, 11 (1-4), 74-76.

Cheng, Y.S. and Liu, M. L., Oils of *Ocimum basilicum* L; *Ocimum basilicum* var. *minimum* L.; and *Ocimum gratissimum* L. grown in Taiwan. In Proceeding 9th International Congress of Essential oils, Singapore, 1983, Essential Oil Technical Papers Book 4.

Chompoo, J., Upadhyay, A., Kishimoto, W., Makise, T. and Tawata, S., Advanced glycation end products inhibitors from *Alpinia zerumbet* rhizomes, *Food Chemistry*, 2011, 129(3), 709-715.

Christen, Y., Oxidative stress and Alzheimer's disease, *The American Journal of Clinical Nutrition*, 2000, 71(2), 621s-629s.

Chung, Y.C., Lin, C.C., Chou, C.C. and Hsu, C.P, The effect of longan seed polyphenols on colorectal arcinoma cells, *European Journal of Clinical Investigation*, 2010, 40, 713-721.

Ciolino, H. and Yeh, G., The flavonoid galangin is an inhibitor of CYP1A1 activity and an agonist/antagonist of the aryl hydrocarbon receptor, *British Journal of Cancer*, 1999, 79(9-10), 1340.

Daisy, P. and Rajathi, M., Hypoglycemic effects of *Clitoria ternatea* Linn.(Fabaceae) in alloxan-induced diabetes in rats. *Tropical Journal of Pharmaceutical Research*, 2009, 8(5), 393-398.

de la Maza, M.P., Garrido, F., Escalante, N., Leiva, L., Barrera, G., Schnitzler, S., Zanolli, M., Verdaguer, J., Hirsch, S. and Jara, N., Fluorescent advanced glycation end-products (ages) detected by spectro-photofluorimetry, as a screening tool to detect diabetic microvascular complications, *Journal of Diabetes Mellitus*, 2012, 2, 221.

Devi, B.P., Boominathan, R. and Mandal, S.C., Anti-inflammatory, analgesic and antipyretic properties of *Clitoria ternatea* root, *Fitoterapia*, 2003, 74(4), 345-349.

Devi, P.U., Ganasoundari, A., Rao, B. and Srinivasan, K., In vivo radioprotection by *Ocimum* flavonoids: survival of mice, *Radiation Research*, 1999, 151(1), 74-78.

Ellis, E.N. and Good, B.H., Prevention of glomerular basement membrane thickening by aminoguanidine in experimental diabetes mellitus, *Metabolism*, 1991, 40(10), 1016-1019.

Esmailzadeh, F., Tahbaz, I., Gaieni, H., Alavi-Majd, L. and Azadbakht, Farideh concentrated pomegranate juice improves lipid profiles in diabetic patients with hyperlipidemia, *Journal of Medicinal Food*, 2004, 7(3), 305-308.

Farnworth, N.R., and Bunyapraphatsara, N., Thai medicinal plants, Prachachon Press, Bangkok, Thailand, 1992.

Farsi, D.A., Harris, C.S., Reid, L., Bennett, S.A., Haddad, P.S., Martineau, L.C. and Arnason, J.T., Inhibition of non-enzymatic glycation by silk extracts from a

Mexican land race and modern inbred lines of maize (*Zea mays*), *Phytotherapy Research*, 2008, 22(1), 108-112.

Fecka, I. and Turek, S., Determination of polyphenolic compounds in commercial herbal drugs and spices from Lamiaceae: thyme, wild thyme and sweet marjoram by chromatographic techniques, *Food Chemistry*, 2008, 108(3), 1039-1053.

Freedman, B.I., Wuerth, J.-P., Cartwright, K., Bain, R.P., Dippe, S., Hershon, K., Mooradian, A.D. and Spinowitz, B.S., Design and baseline characteristics for the aminoguanidine clinical trial in overt Type 2 diabetic nephropathy (ACTION II). *Controlled Clinical Trials*, 1999, 20(5), 493-510.

Garret, R.H. and Charles, M.G., *Biochemistry*, Belmont: Thomson Brooks/Cole, 2005, 382-394.

Gella, A. and Durany, N., Oxidative stress in Alzheimer's disease. *Cell Adhesion & Migration*, 2009, 3(1), 88-93.

Gil, M.I., Tomás-Barberán, F.A., Hess-Pierce, B., Holcroft, D.M. and Kader, A.A., Antioxidant activity of pomegranate juice and its relationship with phenolic composition and processing, *Journal of Agricultural and Food Chemistry*, 2000, 48(10), 4581-4589.

Giugliano, D., Ceriello, A., Paolisso, G., Oxidative stress and diabetic vascular complications, *Diabetes Care*, 1996, 19(3), 257-267.

Gkogkolou, P. and Böhm, M., Advanced glycation end products; Key player for skin aging ?, *Dermato-Endocrinology*, 2012, 4(3), 259-270.

Goh, S.H., Chuah, Mok, J.S.L. and Soepadmo, E., *Malaysian medicinal plants for the treatment of cardiovascular diseases*, Academe Art and Printing Services Sdn. Bhd, Kuala Lumpur, 1995.

Goh, S.-Y. and Cooper, M.E., The role of advanced glycation endproducts in progression and complications of diabetes, *Journal of Clinical Endocrinology & Metabolism*, 2008, 93(4), 1143-1152.

Goldin, A., Beckman, J.A., Schmidt, A.M. and Creager, M.A., Advanced glycation endproducts sparking the development of diabetic vascular injury, *Circulation*, 2006, 114(6), 597-605.

Golej, J., Hoeger H Fau - Radner, W., Radner W Fau - Unfried, G., Unfried G Fau - Lubec, G. and Lubec, G., Oral administration of methylglyoxal leads to kidney collagen accumulation in the mouse, *Life Science*, 1998, 63(3), 801-807.

Grayer, R.J., Bryan, S.E., Veitch, N.C., Goldstone, F.J., Paton, A. and Wollenweber, E., External flavones in sweet basil, *Ocimum basilicum* and related taxa, *Phytochemistry*, 1996, 43(5), 1041-1047.

Gugliucci, A. and Bendayan, M., Renal fate of circulating advanced glycated end products (AGEs): evidence for reabsorption and catabolism of AGE-peptides by renal proximal tubular cells, *Diabetologia*, 1996, 39(2), 149-160.

Gugliucci, A., Bastos, D.H.M., Schulze, J. and Souza, M.F.F., Caffeic and chlorogenic acids in *Ilex paraguariensis* extracts are the main inhibitors of AGE generation by methylglyoxal in model proteins, *Fitoterapia*, 2009, 80(6), 339-344

Gugliucci, A., Glycation as the glucose link to diabetic complications, *JAOA: Journal of the American Osteopathic Association*, 2000, 100(10), 621-634.

Gupta, P., Yadav, D.K., Siripurapu, K.B., Palit, G., Maurya, R., Constituents of *Ocimum sanctum* with Antistress Activity, *Journal of Natural Products*, 2007, 70(9), 1410-1416.

Gülçın, İ., Oktay, M., Kireççi, E., Küfrevioğlu, Ö. İ., Screening of antioxidant and antimicrobial activities of anise (*Pimpinella anisum* L.) seed extracts, *Food Chemistry*, 2003, 83, 371-382

Hamelin, M. Borot-Laloi, C., Friguet, B. and Bakala, H., Increased level of glyoxidation product N^ε(epsilon)-(carboxymethyl)lysine in rat serum and urine proteins with aging: link with glyoxidative damage accumulation in kidney, *Archives of Biochemistry and Biophysics*, 2003, 411, 215-222.

Hamid, A.A., Shah, Z.M., Muse, R. and Mohamed, S., Characterisation of antioxidative activities of various extracts of *Centella asiatica* (L) Urban, *Food Chemistry*, 2002, 77(4), 465-469.

Harborne, J.B., *Phytochemical methods a guide to modern techniques of plant analysis*, Chapman & Hall, 1998.

Hartog, J.W., Voors, A.A., Bakker, S.J., Smit, A.J. and Veldhuisen, D.J., Advanced glycation end-products (AGEs) and heart failure: pathophysiology and clinical implications, *European journal of Heart Failure*, 2007, 9(12), 1146-1155.

Hedges, L. and Lister, C., Nutritional attributes of herbs, Crop and Food Research Confidential Report (1891), Christchurch, New Zealand, 2007.

Hiltunen, R. and Holm, Y., *Basil: the genus Ocimum*, CRC Press, 2003.

Hipkiss, A.R., Michaelis, J. and Syrris, P., Non-enzymatic glycosylation of the dipeptide L-carnosine, a potential anti-protein-cross-linking agent, *FEBS letters*, 1995, 371(1), 81-85.

Ho, S.C., Wu, S.P., Lin, S.M. and Tang, Y.L., Comparison of anti-glycation capacities of several herbal infusions with that of green tea, *Food Chemistry*, 2010, 122(3), 768-774.

Hossain, M.B., Rai, D.K., Brunton, N.P., Martin-Diana, A.B. and Barry-Ryan, C., Characterization of Phenolic Composition in Lamiaceae Spices by LC-ESI-MS/MS, *Journal of Agricultural and Food Chemistry*, 2010, 58(19), 10576-10581.

Hsieh, C.-L., Peng, C.-H., Chyau, C.-C., Lin, Y.-C., Wang, H.-E. and Peng, R.Y., Low-density lipoprotein, collagen, and thrombin models reveal that *Rosemarinus officinalis* L. exhibits potent antiglycative effects, *Journal of Agricultural and Food Chemistry*, 2007, 55(8), 2884-2891.

Huang, R.Q., Zou, X.Y. and Liu, X.M., Study on the hypoglycemic effect of longan seed extract, *Natural Product Research and Development*, 2006, 18, 991–992.

Huda-Faujan, N., Noriham, A., Norrakiah, A.S. and Babji, A.S., Antioxidant activity of plants methanolic extracts containing phenolic compounds, *African Journal of Biotechnology*, 2009, 8(3), 484-489.

Hunt, J.V., Bottoms, M.A. and Michinson, M.J., Oxidative alterations in the experimental glycation model of diabetes mellitus are due to protein-glucose adduct oxidation, *Biochemical Journal*, 1993, 291, 529-535.

Hunter, M.V., Brophy, J.J., Ralph, B.J. and Bienvenu, F.E., Composition of *Polygonum odoratum* Lour. from southern Australia, *Journal of Essential Oil Research*, 1997, 9(5), 603-604.

Hussain, J., Ullah, R., Khan, F.U., Muhammad, Z., Khan, N., ur Rehman, N. and Hussain, H., Antiglycation and Antimicrobial Activities of the Crude Extract of *Phlomis bracteosa*, *American-Eurasian Journal of Agricultural & Environmental Sciences*, 2010, 7(6), 634-636.

Jacobson, E., Jacobson, M. and Wondrak, G., Method for identifying regulators of protein-age formation, 2001,

Jafri, M.A., Aslam, M., Javed, K. and Singh, S., Effect of *Punica granatum* Linn. (flowers) on blood glucose level in normal and alloxan-induced diabetic rats, *Journal of Ethnopharmacology*, 2000, 70 (3) , 309–314.

Jaggi, R.K., Madaan, R., Singh, B., Anticonvulsant potential of holy basil, *Ocimum sanctum* Linn., and its cultures, *Indian Journal of Experimental Biology*, 2003, 41, 1329-1333.

Jain, R.C. and Vyas, S.R., Garlic in alloxan-induced diabetic rabbits, *American Journal of Clinical Nutrition*, 1975, 28, 684-685.

Jalal, R., Bagheri, S.M., Moghimi, A. and Rasuli, M.B., Hypoglycemic effect of aqueous shallot and garlic extracts in rats with fructose-induced insulin resistance, *Journal of Clinical Biochemistry and Nutrition*, 2007, 41(3), 218-223.

Kao, T., Huang, S., Inbaraj, B.S. and Chen, B., Determination of flavonoids and saponins in *Gynostemma pentaphyllum* (Thunb.) Makino by liquid chromatography-mass spectrometry, *Analytica Chimica Acta*, 2008, 626(2), 200-211.

Kartnig, T., Clinical application of *Centella asiatica* (L.) Urb., *Herbs, Spices and Medicinal Plants*, 1988, 3, 145-173.

Khalifah, R.G., Todd, P., Booth, A.A., Yang, S.X., Mott, J.D. and Hudson, B.G., Kinetics of nonenzymatic glycation of ribonuclease a leading to advanced glycation end products. Paradoxical inhibition by ribose leads to facile isolation of protein intermediate for rapid post-Amadori studies, *Biochemistry*, 1996, 35(15), 4645-4654.

Kim, H.Y. and Kim, K., Protein glycation inhibitory and antioxidative activities of some plant extracts *in vitro*, *Journal of Agricultural and Food Chemistry*, 2003, 51(6), 1586-1591.

Kim, S.-S., Gallaher, D. and Csallany, A.S., Vitamin E and probucol reduce urinary lipophilic aldehydes and renal enlargement in streptozotocin-induced diabetic rats, *Lipids*, 2000, 35(11), 1225-1237.

Kislinger, T., Fu, C., Huber, B., Qu, W., Taguchi, A., Du Yan, S., Hofmann, M., Yan, S.F., Pischetsrieder, M. and Stern, D., N^ε-(carboxymethyl) lysine adducts of proteins are ligands for receptor for advanced glycation end products that activate cell signaling pathways and modulate gene expression, *Journal of Biological Chemistry*, 1999, 274(44), 31740-31749.

Knerr, T. and Severin, T., Reaction of glucose with guanosine, *Tetrahedron Letters*, 1993, 34(46), 7389-7390.

Kumar, M., Reddy, P., Kumar, P., Surolia, I. and Reddy, G., Effect of dicarbonyl-induced browning on alpha-crystallin chaperone-like activity: physiological significance and caveats of *in vitro* aggregation assays, *Biochemical Journal*, 2004, 379, 273-282.

Kumari, K. and Augusti, K.T., Antidiabetic and antioxidant effects of *S*-methyl cysteine sulfoxide isolated from onions (*Allium cepa* Linn.) as compared to standard drugs in alloxan diabetic rats, *Indian Journal of Experimental Biology*, 2002, 40(9), 1005-1009.

Lee, S., Park, H.-S., Notsu, Y., Ban, H.S., Kim, Y.P., Ishihara, K., Hirasawa, N., Jung, S.H., Lee, Y.S., Lim, S.S., Park, E.-H., Shin, K.H., Seyama, T., Hong, J. and Ohuchi, K., Effects of hyperin, isoquercitrin and quercetin on lipopolysaccharide-induced nitrite production in rat peritoneal macrophages, *Phytotherapy Research*, 2008, 22(11), 1552-1556.

Lin, L., Dong, Y., Zhao, H., Wen, L., Yang, B. and Zhao, M., Comparative evaluation of rosmarinic acid, methyl rosmarinate and pedalitin isolated from *Rabdosia serra* (MAXIM.) HARA as inhibitors of tyrosinase and α -glucosidase, *Food Chemistry*, 2011, 129(3), 884-889.

Litchfield, J.E., Thorpe, S.R. and Baynes, J.W., Oxygen is not required for the browning and crosslinking of protein by pentoses: relevance to Maillard reactions

in vivo, *The International Journal of Biochemistry & Cell Biology*, 1999, 31(11), 1297-1305.

Lo, C.Y., Li, S., Tan, D., Pan, M.H., Sang, S. and Ho, C.T., Trapping reactions of reactive carbonyl species with tea polyphenols in simulated physiological conditions, *Molecular Nutrition & Food Research*, 2006, 50(12), 1118-1128.

Lunceford, N. and Gugliucci, A., *Ilex paraguariensis* extracts inhibit AGE formation more efficiently than green tea, *Fitoterapia*, 2005, 76(5), 419-427.

Luthra, M. and Balasubramanian, D., Nonenzymatic glycation alters protein structure and stability, A study of two eye lens crystallins, *Journal of Biological Chemistry*, 1993, 268(24), 18119-18127.

Ly, L., Gu, X., Tang, J. and Ho, C.-T., Antioxidant activity of stilbene glycoside from *Polygonum multiflorum* Thunb. *vivo*, *Food Chemistry*, 2007, 104(4), 1678-1681.

Lv, L., Shao, X., Wang, L., Huang, D., Ho, C.-T. and Sang, S., Stilbene glucoside from *Polygonum multiflorum* Thunb.: a novel natural inhibitor of advanced glycation endproduct formation by trapping of methylglyoxal. *Journal of Agricultural and Food Chemistry*, 2010, 58(4), 2239-2245.

Lyles, G.A. and Chalmers, J., The metabolism of aminoacetone to methylglyoxal by semicarbazide-sensitive amine oxidase in human umbilical artery, *Biochemical Pharmacology*, 1992, 43(7), 1409-1414.

Ma, H.-Y., Gao, H.-Y., Sun, L., Huang, J., Xu, X.-M. and Wu, L.-J., Constituents with α -glucosidase and advanced glycation end-product formation inhibitory activities from *Salvia miltiorrhiza* Bge, *Journal of Natural Medicines*, 2011, 65(1), 37-42.

Mammadov, R., Ili, P. and Ertem Vaizogullar, H. Antioxidant activity and total phenolic content of *Gagea fibrosa* and *Romulea ramiflora*, *Iranian Journal of Chemical Engineering*, 2011, 30, 57-62.

Manaharan, T., Teng, L.L., Appleton, D., Ming, C.H., Masilamani, T., Palanisamy, U.D., Antioxidant and antiglycemic potential of *Peltophorum pterocarpum* plant parts, *Food Chemistry*, 2011, 129(4), 1355-1362.

Manohan, R., Papanuvej, C. and Ruangrunsi, N., Genotoxic and cytotoxic effects of Ben-Cha-Moon-Yai remedy and its ingredients, *Research Journal of Pharmaceutical, Biological and Chemical Science*, 2013, 4(1), 1144-1155.

Martinez, G., Delgado, R., Perez, G., Garrido, G., Nunez Selles, A.J. and Leon, O.S., Evaluation of the in vitro antioxidant activity of *Mangifera indica* L. extract, *Phytotherapy Research*, 2000, 14(6), 424-427.

Matsuda, H., Wang, T., Managi, H. and Yoshikawa, M., Structural requirements of flavonoids for inhibition of protein glycation and radical scavenging activities, *Bioorganic & Medicinal Chemistry*, 2003, 11(24), 5317-5323.

Matsuura, N., Aradate, T., Sasaki, C., Kojima, H., Ohara, M., Hasegawa, J. and Ubukata, M., Screening System for the Maillard Reaction Inhibitor from Natural Product Extracts, *Journal of Health Science*, 2002, 48(6), 520-526.

McKay, D.L. and Blumberg, J.B., The role of tea in human health: an update, *Journal of the American College of Nutrition*, 2002, 21(1), 1-13.

Megalli, S., Aktan, F., Davies, N.M., Roufogalis, B.D., Phytopreventive anti-hyperlipidemic effects of *Gynostemma pentaphyllum* in rats. *Journal of Pharmacy and Pharmaceutical Sciences*, 2005, 8(3), 507-515.

Metz, T.O., Alderson, N.L., Chachich, M.E., Thorpe, S.R. and Baynes, J.W., Pyridoxamine traps intermediates in lipid peroxidation reactions in vivo evidence on the role of lipids in chemical modification of protein and development of

diabetic complications, *Journal of Biological Chemistry*, 2003, 278(43), 42012-42019.

Miroliaei, M., Khazaei, S., Moshkelgosha, S. and Shirvani, M., Inhibitory effects of Lemon balm (*Melissa officinalis*, L.) extract on the formation of advanced glycation end products, *Food Chemistry*, 2011, 129(2), 267-271.

Mittelmaier, S., Fünfroeken, M., Fenn, D., Berlich, R., Pischetsrieder, M., Quantification of the six major α -dicarbonyl contaminants in peritoneal dialysis fluids by UHPLC/DAD/MSMS, *Analytical and Bioanalytical Chemistry*, 2011, 401(4), 1183-1193.

Miyata, T., Ueda, Y., Horie, K., Nangaku, M., Tanaka, S., de Strihou, C.v.Y. and Kurokawa, K., Renal catabolism of advanced glycation end products: the fate of pentosidine, *Kidney International*, 1998, 53(2), 416-422.

Modak, M., Dixit, P., Londhe, J., Ghaskadbi, S. and Devasagayam, T.P.A., Indian herbs and herbal drugs used for the treatment of diabetes, *Journal of Clinical Biochemistry and Nutrition*, 2007, 40(3), 163-173.

Monnier, V.M., Intervention against the Maillard reaction *in vivo*, *Archives of Biochemistry and Biophysics*, 2003, 419, 1-15.

Muruganandan, S., Srinivasan, K, Gupta, S., Gupta, D.K. and Lal, J., Effect of mangiferin on hyperglycemia and atherogenecity in streptozotocin diabetic rats. *Journal of Ethnopharmacology*, 2005, 97, 497–501.

Muthenna, P., Akileshwari, C., Saraswat, M. and Bhanuprakash Reddy, G., Inhibition of advanced glycation end-product formation on eye lens protein by rutin, *British Journal of Nutrition*, 2011, 25: 1-9.

Nanasombat, S. and Teckchuen, N., Antimicrobial, antioxidant and anticancer activities of Thai local vegetables, *Journal of Medicinal Plants Research*, 2009, 3(5), 443-449.

Nawale, R., Mourya, V. and Bhise, S., Non-enzymatic glycation of proteins: a cause for complications in diabetes, *Indian Journal of Biochemistry and Biophysics*, 2006, 43(6), 337.

Noor, H. and Ashcroft, S. J. H., Pharmacological characterisation of the antihyperglycaemic properties of *Tinospora crispa* extract, *Journal of Ethnopharmacology*, 1998, 62, 7-13.

Oh, W.K., Lee, C.H., Lee, M.S., Bae, E.Y., Sohn, C.B., Oh, H., Kim, B.Y. and Ahn, J.S., Antidiabetic effects of extracts from *Psidium guajava*, *Journal of Ethnopharmacology*, 2005, 96 (3), 411-415..

Ojewole, J.A.O., Antiinflammatory, analgesic and hypoglycemic effects of *Mangifera indica* Linn. (Anacardiaceae) stem-bark aqueous extract, *Methods and Findings in Experimental and Clinical Pharmacology*, 2005, 27 (8), 547-554.

Oki, T., Matsui, T. and Osajima, Y., Inhibitory effect of α -glucosidase inhibitors varies according to its origin, *Journal of Agricultural and Food Chemistry*, 1999, 47(2), 550-553.

Okuda, Y., Sakoda, S., Fujimura, H. and Yanagihara, T., Aminoguanidine, a selective inhibitor of the inducible nitric oxide synthase, has different effects on experimental allergic encephalomyelitis in the induction and progression phase, *Journal of Neuroimmunology*, 1998, 81(1-2), 201-210.

Ou, P. and Wolff, S.P., Aminoguanidine: A drug proposed for prophylaxis in diabetes inhibits catalase and generates hydrogen peroxide in vitro. *Biochemical Pharmacology*, 1993, 46(7), 1139-1144.

Pari, L. and Maheswari, J.U., Antihyperglycemic activity of *Musa sapientum* flowers: effect on lipid peroxidation in alloxan diabetic rats, *Phytotherapy Research*, 2000, 14(2), 136-138.

Pashikanti, S., Boissonneault, G.A. and Cervantes-Laurean, D., (2011) Ex vivo detection of histone H1 modified with advanced glycation end products, *Free Radical Biology and Medicine*, 2011, 50(10), 1410-1416.

Pashikanti, S., de Alba, D.R., Boissonneault, G.A. and Cervantes-Laurean, D., Rutin metabolites: novel inhibitors of nonoxidative advanced glycation end products, *Free Radical Biology and Medicine*, 2010, 48(5), 656-663.

Pashikanti, S., de Alba, D.R., Boissonneault, G.A. and Cervantes-Laurean, D., Rutin metabolites: novel inhibitors of nonoxidative advanced glycation end products, *Free Radical Biology and Medicine*, 2010, 48(5), 656-663.

Pattanayak, P., Behera, P., Das, D., Panda, S.K., *Ocimum sanctum* Linn. A reservoir plant for therapeutic applications: An overview, *Pharmacognosy Reviews*, 2010, 4(7), 95-105.

Peng, X., Cheng, K.W., Ma, J., Chen, B., Ho, C.T., Lo, C., Chen, F. and Wang, M., Cinnamon bark proanthocyanidins as reactive carbonyl scavengers to prevent the formation of advanced glycation endproducts, *Journal of Agricultural and Food Chemistry*, 2008a, 56(6), 1907-1911.

Peng, X., Ma, J., Chao, J., Sun, Z., Chang, R.C.C., Tse, I., Li, E.T.S., Chen, F. and Wang, M., Beneficial effects of cinnamon proanthocyanidins on the formation of specific advanced glycation endproducts and methylglyoxal-induced impairment on glucose consumption, *Journal of Agricultural and Food Chemistry*, 2010, 58(11), 6692-6696.

Peng, X., Zheng, Z., Cheng, K.W., Shan, F., Ren, G.X., Chen, F. and Wang, M. Inhibitory effect of mung bean extract and its constituents vitexin and isovitexin on the formation of advanced glycation endproducts, *Food Chemistry*, 2008b, 106(2), 475-481.

Peungvicha, P., Thirawarapan, S. S., Temsiririrkkul, R., Watanabe, H., Prasain, J. K. and Kadota S., Hypoglycemic effect of the water extract of *Piper sarmentosum* in rats, *Journal of Ethnopharmacology*, 1998, 60, 27-32.

Porkorný, J., Yanishlieva N., and Gordon, M., *Antioxidants in food: practical applications*, Elsevier, 2001.

Pongmarutai, M., Studying antidiabetic action of *Piper rostratum*, M.Sc. Thesis, Mahidol University, Bangkok, Thailand, 1980.

Pripdeevech, P., Chumpolsri, W., Suttiarporn, P. and Wongpornchai, S., The chemical composition and antioxidant activities of basil from Thailand using retention indices and comprehensive two-dimensional gas chromatography, *Journal of the Serbian Chemical Society*, 2010, 75(11), 1503-1513.

Rabbani, N. and Thornalley, P.J., Dicarbonyls linked to damage in the powerhouse: glycation of mitochondrial proteins and oxidative stress, *Biochemistry Society Transactions*, 2008, 36, 1045-1050.

Rahman, Md. S. Islam, Md. R., Kamruzzaman, Md., Alam Md. K. and Hena Md. A. , *Ocimum sanctum* L.: A review of its phytochemical and pharmacological profile, *Research Journal of Medicinal Plant*, 2011, 1-15.

Rai, V., Iyer U. and Mani U.V. Effect of Tulasi (*Ocimum sanctum*) leaf powder supplementation on blood sugar levels, serum lipids and tissue lipid in diabetic rats, *Plant Food For Human Nutrition*, 1997, 50, 9–16.

Rajathi, M. and Daisy, P., Effect of plant extracts on the blood glucose and cholesterol level of alloxan diabetic rabbit, *Indian Journal of Comparative Animal Physiology*, 2000, 18, 14-17.

Rangkadilok, N., Sitthimonchai, S., Worsuttayangkurn, L., Mahidol, C., Ruchirawat, M. and Satayavivad, J., Evaluation of free radical scavenging and

antityrosinase activities of standardized longan fruit extract, *Food and Chemical Toxicology*, 2007, 45, 328–336.

Rasheed, Z., Anbazhagan, A.N., Akhtar, N., Ramamurthy, S., Voss, F.R., Haqqi, T.M., Green tea polyphenol epigallocatechin-3-gallate inhibits advanced glycation end product-induced expression of tumor necrosis factor- α and matrix metalloproteinase 13 in human chondrocytes, *Arthritis Research & Therapy*, 2009, 11(3),R71.

Reddy, V.P. and Beyaz, A., Inhibitors of the Maillard reaction and AGE breakers as therapeutics for multiple diseases, *Drug Discovery Today*, 2006, 11(13), 646-654.

Rolo, A.P. and Palmeira, C.M., Diabetes and mitochondrial function: role of hyperglycemia and oxidative stress, *Toxicology and Applied Pharmacology*, 2006, 212(2), 167-178.

Roman-Ramos, R., Flores-Saenz, J.L. and Alarcon-Aguilar, F.J., Anti-hyperglycemic effect of some edible plants, *Journal of Ethnopharmacology*, 1995, 48, 25-32.

Rukachaisirikul, T., Siriwatanakit, P., Sukcharoenphol, K., Wongvein, C., Ruttanaweang, P., Wongwattanavuch, P. and Suksamrarn, A., Chemical constituents and bioactivity of *Piper sarmentosum*, *Journal of Ethnopharmacology*, 2004, 93(2), 173-176.

Saito, A., Takeda, T., Sato, K., Hama, H., Tanuma, A., Kaseda, R., Suzuki, Y. and Gejyo, F., Significance of proximal tubular metabolism of advanced glycation end products in kidney diseases. *Annals of the New York Academy of Sciences*, 2005, 1043(1), 637-643.

Sang, S., Shao, X., Bai, N., Lo, C.-Y., Yang, C.S. and Ho, C.-T., Tea polyphenol (–)-epigallocatechin-3-gallate: a new trapping agent of reactive dicarbonyl species, *Chemical Research in Toxicology*, 2007, 20(12), 1862-1870.

Saraswat, M., Suryanarayana, P., Reddy, P.Y., Patil, M.A., Balakrishna, N. and Reddy, G.B., Antiglycating potential of *Zingiber officinalis* and delay of diabetic cataract in rats, 2010, 16, 1525-1537.

Sawa, T., Nakao, M., Akaike, T., Ono, K. and Maeda, H., Alkylperoxyl radical-scavenging activity of various flavonoids and other phenolic compounds: implications for the anti-tumor-promoter effect of vegetables, *Journal of Agricultural and Food Chemistry*, 1999, 47(2), 397-402.

Segal, I., Enzyme Kinetics. John Wiley & Sons, Inc., 1993.

Seidel, W. and Pischetsrieder, M., Reaction of guanosine with glucose under oxidative conditions, *Bioorganic and Medicinal Chemistry Letters*, 1998, 8(15), 2017-2022.

Shavandi, M.A., Haddadian, Z. and Ismail, M.H.S., *Eryngium Foetidum* L. *Coriandrum Sativum* and *Persicaria Odorata* L.: A Review, *Journal of Asian Scientific Research*, 2012, 2(8), 410-426.

Shimizu, K., Ozeki, M., Tanaka, K., Itoh, K., Nakajyo, S., Urakawa, N., and Atsuchi, M., Suppression of glucose absorption by extracts from the leaves of *Gymnema inodorum*, *The Journal of veterinary medical science/the Japanese Society of Veterinary Science*, 1997, 59(9), 753-757.

Silván, J.M., Assar, S.H., Srey, C., Dolores del Castillo, M. and Ames, J.M., Control of the Maillard reaction by ferulic acid, *Food Chemistry*, 2011, 128(1), 208-213.

Sivaci, A. and Duman, S., Evaluation of seasonal antioxidant activity and total phenolic compounds in stem and leaves of some almond (*Prunus amygdalus* L.) varieties, *Biological Research*, 2014, 47(9), 1-5.

Skeggs, L.T., Khan, J.R. and Shumway, N.P., The preparation and function of the hypertension – converting enzyme. *Journal of Experimental Medicine*, 1956, 103, 295-299.

Soman, S., Rauf, A.A., Indira, M. and Rajamanickam, C., Antioxidant and antiglycative potential of ethyl acetate fraction of *Psidium guajava* leaf extract in streptozotocin-induced diabetic rats, *Plant Foods for Human Nutrition*, 2010, 65(4), 386-391.

Sookkongwaree, K., Geitmann, M., Roengsumran, S., Petsom, A. and Danielson, U.H., Inhibition of viral proteases by Zingiberaceae extracts and flavones isolated from *Kaempferia parviflora*, *Pharmazie*, 2006, 61(8), 717-721.

Soong, Y.Y. and Barlow, P.J., Isolation and structure elucidation of phenolic compounds from longan (*Dimocarpus longan* Lour.) seed by high-performance liquid chromatography-electrospray ionization mass spectrometry, *Journal of Chromatography A*, 2005, 1085, 270–277.

Soulis, T., Cooper, M.E., Vranes, D., Bucala, R. and Jerums, G., Effects of aminoguanidine in preventing experimental diabetic nephropathy are related to the duration of treatment, *Kidney International*, 1996, 50(2), 627-634.

Srivastava, A.K., Srivastava, P., Mishra, J., Singh, M.P., Sharma, A.K. and Shrivastava, A., Antidiabetic activity of the stem extracts of *Cissus quadrangularis* linn. *Journal of Pharmacy Research*, 2011,4(11), 3873-3874.

Sudjaroen, Y., Hull, W.E., Erben, G., Würtele, G., Changbumrung, S., Ulrich, C.M. and Owen, R.W., Isolation and characterization of ellagitannins as the major polyphenolic components of Longan (*Dimocarpus longan* Lour.) seeds, *Phytochemistry*, 2012, 77, 226–237.

Sun, Z., Peng, X., Liu, J., Fan, K.-W., Wang, M. and Chen, F., Inhibitory effects of microalgal extracts on the formation of advanced glycation endproducts (AGEs), *Food Chemistry*, 2010, 120(1), 261-267.

Syamsudin, Summarny, R. and Simanjuntak, P., Antidiabetic activity of active fractions of *Leucaena leucocephala* (lmk) Dewit seeds in experiment model, *European Journal of Scientific Research*, 2010, 43(3), 384-391.

Tadera, K., Minami, Y., Takamatsu and K., Matsuoka, T., Inhibition of α -glucosidase and α -amylase by flavonoids. *Journal of Nutritional Science and Vitaminology*, 2006, 52(2), 149-153.

Tang, S.Y., Whiteman, M., Peng, Z.F., Jenner, A., Yong, E.L. and Halliwell, B., Characterization of antioxidant and antiglycation properties and isolation of active ingredients from traditional Chinese medicines, *Free Radical Biology and Medicine*, 2004, 36(12), 1575-1587.

Thitilertdecha, N., Teerawutgulrag, A. and Rakariyatham, N., Antioxidant and antibacterial activities of *Nephelium lappaceum* L. extracts, *LWT-Food Science and Technology*, 2008, 41(10), 2029-2035.

Thornalley, P.J., Use of aminoguanidine (Pimagedine) to prevent the formation of advanced glycation endproducts, *Archives of Biochemistry and Biophysics*, 2003, 419(1), 31-40.

Tsuji-Naito, K., Saeki, H. and Hamano, M., Inhibitory effects of Chrysanthemum species extracts on formation of advanced glycation endproducts, *Food Chemistry*, 2009, 116(4), 854-859.

Uribarri, J., Cai, W., Peppas, M., Goodman, S., Ferrucci, L., Striker, G. and Vlassara, H., Circulating glycotoxins and dietary advanced glycation endproducts: two links to inflammatory response, oxidative stress, and aging, *The Journals of Gerontology Series A: Biological Sciences and Medical Sciences*, 2007, 62(4), 427-433.

Usha, V., Vijayammal, P.L. and Kurup, P.A., Effect of dietary fiber from banana (*Musa paradisiaca*) on metabolism of carbohydrates in rats fed cholesterol free diet, *Indian Journal of Experimental Biology*, 1989, 27 (5) , 445-449.

Vashishth, D., Advanced glycation end-products and bone fractures, *IBMS BoneKEy*, 2009, 6(8), 268-278.

Vats V., Grover J.K. and Rathi S.S. Evaluation of antihyperglycemic and hypoglycemic effect of *Trigonella foenum-graecum* Linn, *Ocimum sanctum* Linn and *Pterocarpus marsupium* Linn in normal and alloxanized diabetic rats, *Journal of Ethnopharmacology*. 2002, 79, 95–100.

Vats, V. and Yadav S.P. Grover, Ethanolic extract of *Ocimum sanctum* leaves partially attenuates streptozotocin induced alteration in glycogen content and carbohydrate metabolism in rats, *Journal of Ethnopharmacology*, 2004, 90, 155–160.

Vijayakumari, P., Shanthi, K., Kayalvizhi, J., Muruganatham, G. Sethuraman, M. and Thirumurugan, V., Studies on the physio-phytochemical and anti diabetic properties of *Cissus quadrangularis* L. and *Solanum torvum* Swartz., *International Journal of Drug Discovery and Herbal Research*, 2012, 2(1), 323-328.

Vimala, S., Adenan, M.L., Ahmad, A.R. and Shahdan, R., Nature's choice to wellness: antioxidant vegetables/ulam. Malaysia, Forest Research Institute, 2003.

Vinson, J.A. and Howard, T.B., Inhibition of protein glycation and advanced glycation end products by ascorbic acid and other vitamins and nutrients, *The Journal of Nutritional Biochemistry*, 1996, 7(12), 659-663.

Voziyan, P.A. and Hudson, B. G., Pyridoxamine as a multifunctional pharmaceutical: targeting pathogenic glycation and oxidative damage, *Cellular and Molecular Life Sciences*, 2005a, 62(15), 1671-1681.

Voziyan, P.A. and Hudson, B.G., Pyridoxamine: the many virtues of a Maillard reaction inhibitor, *Annals of the New York Academy of Sciences*, 2005b, 1043(1), 807-816.

Wang, R., Ding, Y., Liu, R., Xiang, L. and Du, L., Pomegranate: constituents, bioactivities and pharmacokinetics, *Fruit, Vegetable and Cereal Science and Biotechnology*, 2010, 77-87.

Wang, W., Yagiz, Y., Buran, T.J., Nunes, C.N. and Gu, L., Phytochemicals from berries and grapes inhibited the formation of advanced glycation end-products by scavenging reactive carbonyls, *Food Research International*, 2011, 44(9), 2666-2673.

Waterman, P.G. and Mole, S., Analysis of phenolic plant metabolites, Oxford, Blackwell Scientific Publications, 1994, 84.

Wautier, M., Massin, P., Guillausseau, P., Huijberts, M., Levy, B., Boulanger, E., Laloi-Michelin, M. and Wautier, J., N^ε-(carboxymethyl) lysine as a biomarker for microvascular complications in type 2 diabetic patients, *Diabetes & Metabolism*, 2003, 29(1), 44-52.

Wei, Y., Han, C.S., Zhou, J., Liu, Y., Chen, L. and He, R.Q., D-ribose in glycation and protein aggregation, *Biochimica et Biophysica Acta (BBA)-General Subjects*, 2012, 1820(4), 488-494.

Wendt, T.M., Tanji, N., Guo, J., Kislinger, T.R., Qu, W., Lu, Y., Bucciarelli, L.G., Rong, L.L., Moser, B. and Markowitz, G.S., RAGE drives the development of glomerulosclerosis and implicates podocyte activation in the pathogenesis of diabetic nephropathy, *The American Journal of Pathology*, 2003, 162(4), 1123-1137.

Whittier, F., Spinowitz, B. and Wuerth, J., Pimagedine safety profile in patients with type 1 diabetes, *Journal of American Society of Nephrology*, 1999, 10(9), 184A.

Williams, M.E., Clinical studies of advanced glycation endproduct inhibitors and diabetic kidney disease, *Current Diabetes Reports*, 2004, 4(6), 441-446.

Wróbel, K., Wróbel, K., Garay-Sevilla, M.E., Nava, L.E. and Malacara, J.M., Novel analytical approach to monitoring advanced glycosylation end products in human serum with on-line spectrophotometric and spectrofluorometric detection in a flow system. *Clinical Chemistry*, 1997, 43(9), 1563-1569.

Wu, C.H. and Yen, G.C., Inhibitory effect of naturally occurring flavonoids on the formation of advanced glycation endproducts, *Journal of Agricultural and Food Chemistry*, 2005, 53(8), 3167-3173.

Wu, C.-H., Huang, S.-M., Lin, J.-A. and Yen, G.-C., Inhibition of advanced glycation endproduct formation by foodstuffs, *Food & Function*, 2011, 2(5), 224-234.

Wu, C.-H., Lin, J.-A., Hsieh, W.-C. and Yen, G.-C., Low-density-lipoprotein (LDL)-bound flavonoids increase the resistance of LDL to oxidation and glycation under pathophysiological concentrations of glucose in vitro, *Journal of Agricultural and Food Chemistry*, 2009a, 57(11): 5058-5064.

Wu, J.W., Hsieh, C.L., Wang, H.Y. and Chen, H.Y., Inhibitory effects of guava (*Psidium guajava* L.) leaf extracts and its active compounds on the glycation process of protein. *Food Chemistry*, 2009b, 113(1), 78-84.

Xu, T.M., Xin, Y., Cui, M.H., Jiang, X. and Gu, L.P., Inhibitory effect of ginsenoside Rg3 combined with cyclophosphamide on growth and angiogenesis ovarian cancer. *Chinese Medical Journal*, 2007, 120(7), 584-588.

Yang, Z.G., Sun, H.X. and Ye, Y.P., Ginsenoside Rd from *Panax notoginseng* is cytotoxic towards hela cancer cells and induces apoptosis, *Chemistry & Biodiversity*, 2006, 3(2), 187-197.


Yoo, N.H., Jang, D.S., Yoo, J.L., Lee, Y.M., Kim, Y.S., Cho, J.-H. and Kim, J.S., Erigeroflavanone, a flavanone derivative from the flowers of *Erigeron annuus* with protein glycation and aldose reductase inhibitory activity, *Journal of Natural Products*, 2008, 71(4), 713-715.

You, Q., Chen, F., Wang, X., Jiang, Y., and Lin, S., Anti-diabetic activities of phenolic compounds in muscadine against alpha-glucosidase and pancreatic lipase, *LWT-Food Science and Technology*, 2012, 46(1), 164-168.

Zhang, X.F. and Tan, B.K., Antihyperglycemic and anti-oxidant properties of *Andrographis paniculata* in normal and diabetic rats, *Clinical and Experimental Pharmacology and Physiology*, 2000, 27, 358-363.

Zheng, G., Wei, X., Xu, L., Li, Z., Liu, G. and Zhang, X., A New Natural Lactone from *Dimocarpus longan* Lour. Seeds, *Molecules*, 2012, 17(8), 9421-9425.

Zhishen, J., Mengcheng, T. and Jianming, W., The determination of flavonoid contents in mulberry and their scavenging effects on superoxide radicals, *Food Chemistry*, 1999, 64(4), 555-559.



ลิขสิทธิ์มหาวิทยาลัยเชียงใหม่
Copyright© by Chiang Mai University
All rights reserved