

REFERENCES

- Abyaneh MR, Ghahfarokhi MS, Rezae MB, Jaimand K, Alinezhad S, Saberi R, Yoshinari T. Chemical composition and antiaflatoxigenic activity of *Carum carvi* L., *Thymus vulgaris* and *Citrus aurantifolia* essential oils. *J. Food control*. 2009; 20: 1018-1024.
- Abyaneh MR, Ghahfarokhi MS, Kawachic M, Eslamifar A, Schmidt OJ, Schmidt A, Allameh A, Yoshinari T. Ultrastructural evidences of growth inhibitory effects of a novel biocide, Akacid, on an aflatoxigenic *Aspergillus parasiticus*. *Toxicon*. 2006; 48: 1075-1082.
- Adams RP. Identification of Essential oil components by gas chromatography/quadrupole mass spectroscopy. Allured, Carol Stream, IL, USA. 2001.
- Aggarwal KK, Khanuja SPS, Ahmad A, Kumar TRS, Gupta VK, Kumar S. Antimicrobial activity profiles of the two enantiomers of limonene and carvone isolated from the oils of *Mentha spicata* and *Anethum sowa*. *Flavour. Fragr. J.* 2002; 17: 59-63.
- Albuquerque CC, Camara TR, Marian RDR, Willadino L, Marcelino C, Ulisses C. Antimicrobial action of the essential oil of *Lippia gracilis* Schauer. *Braz. Arch. Biol. Technol.* 2006; 49: 527-535.
- Aligiannis N, Kalpoutzakis E, Mitaku S, Chinou I.B. Composition and Antimicrobial Activity of the Essential Oils of Two *Origanum* Species. *J. Agric. Food Chem.* 2001; 49: 4168-4170.
- Almeida RRP, Souto RNP, Bastos CN, Silvado MHL, Maia JGS. Chemical variation in *Piper aduncum* and biological properties of its dillapiolene-rich essential oil. *Chem Biodiv.* 2009; 6: 1427-1434.
- Amaike S, Keller NP. *Aspergillus flavus*. *Annu. Rev. Phytopathol.* 2011. 49: 107-133.

- Bajpai VK, Kang SC. *In Vitro* and *In Vivo* inhibition of plant pathogenic fungi by essential oil and extracts of *Magnolia liliflora* Desr. *J. Agr. Sci. Tech.* 2012; 14: 845-856.
- Bakkali F, Averbeck S, Averbeck D, Idaomar M. Biological effects of essential oils. *Rev. Food Chem Toxicol.* 2008; 46: 446-475.
- Bassolé IHN, Juliani HR. Essential oils in combination and their antimicrobial properties. *Molecules.* 2012; 17: 3989-4006.
- Bhat R, Rai RV, Karim AA. Mycotoxins in Food and Feed: Present Status and Future Concerns. *J. Comp. Rev. Food Sci. Food Saf.* 2010; 9: 57-81.
- Bluma R, Amaiden MR, Etcheverry M. Screening of argentine plant extracts: impact on growth parameters and aflatoxin B1 accumulation by *Aspergillus* section *Flavi*. *Int. J. Food Microbiol.* 2008; 122: 114-125.
- Boonlue S, Surapat W, Pukahuta C, Suwanarit P, Suwanarit A, Morinaga T. Diversity and efficiency of arbuscular mycorrhizal fungi in soils from organic chili (*Capsicum frutescens*) farms. *J. Mycosci.* 2012; 53: 10-16.
- Burt S. Essential oils: their antibacterial properties and potential applications in foods-a review. *Int. J. Food Microbiol.* 2004; 94: 223-253.
- Cakir A, Kordali S, Zengin H, Izumi S, Hirata T. Composition and antifungal activity of essential oils isolated from *Hypericum hyssopifolium* and *Hypericum heterophyllum*. *Flavour Fragr. J.* 2004; 19: 62-68.
- Callan NW, Johnson DL, Westcott MP, Welty LE. Herb and oil composition of dill (*Anethum graveolens* L.): Effects of crop maturity and plant density. *Ind. Crops Prod.* 2007; 25: 282-287.
- Cardile V, Russo A, Formisano C, Rigano D, Senatore F, Arnold NA, Piozzie F. Essential oils of *Salvia bracteata* and *Salvia rubifolia* from Lebanon: chemical composition, antimicrobial activity and inhibitory effect on human melanoma cells. *J. Ethnopharmacol.* 2009; 126: 265-272.

- Carmo ES, Lima EDO, Souza ELD, Sousa FBD. Effect of *Cinnamomum zeylanicum* essential oil on the growth morphogenesis of some potentially *Aspergillus* species. *Braz. J. Microbiol.* 2008; 39: 91-97.
- Chao LK, Hua KF, Hsu HY, Cheng SS, Lin JY, Chang ST. Study on the anti-inflammatory activity of essential oil from leaves of *Cinnamomum osmophloeum*. *J. Agri. Food Chem.* 2005; 53: 7274-7278.
- Charoenying P, Laosinwattana C, Phuwiwat W, Lomratsiri J. Biological activities of *Zanthoxylum limonella* alston fruit extracts. *KMITL Sci. J.* 2008; 8: 12-15.
- Chemat S, Esveld EDC. Contribution of microwaves or ultrasonics on carvone and limonene recovery from dill fruits (*Anethum graveolens* L.). *Innov. Food Sci. Emer. Technol.* 2013; 17: 114-117.
- Cheng M, Kuo L, Ian T, Ih C. Two new sesquiterpenoids and ant-HIV principles from the root bark of *Zanthoxylum ailanthoides*. *Bioorg. Med. Chem.* 2005; 13: 5915-5920.
- Choi HJ, Song JH, Kwon DH, Baek SH, Ahn YJ. Antiviral activity of *Zanthoxylum* species against influenza virus. *Koran. J. Med. Crop Sci.* 2008; 16: 273-278.
- Choochote W, Chaithong U, Kamsuk K, Jitpakdi A, Tippawangkosol P, Tuetun B, Champakaew D, Pitasawat B. Repellent activity of selected essential oils against *Aedes aegypti*. *Fitoterapia.* 2007; 78: 359-364.
- Chun HS, Kim HJ, Ok HE, Hwang JB, Chung DH. Determination of aflatoxin levels in nuts and their products consumed in South Korea. *Food Chem.* 2007; 102: 385-391.
- Chung MS. Volatile compounds of *Zanthoxylum piperitum* A.P. DC. 2005. *Food Sci. Biotechnol.* 14: 529-532.
- Cowan MM. Plant products as antimicrobial agents. *Clin. Microbiol. Rev.* 1999; 12: 564-582.
- Delaquis PJ, Stanich K, Girard B, Mazza G. Antimicrobial activity of individual and mixed fractions of dill, cilantro, coriander and eucalyptus essential oils. *Int. J. Food Microbiol.* 2002; 74: 101-109.

- Devi KP, Nisha SA, Sakthivel R, Pandian SK. Eugenol (an essential oil of clove) acts as an antibacterial agent against *Salmonella typhiby* disrupting the cellular membrane. *J. Ethnopharmacol.* 2010; 130: 107-115.
- Diao WR, Hu QP, Zhang H, Xu JG. Chemical composition, antibacterial activity and mechanism of action of essential oil from seeds of fennel (*Foeniculum vulgare* Mill). *Food Control.* 2014; 35: 109-116.
- Diaz TML, Gonzalez CJ, Moreno B, Otero A. Effect of temperature, water activity, pH and some antimicrobials on the growth of *Penicillium olsonii* isolated from the surface of Spanish fermented meat sausage. *Food Microbiol.* 2002; 19: 1-7.
- Dikbas N, Kotan R, Dadasoglu F, Sahin F. Control of *Aspergillus flavus* with essential oil and methanol extract of *Satureja hortensis*. *Int. J. Food Microbiol.* 2008; 124: 179-182.
- Dimic GR, Kocic SD, Karalic TD. Growth inhibition of some *Eurotium* and *Aspergillus* species with spice extracts. Proceedings of the National Academy of Sciences of the United States of America. 2007; 113: 63-70.
- Doymaz I, Tugrul N, Pala M. Drying characteristics of dill and parsley leaves. *J. Food Eng.* 2006; 77: 559-565.
- Dupont S, Lemetais G, Ferreira T, Cayot P, Gervais P, Beney L. Eegosterol biosynthesis: A fungal pathway for life on land. *Evolution.* 2012; 66-9: 2961-2968.
- Elgayyar M, Dreughon FA, Golden DA, Mount JP. Antimicrobial activity of essential oils from plants against selected pathogenic and saprophytic microorganisms. *J. Food Protect.* 2001; 64: 1019-1024.
- Fatope MO, Marwah RG, Onifade AK, Ochei JE, Mahroqi YKSA. ¹³C NMR analysis and antifungal and insecticidal activities of Oman dill herb oil. *Pharmaceutical Biol.* 2006; 44: 44-4.
- Friedman M, Henika PR, Levin CE, Mandrell RE. Antibacterial activities of plant essential oils and their components against *Escherichia coli* O157:H7 and *Salmonella entericain* apple juice. *J. Agric. Food Chem.* 2004; 52: 6042-6048.

- Frisvad JC, Skouboe P, Samson RA. Taxonomic comparison of three different groups of aflatoxin producers and a new efficient producer of aflatoxin B₁, sterigmatocystin and 3-O-methylsterigmatocystin, *Aspergillus rambellii* sp. nov. *System. Appl. Microbiol.* 2005; 28: 442-453.
- Gill AO, Delaquis P, Russo P, Holley RA. Evaluation of antilisterial action of cilantro oil on vacuum packed ham. *Int. J. Food Microbiol.* 2002; 73: 83-92.
- Glisic SB, Milojevic SZ, Dimitrijevic SI, Orlovic AM, Skala DU. Antimicrobial activity of the essential oil and different fractions of *Juniperus communis* L. and a comparison with some commercial antibiotics. *J. Serb. Chem. Soc.* 2009; 72: 311-320.
- Gonza'lez SL, Vargas M, Marti'nez GC, Chiralt A, Cha'fer M. Use of essential oils in bioactive edible coatings. *Food Eng.* 2011; 3: 1-16.
- Gundidza M, Gweru N, Magwa ML, Mmbengwa V, Samie A. The chemical composition and biological activities of essential oil from the fresh leaves of *Schinus terebinthifolius* from Zimbabwe. *African J. Biotechnol.* 2009; 8: 7164-7169.
- Hitokoto HT, Morozumi S, Wauke T, Sakai S, Kurata H. Inhibitory effects of spices on growth and toxin production of toxigenic fungi. *Appl. Environ. Microbiol.* 1980; 39: 818-822.
- Hosseinzadeh H, Karimi GR, Ameri M. Effects of *Anethum graveolens* L. seed extracts on experimental gastric irritation models in mice. *BMC Pharmacol.* 2002; 2: 1-5.
- Hur JM, Park JG, Yang KH, Park JC, Park JR, Chun, SS, Choi JS, Choi JW. Effect of methanol extract of *Zanthoxylum piperitum* leaves and of its compound, protocatechuic acid, on hepatic drug metabolizing enzymes and lipid peroxidation in rats. *Biosci. Biotechnol. Biochem.* 2003; 67: 945-950.
- Hurtado RD, Garrido GG, Gonzalez SP, Iznaga Y, Gonzalez L, Torres JM, Curini M, Epifano F, Marcotullio MC. Antifungal activity of some Cuban *Zanthoxylum* species. *Fitoterapia.* 2003; 74: 384-386.

- Hwang ES, Kim GH. Safety evaluation of *Zanthoxylum piperitum*-derived essential oil by assessing micronucleus abnormalities, mutagenicity, and chromosomal aberration. *Food Res. Int.* 2012; 47: 267-271.
- Ibrahim B, Naser ZA. Analysis of fruits *Schinus molle* extractions and the efficacy in inhibition of growth the fungi in laboratory. *Int. J. Chem Tech. Res.* 2014; 6: 2799-2806.
- Iheshiulor OOM, Esonu BO, Chuwuka OK, Omede AA, Okoli IC, Ogbuewu IP. Effects of mycotoxins in animal nutrition: A review. *Asian J. Anim. sci.* 2011; 5: 19-33.
- Inouye S, Takizawa T, Yamaguchi H. Antibacterial activity of essential oils and their constituents against respiratory tract pathogens by gaseous contract. *J. Antimicrob. Chemother.* 2001; 47: 565-573.
- Inouye S, Takizawa T, Yamaguchi H. Inhibitory effect of essential oils on apical growth of *Aspergillus fumigatus* by vapour contact. *Mycoses.* 2000; 43: 17-23.
- Ittipanichpong C, Ruangrungsi N, Pattanaautsahakit C. Chemical compositions and pharmacological effect of essential oil from the fruit of *Zanthoxylum limonella*. *J. Med. Assoc. Thai.* 2002; 85: 344-353.
- Jang SY, Ryu SY, Kim SD. Antifungal activity of plant extracts against *Pityrosporum ovale* and *Candida albicans*. *Koran. J. Pharmacogno.* 2003; 34: 303-307.
- Jeong CH, Kwa, JH, Kim JH, Choi GN, Kim DO, Heo HJ. Neuronal cell protective and antioxidant effects of phenolics obtained from *Zanthoxylum piperitum* leaf using in vitro model system. *Food Chem.* 2011; 125: 417-422.
- Jiang J, Kubota K. Differences in the volatile components and their odor characteristics of green and ripe fruits and dried pericarp of Japanese Pepper (*Xanthoxylum piperitum* DC.). *J. Agric. Food Chem.* 2004; 52: 4197-4203.
- Joseph, GS, Jayaprakasha, GK, Selvi, AT, Jena, BS, Sakariah, KK. Antiaflatoxigenic and antioxidant activities of *Garcinia extracts*. *Int. J. Food Microbiol.* 2005; 101: 153-160.

- Joshi RK. Chemical composition and antimicrobial activity of the essential oil of *Plectranthus mollis* (Lamiaceae) from Western Ghats region, Karnataka, India. *Rev. Biol. Trop.* 2014; 62: 423-431.
- Kagan IA, Michel A, Prause A, Scheffler BE, Pace P, Duke SO. Gene transcription profiles of *Saccharomyces cerevisiae* after treatment with plant protection fungicides that inhibit ergosterol biosynthesis. *Pestic. Biochem. Physiol.* 2005; 82: 133-153.
- Kalemba D, Kunicka A. Antibacterial and antifungal properties of essential oils. *Curr. Med. Chem.* 2003; 10: 813-829.
- Kamsuk K, Choochote W, Chaithong U, Jitpakdi A, Tippawangkosol P, Riyong D, Pitasawat B. Effectiveness of *Zanthoxylum piperitum*-derived essential oil as an alternative repellent under laboratory and field applications. *Parasitol. Res.* 2007; 100: 339-345.
- Kedia A, Prakash B, Mishra PK, Dubey NK. Antifungal and antiaflatoxic properties of *Cuminum cyminum* (L.) seed essential oil and its efficacy as a preservative in stored commodities. *Int. J. Food Microbiol.* 2014; 168-169: 1-7.
- Khalaf AF. Toxicological efficacy of some indigenous dill compounds against the flesh fly. *Parasarcophaga dux* Thomson. *J. Egypt. Soc. Parasitol.* 2004; 34: 227-37.
- Khan MSA, Ahmad I. *In vitro* antifungal, anti-elastase and anti-keratinase activity of essential oils of *Cinnamomum*-, *Syzygium*- and *Cymbopogon*-species against *Aspergillus fumigatus* and *Trichophyton rubrum*. *Phytomedicine.* 2011; 19: 48-55.
- Kim JG. Anti-aflatoxic activity of some bacteria related with Fermentation. In: Méndez-Vilas A, editor. Communicating current research and educational topics and trends in applied microbiology. Badajoz: Formatex. 2007; 322-328.
- Klich MA. *Aspergillus flavus*: the major producer of aflatoxin. *Mol. Plant Pathol.* 2007; 8: 713-722.

- Kraikruan W, Sukprakarn, S, Mongkolporn O, Wasee S. Capsaicin and dihydropyridine contents of Thai chili cultivars. *Kasetsart J. (Nat. Sci.)*. 2008; 42: 611-616.
- Kumar SN, Mohandas C, Nambisan B. Purification of an antifungal compound, cyclo (l-Pro-d-Leu) for cereals produced by *Bacillus cereus* subsp. *thuringiensis* associated with entomopathogenic nematode. *Microbiol. Res.* 2013; 168: 278-288.
- Kusuda M, Inada K, Ogawa T, Yoshida T, Shiota S, Tsuchiya T, Hatano T. Polyphenolic constituent structures of *Zanthoxylum piperitum* fruit and the antibacterial effects of its polymeric procyanidin on methicillin-resistant *Staphylococcus aureus*. *Biosci. Biotechnol. Biochem.* 2006; 70: 1423-1431.
- Latge' JP, Beauvais A. Functional duality of the cell wall. *Curr. Opin. Microbiol.* 2014; 20: 111-117.
- Ling W. Health risk evaluation of pesticide contamination in drinking water. *Gesunde Pflangen*. 1991; 43: 21-25.
- Loopez P, Saanchez C, Batlle R, Nerian C. Solid- and vapor-phase antimicrobial activities of six essential oils: susceptibility of selected foodborne bacterial and fungal strains. *J. Agric. Food Chem.* 2005; 53: 6939-6946.
- Mahmoud, ALE. Antifungal action and antiaflatoxigenic properties of some essential oil constituents. *Lett. Appl. Microbiol.* 1994; 19: 110-113.
- Mello D. Mycotoxins. *Boletim científico*. 2000; 47: 1-7.
- Misra LN, Wouatsa NAV, Kumar S, Kumar RV, Tchoumboungang F. Antibacterial, cytotoxic activities and chemical composition of fruits of two Cameroonian *Zanthoxylum* species. *J. Ethnopharmacol.* 2013; 148: 74-80.
- Monsefi M, Ghasemi M, Bahaoddinu A. The effects of *Anethum graveolens* L. on female reproductive system of rats. *DARU J. Pharmaceutical Sci.* 2006; 14: 131-135.
- Mourey A, Canillac N. Anti-listeria monocytogenes activity of essential oils components of conifers. *Food Control*. 2002; 13: 289-292.

- Nes WD, Zhou W, Ganapathy K, Liu J, Vatsyayan R, Chamala S, Hernandez K, Miranda M. Sterol 24-C-methyltransferase: An enzymatic target for the disruption of ergosterol biosynthesis and homeostasis in *Cryptococcus neoformans*. *Arch. Biochem. Biophys.* 2009; 481: 210-218.
- Nieblas MMO, Burgueño MRR, Félix EA, León AG, Trejo AM, Moreno LV. Chemical composition and antimicrobial activity of oregano (*Lippia palmeri* S. WATS) essential oil. *Rev. Fitotec. Mex.* 2011; 34: 11-17.
- Nogueira JHC, Gonçalves E, Galletti SR, Facanali R, Marquesc MOM, Felício JD. *Ageratum conyzoides* essential oil as aflatoxin suppressor of *Aspergillus flavus*. *Int. J. Food Microbiol.* 2010; 137: 55-60.
- Omran SM, Moodi MA, Mohammad S, Amiri BN, Mosavi SJ, Ghazi SAM, Saeed M, Shiade SMJ, Kheradi E, Salehi M. The Effects of Limonene and Orange Peel Extracts on Some Spoilage Fungi. *Int. J. Mol. Clin. Microbiol.* 2011; 1: 82-86.
- Orhan IE, Senol FS, Ozturk N, Celik SA, Pular A, Kan Y. Phytochemical contents and enzyme inhibitory and antioxidant properties of *Anethum graveolens* L. (dill) samples cultivated under organic and conventional agricultural conditions. *Food Chem Toxicol.* 2013; 59: 93-106.
- Ouazzou AA, Cherrat L, Espina L, Lorán S, Rota C, Pagán R. The antimicrobial activity of hydrophobic essential oil constituents acting alone or in combined processes of food preservation. *Innovative Food Sci. Emer. Technol.* 2011; 12: 320-329.
- Park HS, Jun DY, Fang Z, Woo MH, Kim YH. Antimicrobial activity from seeds of *Zanthoxylum piperitum* against oral pathogen *Streptococcus mutans*. *J. life Sci.* 2008; 18: 167-174.
- Park YD, Lee WS, An S, Jeong TS. Human acyl-Co A: Cholesterol acyltransferase inhibitory activities of aliphatic acid amides from *Zanthoxylum piperitum* DC. *Biol. Pharm. Bull.* 2007; 30: 205-207.
- Pauli A. Antimicrobial properties of essential oil constituents. *Int. J. Aromather.* 2001; 3: 126-133.

- Peter KV. Handbook of herbs and spices Volume 2. 2004. Woodhead Publishing Limited.
- Peter KV. Handbook of herbs and spices Volume 3. 2006. Woodhead Publishing Limited.
- Pinto E, Vale-Silva L, Cavaleiro C, Salgueiro L. Antifungal activity of the clove essential oil from *Syzygium aromaticum* on *Candida*, *Aspergillus* and dermatophyte species. *J. Med. Microbiol.* 2009; 58: 1454-1462.
- Prakash B, Mishra PK, Kedia A, Dubey NK. Antifungal, antiaflatoxin and antioxidant potential of chemically characterized *Boswellia carterii* Birdw essential oil and its in vivo practical applicability in preservation of *Piper nigrum* L. fruits. *LWT - Food Sci. Techno.* 2014; 56: 240-247.
- Prakash B, Singh P, Kedia A, Dubey NK. Assessment of some essential oils as food preservatives based on antifungal, antiaflatoxin, antioxidant activities and in vivo efficacy in food system. *Food Res. Int.* 2012; 49: 201-208.
- Prakash B, Singh P, Mishra PK, Dubey NK. Safety assessment of *Zanthoxylum alatum* Roxb. essential oil, its antifungal, antiaflatoxin, antioxidant activity and efficacy as antimicrobial in preservation of *Piper nigrum* L. fruits. *Int. J. Food Microbiol.* 2012; 153: 183-191.
- Przygodzka M, Zielinska D, Ciesarová Z, Kukurová K, Zielinski H. Comparison of methods for evaluation of the antioxidant capacity and phenolic compounds in common spices. *LWT - Food Sci. Techno.* 2014; 58: 321-326.
- Radulescu V, Popescu M.L, Ilies D.C. Chemical composition of the volatile oil from different plant parts of *Anethum graveolens* L. (Umbelliferae) cultivated in Romania. *Farmacia.* 2010; 58: 594-600.
- Raghavan S. Handbook of spices, Seasonings and Flavorings second edition. Taylor & Francis Group, LLC. 2007.
- Rahman, A, Kang S.C. In vitro control of food-borne and food spoilage bacteria by essential oil and ethanol extracts of *Lonicera japonica* Thunb. *Food Chem.* 2009; 116: 670-675.

- Regnier T, Plooy W, Combrinck S, Botha B. Fungitoxicity of *Lippia scaberrima* essential oil and selected terpenoid components on two mango postharvest spoilage pathogens. *Postharv. Biol. Technolo.* 2008; 48: 254-258.
- Rodrigues P, Soares C, Kozakiewicz Z, Paterson RRM, Lima N, Venâncio A. Identification and characterization of *Aspergillus flavus* and aflatoxins. Communicating Current Research and Educational Topics and Trends in Applied Microbiology. A. Méndez-Vilas (Ed.). 2007. 527-534.
- Rogg LE, Fortwende JR, Juvvadi PR, Lilley A, Steinbach WJ. The chitin synthase genes *chsA* and *chsC* are not required for cell wall stress responses in the human pathogen *Aspergillus fumigatus*. *Biochem. Biophys. Res. Commun.* 2011; 411: 549-554.
- Saggdic O, Ozcan M. Antibacterial activity of Turkish spice hydrosols. *Food Control.* 2003; 14: 141-143.
- Sardiñas N, Serna JG, Santos L, Ramos AJ, Jaén MTG, Patiño B, Vázquez C. Detection of potentially mycotoxigenic *Aspergillus* species in *Capsicum* powder by a highly sensitive PCR-based detection method. *J. Food control.* 2011; 22: 1363-1366.
- Schachtschabel D, Arentshorst M, Lagendijk EL, Ram AFJ. Vacuolar H⁺-ATPase plays a key role in cell wall biosynthesis of *Aspergillus niger*. *Fungal Genet. Biol.* 2012; 49: 284-293.
- Senatore F, Oliviero F, Scandolera E, Scafati T.O, Roscigno G, Zaccardelli M, Falco E.D. Chemical composition, antimicrobial and antioxidant activities of anethole-rich oil from leaves of selected varieties of fennel [*Foeniculum vulgare* Mill. ssp.vulgare var. azoricum (Mill.) Thell]. *Fitoterapia.* 2013; 90: 214-219.
- Sharma N, Tripathi A. Effects of *Citrus sinensis* (L.) Osbeck epicarp essential oil on growth and morphogenesis of *Aspergillus niger* (L.) Van Tieghem. *Microbiol. Res.* 2006; 163: 337-344.
- Shukla R, Singh P, Prakash B, Dubey N.K. Antifungal, aflatoxin inhibition and antioxidant activity of *Callistemon lanceolatus* (Sm.) sweet essential oil and its major component 1,8-cineole against fungal isolates from chickpea seeds. *J. Food Control.* 2012; 25: 27-33.

- Shyu YS, Lin JT, Chang YT, Chiang CJ, Yang DJ. Evaluation of antioxidant ability of ethanolic extract from dill (*Anethum graveolens* L.) flower. *Food Chem.* 2009; 115: 515-521.
- Silva F, Ferreira S, Duarte A, Mendonca ID, Domingues CF. Antifungal activity of *Coriandrum sativum* essential oil, its mode of action against *Candida* species and potential synergism with amphotericin B. *Phytomedicine.* 2011; 19: 42-47.
- Silveira SMD, Júnior AC, Scheuermann GN, Secchi FL, Vieira CRW. Chemical composition and antimicrobial activity of essential oils from selected herbs cultivated in the South of Brazil against food spoilage and foodborne pathogens. *Ciência Rural, Santa Maria.* 2012; 42: 1300-1306.
- Singh G, Maurya S, De Lampasona M.P, Catalan C. Chemical constituents, antimicrobial investigations, and antioxidative potentials of *Anethum graveolens* L. essential oil and acetone extract. *J. Food Sci.* 2005; 70: 208-215.
- Soliman KM, Badea RI. Effect of oil extracted from some medicinal plants on different mycotoxigenic fungi. *Food Chem Toxicol.* 2002; 40: 1669–1675.
- Sreelekha M, Anto AP, Anto RJ, Shafi PM. Cytotoxicity of 6-acetonyl dihydrochelerythrin, arnottianamide and 6-(2-hydroxypropyl)-dihydrochelerythrine towards human cancer cell lines. *Indian J. Chem.* 2014; 53: 647-651.
- Tangjitjaroenkun J, Chantarasriwong O, Chavasiri W. Chemical constituents of the stems of *Zanthoxylum limonella* Alston. *Phytochem Lett.* 2012; 5: 443-445.
- Tangjitjaroenkun J, Supabphol R, Chavasiri W. Antioxidant effect of *Zanthoxylum limonella* Alston. *J. Med. Plants Res.* 2012; 6: 1407-1414.
- Tanruean K, Kaewnarin K, Rakariyatham N. Antibacterial and Antioxidant Activities of *Anethum graveolens* L. Dried Fruit Extracts. *Chiang Mai J. Sci.* 2014; 41: 649-660.
- Tatsadjieu LN, Ngang JJE, Ngassoum MB, Etoa FX. Antibacterial and antifungal activity of *Xylopi aethiopica*, *Monodora myristica*, *Zanthoxylum xanthoxylodes* and *Zanthoxylum leprieurii* from Cameroon. *Fitoterapia.* 2003; 74: 469-472.

- Tatsadjieu NL, Dongmo PMJ, Ngassoum MB, Etoa FX, Mbofung CMF. Investigations on the essential oil of *Lippia rugosa* from Cameroon for its potential use as antifungal agent against *Aspergillus flavus* Link ex. Fries. *Food Control*. 2009; 20: 161-166.
- Thakaew R, Niamsup H. Inhibitory activity of *Bacillus subtilis* BCC 6327 metabolites against growth of aflatoxigenic fungi isolated from bird chili powder. *Int. J. Biosci. Biochem. Bioinforma*. 2013; 3: 27-32.
- Tian J, Ban X, Zeng H, Huang B, He J, Wang Y. In vitro and in vivo activity of essential oil from dill (*Anethum graveolens* L.) against fungal spoilage of cherry tomatoes. *Food Control*. 2011; 22: 1992-1999.
- Tian J, Huang B, Luo X, Zang H, Ban BX, He J, Wang Y. The control of *Aspergillus flavus* with *Cinnamomum jensenianum* Hand.-Mazz essential oil and its potential use as a food preservative. *Food Chem*. 2012; 130: 520-527.
- Tolouee M, Alinezhad S, Saberi R, Eslamifar A, Zad SJ, Jaimand K, et al. Effect of *Matricaria chamomilla* L. flower essential oil on the growth and ultrastructure of *Aspergillus niger* van Tieghem. *Int. J. Food Microbiol*. 2010; 139: 127-133.
- Tripathi A, Sharma N, Sharma V. In vitro efficacy of *Hyptis suaveolens* L. (Poit.) essential oil on growth and morphogenesis of *Fusarium oxysporum* f.sp gladioli (Massey) Snyder & Hansen. *World J. Microbiol. Biotechnol*. 2009; 25: 503-512.
- Tzortzakis NG. Impact of cinnamon oil-enrichment on microbial spoilage of fresh produce. *Innov. Food Sci. Emer. Technol*. 2009; 10: 97-102.
- Uhl SR. Handbook of spices, Seasonings and Flavorings. Technomic Publishing Company. Lancaster. 2000: 62-64.
- Veen M, Lang C. Interactions of the ergosterol biosynthetic pathway with other lipid pathways. *Biochem. Soc. Transact*. 2005; 33: 1178-1181.
- Victório CP, Alviano DS, Alviano CS, Lage CLS. Chemical composition of the fractions of leaf oil of *Alpinia zerumbet* (Pers.) B.L. Burtt & R.M. Sm. and antimicrobial activity. *Braz. J. Pharmacog*. 2008; 19: 697-701.

- Wangcharoen W, Morasuk W. Antioxidant capacity changes of bird chili (*Capsicum frutescens* Linn.) during hot air drying. *Kasetsart J. (Nat. Sci.)*. 2009; 43: 12-20.
- Weiss EA. Spice crops. CABI Publishing. 2002.
- Wu F, Narrod C, Tiongco M, Liu Y. The Health economics of aflatoxin: global burden of disease. *Int. Food Policy Res. Institute*. 2011; 1-16.
- Yamazaki E, Inagaki M, Kurita O, Inoue T. Antioxidant activity of Japanese pepper (*Zanthoxylum piperitum* DC.) fruit. *Food Chem*. 2007; 100: 171-177.
- Yang VW, Clausen CA. Antifungal effect of essential oils on southern yellow pine. *Int. Biodeterior. Biodegrad*. 2007; 59: 302-306.
- Yang Y.C, Park K, Kim E.H, Lee H.S, Ahn Y.J. Larvicidal activity of medicinal plant extracts against *Aedes aegypti*, *Ochlerotatus togoi*, and *Culex pipiens pallens* (Diptera: Culicidae). *J. Asia-Pacific. Entomol*. 2004; 7: 227-232.
- Yazdanparast R, Bahramikia S. Evaluation of the effect of *Anethum graveolens* L. crude extracts on serum lipids and lipoproteins profiles in hypercholesterolaemic rats. *DARU J. Pharmaceutical Sci*. 2008; 16: 88-94.
- Yeboa NFAB, Arnason JT, Baker J, Smith ML. Antifungal constituents of Northern prickly ash, *Zanthoxylum americanum* Mill. *Phytomedicine*. 2005; 12: 370-377.
- Yun Y, Yin D, Dawood DH, Liu X, Chen Y, Ma Z. Functional characterization of *FgERG3* and *FgERG5* associated with ergosterol biosynthesis, vegetative differentiation and virulence of *Fusarium graminearum*. *Fungal Genet. Biol*. 2014; 68: 60-70.
- Zain ME. Impact of mycotoxins on humans and animals. *J. Saudi Chem Soc*. 2011; 15: 129-144.
- Zambonelli A, Aulerio AZ, Bianchi A, Alsabini A. Effect of essential oils on phytopathogenic fungi in vitro. *J. Phytopathol*. 1996; 144: 491-494.
- Zheng G, Kenney PM, Lam LKT. Anethofuran, Carvone and Limonene: Potential cancer chemopreventive agents from dill weed oil and caraway oil. *Planta Med*. 1992; 58: 339-341.