

REFERENCES

- [1] Hosni, K., Hassen, I., Chaabane, H., Jemli, M., Dallali, S., Sebei, H., and Casabianca, H. 2013. Enzyme-assisted extraction of essential oils from thyme (*Thymus capitatus* L.) and rosemary (*Rosmarinus officinalis* L.): Impact on yield, chemical composition and antimicrobial activity. *Industrial Crops and Products*, 47, 291-299.
- [2] Politeo, O., Jukic, M., and Milos, M. 2007. Chemical composition and antioxidant capacity of free volatile aglycones from basil (*Ocimum basilicum* L.) compared with its essential oil. *Food Chemistry*, 101, 379-385.
- [3] Bouaziz, M., Yangui, T., Sayadi, S., and Dhouib, A. 2009. Disinfectant properties of essential oils from *Salvia officinalis* L. cultivated in Tunisia. *Food and Chemical Toxicology*, 47, 2755-2760.
- [4] Tsai, T-H., Tsai, T-H., Wu, W-H., Tseng, J. T-P., and Tsai, P-J. 2010. *In vitro* antimicrobial and anti-inflammatory effects of herbs against *Propionibacterium acnes*. *Food Chemistry*, 119, 964-968
- [5] Singha-Aussavarat, P. 2009. Antioxidant activity, antibacterial activity against acne-inducing bacteria and HPLC analysis of green tea extract and its cream product. Master's thesis. Chiang Mai University, Chiang Mai.
- [6] Sukatta, U., Rugthaworn, P., Leebunyanon, M., and Pitpiangchan, P. 2010. Bioactive compounds content, free radical scavenging and anti-acne inducing bacteria activities of some fruit peels extract. *Proceedings of the 48th Kasetsart University Annual Conference*, Bangkok, Thailand, pp. 364-373.

- [7] Daud, F. S., Pande, G., Joshi, M., Pathak, R., and Wankhede, S. 2013. A study of antibacterial effect of some selected essential oils and medicinal herbs against acne causing bacteria. *International Journal of Pharmaceutical Science Invention*, 2, 27-34.
- [8] Kalembe, D., and Kunicka, A. 2003. Antibacterial and antifungal properties of essential oils. *Current Medicinal Chemistry*, 10, 813-829.
- [9] Liu, A-L., Lee, S. M. Y., Wang, Y-T., and Du, G-H. 2007. *Elsholtzia*: review of traditional uses, chemistry and pharmacology. *Chinese Pharmaceutical Sciences*, 16, 73-78.
- [10] Arbab, I. A., Abdul, A. B., Aspollah, M., Abdullah, R., Abdelwahab, S. I., Mohan, S., and Abdelmageed, A. H. A. 2011. *Clausena excavata* Burm. f. (Rutaceae): a review of its traditional uses, pharmacological and phytochemical properties. *Journal of Medicinal Plants Research*, 5, 7177-7184.
- [11] Guo, Z., Liu, Z., Wang, X., Liu, W., Jiang, R., Cheng, R., and She, G. 2012. *Elsholtzia*: phytochemistry and biological activities. *Chemistry Central Journal*, 6:147, 1-8.
- [12] Zouari, N., Fakhfakh, N., Zouari, S., Bougatef, A., Karray, A., Neffati, M., and Ayadi, M.A. 2011. Chemical composition, angiotensin I-converting enzyme inhibitory, antioxidant and antimicrobial activities of essential oil of Tunisian *Thymus algeriensis* Boiss. et Reut. (Lamiaceae). *Food and Bioproducts Processing*, 89, 257-265.
- [13] Goren, A. C., Piozzi, F., Akcicek, E., Kilic, T., Carikci, S., Mozioglu, E., and Setzer, W. N. 2011. Essential oil composition of twenty-two *Stachys* species (mountain tea) and their biological activities. *Phytochemistry Letters*, 4, 448-453.

- [14] Baser, K. H. C., Buchbauer, G. 2010. Hand book of ESSENTIAL OILS Science, Technology, and Applications, Taylor and Francis Group, LLC, pp. 95-140.
- [15] Bouayed, J., and Bohn, T. 2012. Nutrition, Well-Being and Health, InTech, Croatia, pp.160-165.
- [16] Handa, S. S., Khanuja, S. P. S., Longo, G., and Rakesh, D. D. 2008. Extraction Technologies for Medicinal and Aromatic Plant, United Nations Industrial Development Organization and the International Centre for Science and High Technology (ICS-UNIDO), Trieste, pp.31-51.
- [17] [Online]. “Enfleurage”, Retrieved April 20, 2014, from <http://www.aromatherapy-school.com/essential-oils/enfleurage.html>
- [18] Gerhards, P., Bons, U., Sawazki, J., Szigan, J., and Wertmann, A. 1999. GC/MS in Clinical Chemistry, Wiley-VCH, Weinheim, pp.22-24.
- [19] [Online]. “Gas Chromatography Mass Spectrometry (GC/MS)”, Retrieved April 20, 2014, from <http://www.bris.ac.uk/nerclsmsf/techniques/gcms.html>
- [20] Heberger, K., Gorgenyi, M., and Kowalska, T. 2002. Temperature dependence of Kovat indices in gas chromatography revisited. *Journal of Chromatography A*, 973, 135-142.
- [21] Adams, R. P. 2007. Identification of essential oil components by gas chromatography/mass spectrometry, 4th Edition, Allured, Carol Stream, Illinois.
- [22] Lee, C-J., Chen, L-G., Chang, T-L., Ke, W-M., Lo, Y-F., and Wang, C-C. 2011. The correlation between skin-care effects and phytochemical contents in Lamiaceae plants. *Food Chemistry*, 124, 833-841.

- [23] Gill, A. K., De, A., and Giri, P. D. 2013. Nature for the treatment of acne vulgaris: The disease of adolescence. *World Journal of Pharmacy and Pharmaceutical Sciences*, 2, 1739-1766.
- [24] Azimi, H., Fallah-Tafti, M., Khakshur, A. A., and Abdollahi, M. 2012. A review of phytotherapy of acne vulgaris: Perspective of new pharmacological treatments. *Fitoterapia*, 83, 1306-1317.
- [25] Ghosh, V. K., Nagore, D. H., and Kadbhane, K. P. 2011. Different approaches of alternative medicines in *acne vulgaris* treatment. *Oriental Pharmacy and Experimental Medicine*, 11, 1-9.
- [26] Atlas, R. M. 1984. Microbiology: Fundamentals and applications, Macmillan, New York, pp. 649-652.
- [27] Presterl, E., Suchomel, M., Eder, M., Reichmann, S., Lassnigg, A., Graninger, W., Rotter, M. 2007. Effects of alcohols, povidone-iodine and hydrogen peroxide on biofilms of *Staphylococcus epidermidis*. *Journal of Antimicrobial Chemotherapy*, 60, 417-420.
- [28] Zu, Y., Yu, H., Liang, L., Fu, Y., Efferth, T., Liu, X., and Wu, N. 2010. Activities of ten essential oils towards *Propionibacterium acnes* and PC-3, A-549 and MCF-7 Cancer Cells. *Molecules*, 15, 3200-3210.
- [29] Upadhyay, R. K., Dwivedi, P., and Ahmad, S. 2010. Screening of antibacterial activity of six plant essential oils against pathogenic bacterial strains. *Asian Journal of Medical Sciences*, 2, 152-158.
- [30] Kim, S-S., Baik, J. S., Oh, T-H., Yoon, W-J., Lee, N. H., and Hyun, C-G. 2008. Biological activities of Korean *Citrus obovoides* and *Citrus natsudaoides* essential oils against acne-inducing bacteria. *Bioscience Biotechnology and Biochemistry*, 72, 2507-2513.

- [31] Viyoch, J., Pisutthanan, N., Faikreua, A., Nupangta, K., Wangtorpol, K., and Ngokkuen, J. 2006. Evaluation of *in vitro* antimicrobial activity of Thai basil oils and their micro-emulsion formulas against *Propionibacterium acnes*. *International Journal of Cosmetic Science*, 28, 125-133.
- [32] Jiang, L. 2011. Comparison of disk diffusion, agar dilution, and broth micro-dilution for antimicrobial susceptibility testing of five chitosans. Master's thesis. Louisiana State University, U.S.
- [33] Seekhaw, P. 2010. The chemical constituents and biological activities of the leaves of *Clausena harmandiana* (Pierre). Master's thesis. Ramkhamhaeng University, Bangkok.
- [34] Thongthoom, T., Songsiang, U., Phaosiri, C., and Yenjai, C. 2010. Biological activity of chemical constituents from *Clausena harmandiana*. *Archives of Pharmacal Research*, 33, 675-680.
- [35] Chokeprasert, P. 2006. Studies on development of mafai jeen [*Clausena lansium* (Lour.) Skeels] products. Ph.D.'s thesis. Kasetsart University, Bangkok.
- [36] Adebajo, A.C., Iwalewa, E.O., Obuotor, E.M., Ibikunle, G.F., Omisore, N.O., Adewunmi, C.O., Obaparusi, O.O., Klaes, M., Adetogun, G.E., Schmidt, T.J., and Verspohl, E.J. 2009. Pharmacological properties of the extract and some isolated compounds of *Clausena lansium* stem bark: Anti-trichomonal, antidiabetic, anti-inflammatory, hepatoprotective and antioxidant effect. *Journal of Ethnopharmacology*, 122, 10-19.
- [37] Prasad K. N., Xie, H., Hao, J., Yang, B., Qiu, S., Wei, X., Chen, F., and Jiang, Y. 2010. Antioxidant and anticancer activities of 8-hydroxypsoralen isolated from wampee [*Clausena lansium* (Lour.) Skeels] peel. *Food Chemistry*, 118, 62-66.


- [38] Han, Y., Li, L., Hao, W., Tang, M., and Wan, S. 2013. Larvicidal activity of lansiumamide B from the seeds of *Clausena lansium* against *Aedes albopictus* (Diptera: Culicidae). *Parasitology Research*, 112, 511-516.
- [39] Songsiang, U., Thongthoom, T., Zeekpudsa, P., Kukongviriyapan, V., Boonyarat, C., Wangboonskul, J., and Yenjai, C. 2012. Antioxidant activity and cytotoxicity against cholangiocarcinoma of carbazoles and coumarins from *Clausena harmandiana*. *ScienceAsia*, 38, 75-81.
- [40] Sriphana, U., Thongsri, Y., Prariyachatigul, C., Pakawatchai, C., and Yenjai, C. 2013. Clauraila E from the roots of *Clausena harmandiana* and antifungal activity against *Pythium insidiosum*. *Archives of Pharmacal Research*, 36, 1078-1083.
- [41] Maneerat, W., Phakhodee, W., Ritthiwigrom, T., Cheenpracha, S., Deachathai, S., Laphookhieo, S. 2013. Phenylpropanoid derivatives from *Clausena harmandiana* fruits. *Phytochemistry Letters*, 6, 18-20.
- [42] Shen, D-Y., Chao, C-H., Chan, H-H., Huang, G-J., and Hwang, T-L. 2012. Bioactive constituents of *Clausena lansium* and a method for discrimination of aldose enantiomers. *Phytochemistry*, 82, 110-117.
- [43] Rajkaran, S., and Jebanesan, A. 2010. Chemical composition and larvicidal activity of leaf essential oil from *Clausena dentata* (Willd) M. Roam. (Rutaceae) against the chikungunya vector, *Aedes aegypti* Linn. (Diptera: Culicidae). *Journal of Asia-Pacific Entomology*, 13, 107-109.
- [44] Cheng, S-S., Chang, H-T., Lin, C-Y., Chen, P-S., Huang, C-G., Chen, W-J., and Chang S-T. 2009. Insecticidal activities of leaf and twig essential oils from *Clausena excavata* against *Aedes aegypti* and *Aedes albopictus* larvae. *Pest Management Science*, 65, 339-343.

- [45] Rahman, Md. A., Chakma, J. S., Bhuiyan, N. I., and Islam, Md. S. 2012. Composition of the essential oil of *Clausena Suffruticosa* leaf and evaluation of its antimicrobial and cytotoxic activities. *Tropical Journal of Pharmaceutical Research*, 11, 739-746.
- [46] Pandey, A. K., Singh, P., Mohan, M., and Tripathi, N. N. 2012. New report on the chemical composition of the essential oil from leaves of *Clausena pentaphylla* from India. *Chemistry of Natural Compounds*, 48, 896-897.
- [47] Govindarajan, M. 2010. Chemical composition and larvicidal activity of leaf essential oil from *Clausena anisata* (Willd.) Hook. F. ex Benth (Rutaceae) against three mosquito species. *Asian Pacific Journal of Tropical Medicine*, 3, 874-877.
- [48] Usman, L.A., Hamid, A.A., Olawore, N.O., Fakunle, C.O., Oladosu, I.A., and Zubair, M.F. 2010. Chemical composition of leaf essential oil of *Clausena anisata* growing in north-central Nigeria. *Journal of Applied Science Research*, 6, 891-894.
- [49] Zhao, J., Nan, P., and Zhong, Y. 2004. Chemical composition of the essential oils of *Clausena lansium* from Hainan Island, China. *Z. Naturforsch*, 59c, 153-156.
- [50] Wu, C. Y., Raven, P. H., and Hong, D. Y. 1994. Flora of China Editorial Committee. Science Press & Missouri Botanical Garden Press, Beijing & St. Louis, 17, 1-378.
- [51] Bongcheewin, B., Chantaranothai, P. 2008. Two new records of *Elsholtzia* Willd. (Lamiaceae) for Thailand. *The Natural History Journal of Chulalongkorn University*, 8, 1-5.

- [52] Kim, S-S., Oh, H. J., Baik, J. S., Oh, T-H., Yun, P-Y., Kim, C-S., Lee, N. H., and Hyun, C-G. 2008. Chemical composition and biological activities of *Elsholtzia splendens* essential oil. *Journal of Applied Biological Chemistry*, 51, 69-72.
- [53] Guang-hui, T. 2013. Chemical constituents in essential oils from *Elsholtzia ciliata* and their antimicrobial activities. *Chinese Herbal Medicines*, 5, 104-108.
- [54] Lorian, V. 1991. Antibiotics in laboratory medicine. Third edition, Maryland, Williams & Wilkins, pp.28-34.
- [55] Barry, A. L. 1976. The antimicrobial susceptibility test: Principles and practices. Philadelphia, Lea & Febiger, pp.182-187.
- [56] Martins, M. D. R., Arantes, S., Candeias, F., Tinoco, M. T., and Cruz-Morais, J. 2014. Antioxidant, antimicrobial and toxicological properties of *Schinus molle* L. essential oils. *Journal of Ethnopharmacology*, 151, 485-492.
- [57] Baker, F. J., and Breach, M. R. 1980. Medical Microbiological Techniques. Butterworth & Co., Ltd., pp.343-350.
- [58] Houghton, P. J., and Raman, A. 1998. Laboratory Handbook for the Fractionation of Natural Extracts. Chapman & Hall, pp.22-31.
- [59] Vimalanathan, S., and Hudson, J. 2012. Anti-influenza virus activities of commercial Oregano oils and their carriers. *Journal of Applied Pharmaceutical Science*, 02, 214-218.

- [60] Economou, G., Panagopoulos, G., Karamanos, A., Tarantilis, P., Kalivas, D., and Kotoulas, V. 2014. An assessment of the behavior of carvacrol-rich wild *Lamiaceae* species from the eastern Aegean under cultivation in two different environments. *Industrial Crops and Products*, 54, 62-69.
- [61] Abbaszadeh, S., Sharifzadeh, A., Shokri, H., Khosravi, A.R., and Abbaszadeh, A. 2014. Antifungal efficacy of thymol, carvacrol, eugenol and menthol as alternative agents to control the growth of food-relevant fungi. *Journal of Medical Mycology*, Available online 26 February 2014, <http://dx.doi.org/10.1016/j.mycmed.2014.01.063>
- [62] Lewinsohn, E., Dudai, N., Tadmor, Y., Katzir, I., Ravid, U., Putievsky, E., and Joel, D. M. 1998. Histochemical localization of citral accumulation in lemongrass leaves (*Cymbopogon citratus* (DC.) Stapf., Poaceae) *Annals of Botany*, 81, 35-39.
- [63] Mesa-Arango, A. C., Montiel-Ramos, J., Zapata, B., Durán, C., Betancur-Galvis, L., and Stashenko, E. 2009. Citral and carvone chemotypes from the essential oils of *Colombian Lippia alba* (Mill.) N.E. Brown: composition, cytotoxicity and antifungal activity. *Memórias do Instituto Oswaldo Cruz*, 104, 878-884.
- [64] Farhath, M. S. S., Vijaya, P. P., and Vimal, M. 2013. Antioxidant activity of Geraniol, Geranial acetate, Gingerol and Eugenol. *Research in Pharmacy* 3, 1-6.
- [65] Panyakaew, J., 2013. Compositional analysis and antioxidant activities of essential oils from genus *Elsholtzia*. Master's thesis. Chiang Mai University, Chiang Mai.
- [66] Verma, R. S., Padalia, R. C., Chauhan, A., and Thul, S. T. 2013. Exploring compositional diversity in the essential oils of 34 *Ocimum* taxa from Indian flora. *Industrial Crops and Products*, 45, 7-19.

- [67] Mathela, C. S., Padalia, R. C., Joshi, S. C., Singh, K. K., Bisht, D. S., Pant, A. K., and Kharkwal, H. 2009. Chemical diversity in Himalayan *Elsholtzia* species. *Chemiatry and Biodiversity*, 6, 2217-2226.
- [68] Rana, V. S., Devi, L. R., Verdeguer, M., and Blazquez, M. A. 2012. *Elsholtzia blanda* Benth: New citral-rich chemotypes from India. *Journal of Herbs, Spices Medicinal Plants*, 18, 132-139.
- [69] Chokeprasert, P., Charles, A. L., Sue, K-H., and Huang, T-C. 2007. Volatile components of the leaves, fruits and seeds of wampee [*Clausena lansium* (Lour.) Skeels]. *Journal of Food Composition and Analysis*, 20, 52-56.
- [70] Nakatsuji, T., Kao, M. C., Fang, J-Y., Zouboulis, C. C., Zhang, L., Gallo, R. L., and Huang, C-M. 2009. Antimicrobial property of lauric acid against *Propionibacterium acnes*: Its therapeutic potential for inflammatory acne vulgaris. *Journal of Investigative Dermatology*, 129, 2480-2488.
- [71] Huang, W-C., Tsai, T-H., Chuang, L-T., Li, Y-Y., Zouboulis, C. C., and Tsai, P. J. 2014. Anti-bacterial and anti-inflammatory properties of capric acid against *Propionibacterium acnes*: A comparative study with lauric acid. *Journal of Dermatological Science*, 73, 232-240.
- [72] Marino, M., Bersani, C., and Comi, G. 2001. Impedance measurements to study the antimicrobial activity of essential oils from *Lamiaceae* and *Compositae*. *International Journal of Food Microbiology*, 67, 187-195.
- [73] Hossain, M. A., Ismail, Z., Rahman, A., and Kang, S. C. 2008. Chemical composition and anti-fungal properties of the essential oils and crude extracts of *Orthosiphon stamineus* Benth. *Industrial Crops and Products*, 27, 328-334.

- 
- The background of the page features a large, faint watermark of the Chiang Mai University seal. The seal is circular, containing a central emblem with a torch and a book, surrounded by Thai script and the English text "CHIANG MAI UNIVERSITY 1964".
- [74] Gill, A.O., Delaquis, P., Russo, P., Holley, R. A. 2002. Evaluation of antilisterial action of cilantro oil vacuum packed ham. *International Journal of Food Microbiology*, 73, 83-92.
- [75] Bury, S. 2004. Essential oils: their antibacterial properties and potential applications in foods-a review. *International Journal of Food Microbiology*, 94, 223-253.
- [76] Lambert, R. J. W., Skandamis, P. N., Coote, P., Nychas, G.-J. E., 2001. A study of the minimum inhibitory concentration and mode of action of oregano essential oil, thymol and carvacrol. *Journal of Applied Microbiology*, 91, 453-462.
- [77] Ballester-Costa, C., Sendra, E., Fernandez-Lopez, J., Perez-Alvarez, J. A., and Viuda-Martos, M. 2013. Chemical composition and *in vitro* antibacterial properties of essential oils of four *Thymus* species from organic growth. *Industrial Crops and Products*, 50, 304-311.