

REFERENCES

- [1] Jun, H., Sheng-Ping, Y., Bo-Jun, X., Shang-Gao, L., Li-Ping, L., Jian, D., Jian-Min, Y., (2008) “Cytotoxic sesquiterpenoids from *Vernonia bockiana*”, *J. Asian. Nat. Prod. Res.*, 10, 571–575.
- [2] Alembert, T. T., Pierre, T., Johnson, F. A., Joseph, D. C., (2003) “Stigmastane derivatives and isovaleryl sucrose esters from *Vernonia guineensis* (Asteraceae)”, *Phytochemistry*, 63, 841–846.
- [3] Olha, K., Victor, C., Renato, M., Luis, P., Irmgard, M., (2006) “*Ent*-kaurane glycosides and sesquiterpene lactones of the hirsutinolide type from *Vernonia triflosculosa*”, *Phytochemistry*, 67, 62–69.
- [4] Erasto, P., Grierson, D. S., Afolayan, A. J., (2006) “Bioactive sesquiterpene lactones from the leaves of *Vernonia amygdalina*”, *J. Ethnopharmacol.*, 106, 117–120.
- [5] http://zipcodezoo.com/Key/Plantae/Vernonia_Genus.asp (August 2013)
- [6] <http://plants.jstor.org/specimen/k000814667> (June 2014)
- [7] Peter, B. K., Leland, J. C. S. W., James, A. D., Harry, L. B. (1999). Natural products from plants, *Phytochemicals: The chemical components of plants* (pp.2–23), New York: CRC Press.
- [8] Jose, N. B., Joao, L., Callegari, L., Walter, V., Diones, A. D., Kenji, N., Wilson, R.C., Werner, H., (1997) “Cadinanolides and other constituents from *Vernonia fruticulosa* and *Vernonanthura discolor*”, *Phytochemistry*, 44, 1535–1536.
- [9] Rabe, T., Mullholland, D., van Staden, J., (2002) “Isolation and identification of antibacterial compounds from *Vernonia colorata* leaves”, *J. Ethnopharmacol.*, 80, 91–94.

- [10] Chukwujekwu, J.C., Lategan, C.A., Smith, P.J., Van Heerden, F.R., Van Staden, J., (2009) “Antiplasmodial and cytotoxic activity of isolated sesquiterpene lactones from the acetone leaf extract of *Vernonia colorata*”, *S. Afr. J. Bot.*, 75, 176–179.
- [11] Kumari, G. N. K., Masilamani, S., Ganesh, M. R., Aravind, S., Sridhar, S. R., (2003) “Zaluzanin D: a fungistatic sesquiterpene from *Vernonia arborea*”, *Fitoterapia*, 74, 479–482.
- [12] Yao-Haur, K., Yu-Jen, K., Ang-Su, Y., Ming-Der, W., Chi-Wi, O., Li-Ming, Y. K., Jo-Ti, H., Chieh-Fu, C., Shyh-Yuan, L., (2003) “Two novel sesquiterpene lactones, cytotoxic Vernolide-A and -B, from *Vernonia cinerea*”, *Chem. Pharm. Bull.*, 51, 425–426.
- [13] Xin, C., Zha-Jun, Z., Jian-Min, Y., (2006) “Sesquiterpenoids from *Vernonia cinerea*”, *Nat. Pro. Res.*, 20, 31–35.
- [14] Russell, B. W., Andrew, N., Carla, S., Joseph, M., James, S. M., Rabodo, A., Vincent, E. R., David, G. I. K., (2005) “Cytotoxic sesquiterpene lactones from *Vernonia pachyclada* from the Madagascar rainforest”, *J. Nat. Prod.*, 68, 1371–1374.
- [15] Aun, C., Sotheara, H., Christophe, L., Laurence, M., Robert, F., Nadine, A., Riad, E., (2006) “Antimalarial Activity of Sesquiterpene Lactones from *Vernonia cinerea*”, *Chem. Pharm. Bull.*, 54, 1437–1439.
- [16] Pamisha, P., Robert, V., Vinesh, J. M., Peter, J. S., Carmen, A. L., Frank, C., Kelly, C., (2007) “Antiplasmodial hirsutinolides from *Vernonia staehelinoides* and their utilization towards a simplified pharmacophore”, *Phytochemistry*, 68, 1200–1205.
- [17] Rokia, S., Maria, P. G., Nunziatina, D. T., Cosimo, P., Rita, A., (1998) “Vernoniosides and an androstane glycoside from *Vernonia kotschyana*”, *Phytochemistry*, 47, 73–78.

- [18] Alembert, T. T., Apollinaire, T., Pierre, T., Johnson, F. A., Joseph, D. C., Olov, S., (2002) “Vernoguinosterol and vernoguinoside, trypanocidal stigmastane derivatives from *Vernonia guineensis* (Asteraceae)”, *Phytochemistry*, 59, 371–374.
- [19] Giuseppina, C., Rokia, S., Drissa, D., Giovanni, R., Nunziatina De, T., (2004) “New compounds from an extract of *Vernonia colorata* leaves with anti-inflammatory activity”, *J. Nat. Prod.*, 67, 389-394.
- [20] Jing, L., Yunbao, L., Yikang, S., Shishan, Y., Jing, Q., Song, X., Youcui, H., Shuanggang, M., (2009) “New vernocuminosides from the stem barks of *Vernonia cumingiana* Benth”, *Steroids*, 74, 51–61.
- [21] Oketch-Rabah, H. A., Lemmich, E., Dossaji, S. F., Theander, T. G., Olsen, C.E., Cornett, C., Kharazmi, A., Christensen, S. B., (1997) “Two new antiprotozoal 5-methylcoumarins from *Vernonia brachycalyx*”, *J. Nat. Prod.*, 60, 458–461.
- [22] Luis, M-E., Alessandra, B., Cosimo, P., Nunziatina, De T., (2007) “New phenolic derivatives from *Vernonia mapirensis* Gleason”, *ARKIVOC*, 7, 349–358.
- [23] Francois, M., Oliver, P., Andrew, M., Geoffrey, M. M., Kurt, H., (1996) “Flavonol glycosides from *Vernonia galamensis* ssp. *nairobiensis*”, *Phytochemistry*, 43, 283–286.
- [24] Guilian, T., Ubin, Z., Tianyou, Z., Fuquan, Y., Yoichiro, I., (2004) “Separation of flavonoids from the seeds of *Vernonia anthelmintica* Willd by high-speed counter-current chromatography”, *J. Chromatogr.*, 1049, 219–222.
- [25] Inboot, W., Taya, S., Chailungka, A., Meepowpan, P., Wongpoomchai, R., (2012) “Genotoxicity and antigenotoxicity of the methanol extract of *Cleistocalyx nervosum* var. *paniala* seed using a Salmonella mutation assay and rat liver micronucleus tests”, *Mol. Cell. Toxicol.*, 8, 19–24.
- [26] Jansom, C., Bhamarapravati, S., Itharat, A., (2008) “Major anthocyanin from ripe berries of *Cleistocalyx nervosum* var. *paniala*”, *Thammasat Med. J.*, 8, 364–370.

- [27] Maisuthisakul, P., (No date) “Effect of Makiang (*Cleistocalyx nervosum* var. *paniala*) seed on rancidity inhibition in soybean oil model”, *University of the Thai Chamber of Commerce journal*, 282–291.
- [28] Nuengchamnong, N., Ingkaninan, K., (2009) “On-line characterization of phenolic antioxidants in fruit wines from family myrtaceae by liquid chromatography combined with electrospray ionization tandem mass spectrometry and radical scavenging detection”, *LWT-Food. Sci. Technol.*, 42, 297–302.
- [29] http://zipcodezoo.com/plants/c/cleistocalyx_nervosum_var_paniala/ (August 2013)
- [30] <http://dokmaidogma.wordpress.com/2012/07/10/confusing-thai-fruits/> (August 2013)
- [31] Chun-Lin, Y., Yan-H. L., Dong-Z. W., (2004) “Flavonoids from *Cleistocalyx operculatus*”, *Phytochemistry*, 65, 445–447.
- [32] Chun-Lin, Y., Jian-W. L., Dong-Z. W., Yan-H. L., Feng, Q., (2004) “In vitro anti-tumor activity of 2',4'-Dihydroxy-6'-methoxy-3',5'-dimethylchalcone against six established human cancer cell lines”, *Pharm. Res.*, 50, 505–510.
- [33] Chun-Lin, Y., Feng, Q., Dong-Z. W., Yan-H. L., Jian-W. L., (2005) “Induction of apoptosis in K562 human leukemia cells by 2',4'-dihydroxy-6'-methoxy-3',5'-dimethylchalcone”, *Leukemia. Res.*, 29, 887–892.
- [34] Byung-Sun, M., Cao, V. T., Nguyen, T. D., Nguyen, H. D., Han-S. J., Tran, M. H., (2008) “Antioxidative flavonoids from *Cleistocalyx operculatus* buds”, *Chem. Pharm. Bull.*, 56, 1725–1728.
- [35] Nikie'ma1, J. B., Vanhaelen-Fastre', R., Vanhaelen, M., Fontaine, J., De Graef, C., Heenen, M., (2001) “Effects of Antiinflammatory triterpenes isolated from *Leptadenia hastata* Latex on keratinocyte proliferation”, *Phytother. Res.*, 15, 131–134.

- [36] Prachayasittikul, S., Saraban, P., Cherdtrakulkiat, R., Ruchirawat, S., Prachayasittikul, V., (2010) “New bioactive triterpenoids and antimalarial activity of *Diospyros Rubba Lec*”, *Excli. J.*, 9, 1–10.
- [37] Taha, S. E., Shahira, M. E., Ahmed, K. H., Aziza, M.M. A., Gehan, M. K., (2011) “Isolation of biologically active constituents from Moringa (Forssk.) Fiori. (family: Moringaceae) growing in Egypt”, *Pharmacogn Mac.*, 7, 109–115.
- [38] Daniel L, L., Elaine CP, L., Mary Anne M, B., Helenicy NH, V., Aline H, S., Luzia Kalyne AM, L., Amanda A, L., Victor CC, A., Gabriela S, S., Gerly, A. B., Glauce B, V., (2010) “Anti-inflammatory effects and possible mechanism of action of lupeol acetate isolated from *Himatanthus drasticus* (Mart.) Plumel”, *J. Inflamm.*, 7, 1–11.
- [39] Ashalatha, K., Venkateswarlu, Y., Moushumi, P. A., Lalitha, P., Krishnaveni, M., Jayachandran, S., (2010) “Anti inflammatory potential of *Decalepis hamiltonii* (Wight and Arn) as evidenced by down regulation of proinflammatory cytokines-TNF- α and IL-2”, *J. Ethnopharmacol.*, 130, 167–170.
- [40] Guoxin, T., Yoshifumi, I., Dian-Jun, L., Wing Ming, K., (2005) “Eugenol and its structural analogs inhibit monoamine oxidase A and exhibit antidepressant-like activity”, *Bioorgan. Med. Chem.*, 13, 4777–4788.
- [41] Chung, J. Y., Choo, J. H., Lee, M. H., Hwang, J. K., (2006) “Anticariogenic activity of macelignan isolated from *Myristica fragrans* (nutmeg) against *Streptococcus mutans*”, *Phytomedicine*, 13, 261–266.
- [42] Jin, D-Q., Lim, C. S., Hwang, J. K., Han, I., Han, J-S., (2005) “Anti-oxidant and anti-inflammatory activities of macelignan in murine hippocampal cell line and primary culture of rat microglial cells”, *Biochem. Bioph. Res. Co.*, 331, 1264–1269.
- [43] Park, B-Y., Min, B-S., Kwon, O-K., Oh, S-R., Ahn, K-S., Kim, T-J., Kim, D-Y., Bae, K. W., Lee, H-K., (2004) “Increase of caspase-3 activity by lignans from *Machilus thunbergii* in HL-60 cells”, *Biol. Pharm. Bull.*, 27, 1305–1307.

- [44] Cho, Y., Kim, K-H., Shim, J-S., Hwang, J-K., (2008) “Inhibitory effects of macelignan isolated from *Myristica fragrans* HOUTT. on melanin biosynthesis”, *Biol. Pharm. Bull.*, 31, 986–989.
- [45] Ma, J., Hwang, Y. K., Cho, W-H., Han, S-H., Hwang, J. K., Han, J-S., (2009) “Macleignan attenuates activations of mitogen-activated protein kinases and nuclear factor kappa B Induced by lipopolysaccharide in microglial cells”, *Biol. Pharm. Bull.*, 32, 1085–1090.
- [46] Cui, C-A., Jin, D-Q., Hwang, Y. K., Lee, I-S., Hwang, J. K., Ha, I., Han, J-S., (2008) “Macleignan attenuates LPS-induced inflammation and reduces LPS-induced spatial learning impairments in rats”, *Neurosci. Lett.*, 448, 110–114.
- [47] Sohn, J. H., Han, K. L., Choo, J. H., Hwang, J-K., (2007) “Macleignan protects HepG2 cells against *tert*-butylhydroperoxide-induced oxidative damage”, *Biofactors*, 29, 1–10.
- [48] Sohn, J. H., Han, K. L., Kim, J-H., Rukayadi, Y., Hwang, J-K., (2008) “Protective effects of macelignan on cisplatin-induced hepatotoxicity is associated with JNK activation”, *Biol. Pharm. Bull.*, 31, 273–277.
- [49] Han, K. L., Choi, J. S., Lee, J. Y., Song, J., Joe, M. K., Jung, M. H., Hwang, J-K., (2008) “Therapeutic potential of peroxisome proliferators-activated receptor- α/γ dual agonist with alleviation of endoplasmic reticulum stress for the treatment of diabetes”, *Diabetes*, 57, 737–745.
- [50] Gomes, A., Saha, A., Chatterjee, I., Chakravarty, A. K., (2007) “Viper and cobra venom neutralization by β -sitosterol and stigmasterol isolated from the root extract of *Pluchea indica* Less. (Asteraceae)”, *Phytomedicine*, 14, 637–643.
- [51] Nguyen, A. T., Malonne, H., Duez, P., Vanhaelen-Fastre, R., Vanhaelen, M., Fontaine, J., (2004) “Cytotoxic constituents from *Plumbago zeylanica*”, *Fitoterapia*, 75, 500–504.

- [52] Malek, S. N. A., Shin, S. K., Wahab, N. A., Yaacob, H., (2009) “Cytotoxic Components of *Pereskia bleo* (Kunth) DC. (Cactaceae) Leaves”, *Molecules*, 14, 1713–1724.
- [53] Kui-Wu, W., (2007) “A new fatty acid ester of triterpenoid from *Celastrus rosthornianus* with anti-tumor activitives”, *Nat. Prod. Res.*, 21, 669–674.
- [54] Tetsuo, I., Yuuichi, O., Chikako, U., Tsunao, H., (1982) “On the constituents of the leaves of *Viburnum urceolatum*”, *Rep. Fac. Sci.*, 15, 63–68.
- [55] Appleton, R. A., Enzell, C. R., (1971) “Triterpenoids and aromatic components of deertongue leaf”, *Phytochemistry*, 10, 447–449.
- [56] Jamal, A. K., Yaacob, W. A., Din, L. B., (2008) “A chemical study on *Phyllanthus reticulates*”, *J. Phys. Sci.*, 19, 45–50.
- [57] Dewick, P. M. (2002). Medicinal natural products, *The mevalonate and deoxyxylulose phosphate pathways: terpenoids and steroids* (pp.173, 192, 213–219), UK: John Wiley & Sons, Ltd.
- [58] Rahimi, A. A., Ashnagar, A., Nikoei, H., (2012) “Isolation and characterization of 4-allyl-2-methoxphenol (eugenol) from clove buds marketed in Tehran city of Iran”, *Int. J. ChemTech Res.*, 4, 105–108.
- [59] Martin, S. (2004). Understanding mass spectra, *A basic approach: second edition* (pp.314), Canada: John Wiley&Sons, Inc.
- [60] Garg, A., Gupta, B., Prakash, R., Sinngh, S., (2010) “Preparation and characterization of hydroxypropyl- β -cyclodextrin inclusion complex of eugenol: differential pulse voltammetry and $^1\text{H-NMR}$ ”, *Chem. Pharm. Bull.*, 58, 1313–1319.
- [61] Locci, E., Lai, S., Piras, A., Maronglu, B., Lai, A., (2004) “ ^{13}C -CPMAS and ^1H -NMR study of the inclusion complexes of β -cyclodextrin with carvacrol, thymol, and eugenol prepared in supercritical carbon dioxide”, *Chem. Biodivers.*, 1, 1354–1366.

- [62] Gang, D. R., Wang, J., Dudareva, N., Nam, K. H., Simon, J. E., Lewinsohn, E., Pichersky, E., (2001) “An investigation of the storage and biosynthesis of phenylpropenes in Sweet Basil”, *Plant. Physiol.*, 125, 539–555.
- [63] Louie, G. V., Baiga, T. J., Bwman, M. E., Koeduka, T., Taylor, J. H., Spassova, S. M., Pichersky, E., Noel, J. P., (2007) “Structure and rection mechanism of Basil eugenol synthase”, *Plos. One.*, 10, 1–2.
- [64] Woo, S. W., Shin, K. H., Wagner, H., Lotter, H., (1987) “The structure of macelignan from *Myristica fragrans*”, *Phytochemistry*, 26, 1542–1543.
- [65] Filleur, F., Pouget, C., Allais, D. P., Kaouadji, M., Chulia, A. J., (2001) “Lignans and neolignans from *Myristica argentea* warb.”, *Nat. Prod. Lett.*, 16, 1–7.
- [66] Dewick, P. M. (2002). Medicinal natural products, *The shikimate pathway: aromatic amino acids and phenylpropanoids* (pp.122, 131–138), UK: John Wiley & Sons, Ltd.
- [67] Pated, U. U., Haruna, A. K., Garba, M., Iliya, I., Sule, I. M., Abubakar, M. S., Ambi, A. A., (2009) “Isolation of stigmasterol, β -sitosterol and 2-hydroxyhexadecanoic acid methyl ester from the rhizomes of *Stylochiton lancifolius* pyer and kotchy (araceae)”, *Nig. Journ. Pharm. Sci.*, 8, 19–25.
- [68] Charles, J. P., Jacqlynn, B. (1993), The Aldrich Library of ^{13}C and ^1H NMR Spectra, *Volume III* (pp.569), USA: Aldrich Chemical Company, Inc.
- [69] Jamal, A. K., Yaacob, W. A., Din, L. B., (2009) “A chemical study on *Phyllanthus columnaris*”, *Eur. J. Sci. Res.*, 28, 76–81.
- [70] Kolak, U., Topcu, G., Birteksoz, S., Otuk, G., Ulubelen, A., (2005) “Terpenoids and steroids from the roots of *Salvia blepharochlaena*”, *Turk. J. Chem.*, 29, 177–186.
- [71] Bu’Lock, J. D. (1965), The biosynthesis of natural products, *Chapter 4 Isoprenoids* (pp. 60–65), London: McGraw-Hill publishing company, Ltd.

- [72] Inboot, W. "Effect of *Cleistocalyx nervosum* var. *paniala* seeds extracts on cancer chemoprevention". M.S. Thesis. Chiang Mai University, 2012.
- [73] Amor, E. C., Villasenor, I. M., Nawaz, S. A., Hussain, M. S., Choudhary, M. I., (2005) "A dihydrochalcone from *Syzygium samarangense* with anticholinesterase activity", *Philipp. J. Sci.*, 134, 105–111.
- [74] Rachana, S., Arun, K. S., Dinesh, K. K., (1995) "Triterpenoids and chalcone from *Syzygium samarangense*", *Phytochemistry*, 38, 687–689.
- [75] Facundo, V. A., Braz-Filho, R., (2004) "C-methylated flavonoids from the roots of *Piper carniconnectivum* C. DC. (Piperaceae)", *Biochem. Syst. Ecol.*, 32, 1215–1217.
- [76] Amor, E.C., Villasenor, I.M., Ghayur, M.N., Gilani, A.H., Choudhary, M.I., (2005) "Spasmolytic flavonoids from *Syzygium samarangense* (Blume) Merr. & L.M. Perry", *Z. Naturforsch.*, 60, 67–71.
- [77] Nguyen, N.T., Pham, V.C., Litaudon, M., Gueritte, F., Bodo, B., Nguyen, V.T., Nguyen, V.H., (2009) "Novel cyclopeptide and unique flavones from *Desmos rostrata*. Total synthesis of desmorostratone", *Tetrahedron*, 65, 7171–7176.
- [78] Fang, J., Paetz, C., Schneider, B., (2011) "C-methylated flavanones and dihydrochalcones from *Myrica gale* seeds", *Biochem. Syst. Ecol.*, 39, 68–70.
- [79] Nakanishi, K., Goto, T., Ito, S., Natori, S., Nozoe, S. (1975), Natural products chemistry Vol.2, Chapter 9 *Carboaromatic ana related compounds* (pp. 218), Tokyo: Kodansha, Ltd.
- [80] Dacunha-Marinho, B., Martinez, A., Estevez, R.J., (2008) "7-Hydroxy-5-methoxy-6,8-dimethylflavanone: a natural flavonoid", *Acta. Cryst.*, 64, 353–356.
- [81] Kang, Y., Mei, Y., Du, Y., Jin, Z., (2003) "Total synthesis of the highly potent anti-HIV natural product daurichromenic acid along with its two chromane derivatives, rhododaurichromanic acids A and B", *Org. Lett.*, 5, 4481–4484.

- [82] Juanjuan, W., Zhiqiang, C., Shunxi, W., (2010) “Malloapeltic acid, a new benzopyran derivative from *Mallotus apelta*”, *Chen. Nat. Compd.*, 46, 7–9.



ลิขสิทธิ์มหาวิทยาลัยเชียงใหม่
Copyright© by Chiang Mai University
All rights reserved