

REFERENCES

- Aguirre J, Hansberg W, Navarro R. “Fungal responses to reactive oxygen species”. *Med Mycol.* 2006; 44: S101-S107.
- Alonso-Monge R, Navarro-García F, Molero G, Diez-Orejas R, Gustin M, Pla J, *et al.* “Role of the mitogen-activated protein kinase Hog1p in morphogenesis and virulence of *Candida albicans*”. *J Bacteriol.* 1999; 181: 3058-3068.
- Alonso-Monge R, Navarro-García F, Román E, Negredo AI, Eisman B, Nombela C, *et al.* “The Hog1 mitogen-activated protein kinase is essential in the oxidative stress response and chlamydospore formation in *Candida albicans*”. *Eukaryot Cell.* 2003; 2: 351-361.
- Arana DM, Alonso-Monge R, Román E, Nombela C, Pla J. MAP kinase pathways in pathogenic fungi: elements, roles and interactions. In: San-Blas G, Calderone RA, editors. *Pathogenic fungi: insights in molecular biology.* 2008; pp.151-170.
- Bahn YS, Kojima K, Cox GM, Heitman J. “Specialization of the HOG pathway and its impact on differentiation and virulence of *Cryptococcus neoformans*”. *Mol Biol Cell.* 2005; 16: 2285-2300.
- Balázs A, Pócsi I, Hamari Z, Leiter É, Emri T, Miskei M, *et al.* “AtfA bZIP-type transcription factor regulates oxidative and osmotic stress responses in *Aspergillus fumigatus*”. *Mol Genet Genomics.* 2010; 283: 289-303.
- Bermejo C, Rodríguez E, García R, Rodríguez-Peña J., Rodríguez de la Concepción ML, Rivas C, *et al.* “The sequential activation of the yeast HOG and SLT2 pathway is required for cell survival to cell wall stress”. *Mol Biol Cell.* 2008; 19: 1113-1124.
- Billmyre RB, Calo S, Feretzaki M, Wang X, Heitman J. “RNAi function, diversity, and loss in the fungal kingdom”. *Chromosome Res.* 2013; 21: 561-572.

- Borneman AR, Hynes MJ, Andrianopoulos A. "An *STE12* homolog from the asexual, dimorphic fungus *Penicillium marneffe* complements the defect in sexual development of an *Aspergillus nidulans steA* mutant". *Genetics*. 2001; 157: 1003-1014.
- Boyce KJ, Andrianopoulos A. "Ste20-related kinases: effectors of signaling and morphogenesis in fungi". *Trends Microbiol*. 2011; 19:400-410.
- Boyce KJ, Andrianopoulos A. "Morphogenetic circuit regulating growth and development in the dimorphic pathogen *Penicillium marneffe*". *Eukaryot Cell*. 2013; 12: 154-160.
- Boyce KJ, Schreider L, Andrianopoulos A. "In vivo yeast cell morphogenesis is regulated by a p21-activated kinase in the human pathogen *Penicillium marneffe*". *PLoS Pathog*. 2009; 5: e1000678.
- Boyce KJ, Schreider L, Kirszenblat L, Andrianopoulos A. "The two-component histidine kinases DrkA and SlnA are required for in vivo growth in the human pathogen *Penicillium marneffe*". *Mol Microbiol*. 2011; 82: 1164-1184.
- Brakhage AA, Langfelder K. "Menacing mold: the molecular biology of *Aspergillus fumigatus*". *Annu Rev Microbiol*. 2002; 56: 433-455.
- Brown SM, Campbell LT, Lodge JK. "*Cryptococcus neoformans*, a fungus under stress". *Curr Opin Microbiol*. 2007; 10: 320-325.
- Brown AJ, Haynes K, Quinn J. "Nitrosative and oxidative stress responses in fungal pathogenicity". *Curr Opin Microbiol*. 2009; 12: 384-391.
- Bugeja HE, Boyce KJ, Weerasinghe H, Beard S, Jeziorowski A, Pasricha S., et al. "Tools for high efficiency genetic manipulation of the human pathogen *Penicillium marneffe*". *Fungal Genet Biol*. 2012; 10: 772-778.
- Campos EG, Jesuino RS, Dantas Ada S, Brígido Mde M, Felipe MS. "Oxidative stress response in *Paracoccidioides brasiliensis*". *GMR*. 2005; 4: 409-429.

- Cánovas D, Andrianopoulos A. The biology of the thermally dimorphic fungal pathogen *Penicillium marneffe*. In: Kavanage K, editor. *New insights in Medical Mycology*, Springer-Verlag Press, Berlin Heidelberg. 2007; pp. 213-226.
- Cao C, Liu W, Li R. “*Penicillium marneffe* Skn7, a novel gene, could complement the hypersensitivity of *S. cerevisiae* *skn7* disruptant strain to oxidative stress”. *Mycopathologia*. 2009; 168: 23-30.
- Cardoza RE, Vizcaino JA, Hermosa MR, Monte E, Gutiérrez S. “A comparison of the phenotypic and genetic stability of recombinant *Trichoderma* spp. Generated by protoplast- and *Agrobacterium*-mediated transformation”. *J Microbiol*. 2006; 44: 383-395.
- Catlett NL, Lee B, Yoder OC, Turgeon BG. “Split-marker recombination for efficient targeted deletion of fungal genes”. *Fungal Genet Newsl*. 2003; 50: 9-11.
- Chauhan N, Calderone R. “Two-component signal transduction proteins as potential drug targets in medically important fungi”. *Infect Immun*. 2008; 76: 4795-4803.
- Chauhan N, Latge JP, Calderone R. “Signalling and oxidant adaptation in *Candida albicans* and *Aspergillus fumigatus*”. *Nat Rev Microbiol*. 2006; 4: 435-44.
- Cooper CR Jr, Vanittanakom N. “Insights into the pathogenicity of *Penicillium marneffe*”. *Future Microbiol*. 2008; 3: 43-55.
- Costa V, Moradas-Ferreira P. “Oxidative stress and signal transduction in *Saccharomyces cerevisiae*: insights into ageing apoptosis and diseases”. *Mol Aspects Med*. 2001; 22: 217-246.
- Du C, Sarfati J, Latge JP, Calderone R. “The role of the *sakA (Hog1)* and *tcsB (sln1)* genes in the oxidants adaptation of *Aspergillus fumigatus*”. *Med Mycol*. 2006; 44: 211-218.
- Eaton CJ, Jourdain I, Foster SJ, Hyams J, Scott B. “Functional analysis of a fungal endophyte stress-activated MAP kinase”. *Curr Genet*. 2008; 53: 163-174.

- Enjalbert B, Smith DA, Cornell MJ, Alam I, Nicholls S, Brown AJ, *et al.* “Role of the Hog1 stress-activated protein kinase in the global transcriptional response to stress in the fungal pathogen *Candida albicans*”. *Mol Biol Cell*. 2006; 17: 1018-1032.
- Flattery-O’Brien J, Collinson LP, Dawes IW. “*Saccharomyces cerevisiae* has an inducible response to menadione which differs from that to hydrogen peroxide”. *J Gen Microbiol*. 1993; 139: 501-507.
- Furukawa K, Hoshi Y, Maeda T, Nakajima T, Abe K. “*Aspergillus nidulans* HOG pathway is activated only by two-component signaling pathway in response to osmotic stress”. *Mol Microbiol*. 2005; 56: 1246-1261.
- Gessler NN, Aver’yanov AA, Belozerskaya TA. “Reactive oxygen species in regulation of fungal development”. *Biochemistry (Mosc)*. 2007; 72: 1091-1109.
- Gravelat FN, Ejzykowicz DE, Chiang LY, Chabot JC, Urb M, Macdonald KD, *et al.* “*Aspergillus fumigatus* MedA governs adherence, host cell interactions and virulence”. *Cell Microbiol*. 2010; 12: 473-488.
- Hagiwara D, Asano Y, Yamashino T, Mizuno T. “Characterization of bZip-type transcription factor AtfA with reference to stress responses of conidia of *Aspergillus nidulans*”. *Biosci Biotechnol Biochem*. 2008; 72: 2756-2760.
- Hamilton AJ, Holdom MD. “Antioxidant systems in the pathogenic fungi of man and their role in virulence”. *Med Mycol*. 1999; 37: 375-389.
- Hamilton AJ, Jeavons L, Youngchim S, Vanittanakom N, Hay RJ. “Sialic acid-dependent recognition of laminin by *Penicillium marneffei* conidia”. *Infect Immun*. 1998; 66: 6024-6026.
- Herrero E, Ros J, Bellí G, Cabisco E. “Redox control and oxidative stress in yeast cells”. *Biochim Biophys Acta*. 2008; 1780: 1217-1235.
- Hohmann S. “Osmotic stress signaling and osmoadaptation in yeasts”. *Microbiol Mol Biol Rev*. 2002; 66: 300-372.

- Ikner A, Shiozaki K. "Yeast signaling pathways in the oxidative stress response". *Mutat Res.* 2005; 569: 13-27.
- Jin FJ, Maruyama J, Juvvadi PR, Arioka M, Kitamoto K. "Adenine auxotrophic mutant of *Aspergillus oryzae*: Development of a novel transformation system with triple auxotrophic hosts". *Biosci Biotechnol Biochem.* 2004; 68: 656-662.
- Kammasook A, Pongpom P, Vanittanakom N. "Cloning, characterization and differential expression of and *hsp70* gene from the pathogenic dimorphic fungus, *Penicillium marneffe*". *DNA Sequence.* 2007; 18: 385-394.
- Kawasaki L, Sánchez O, Shiozaki K, Aguirre J. "SakA MAP kinase is involved stress signal transduction, sexual development and spore viability in *Aspergillus nidulans*". *Mol Microbiol.* 2002; 45: 1153-1163.
- Krappmann S. "Gene targeting in filamentous fungi: the benefits of impaired repair". *Fungal Biol Rev.* 2007; 21: 25-29.
- Kruppa M, Goins T, Cutler JE, Lowman D, Williams D, Chauhan N, *et al.* "The role of the *Candida albicans* histidine kinase (*CHK1*) gene in the regulation of cell wall mannan and glucan biosynthesis". *FEMS Yeast Res.* 2003; 3: 289-299.
- Kück U, Hoff B. "New tools for the genetic manipulation of filamentous fungi". *Appl Microbiol Biotechnol.* 2010; 86: 51-62.
- Kuge S, Jones N. "YAP1 dependent activation of TRX2 is essential for the response of *Saccharomyces cerevisiae* to oxidative stress by hydroperoxide". *EMBO J.* 1994; 13: 655-664.
- Kumar S, McLaughlin MM, McDonnell PC, Lee JC, Livi GP, Young PR. "Human mitogen-activated protein kinase CSBP1, but not CSBP2, complements a *hog1* deletion in yeast". *J Biol Chem.* 1995; 270: 29043-29046.
- Lara-Rojas, F, Sánchez O, Kawasaki L, Aguirre J. "*Aspergillus nidulans* transcription factor AtfA interacts with the MAPK SakA to regulate general responses, development and spore functions". *Mol Microbiol.* 2010; 80: 436-454.

- Lau SK, Tse H, Chan JS, Zhou AC, Curreem SO, Lau CC, *et al.* “Proteome profiling of the dimorphic fungus *Penicillium marneffei* extracellular proteins and identification of glyceraldehyde-3-phosphate dehydrogenase as an important adhesion factor for conidial attachment”. *FEBS J.* 2013; 280: 6613-6626.
- Li D, Gurkovska V, Sheridan M, Calderone R, Chauhan N. “Studies on the regulation of the two-component histidine kinase gene *CHK1* in *Candida albicans* using the heterologous *lacZ* reporter gene”. *Microbiology.* 2004; 150: 3305-3313.
- Lin X, Ran Y, Gou L, He F, Zhang R, Wang P, *et al.* “Comprehensive transcription analysis of human pathogenic fungus *Penicillium marneffei* in mycelial and yeast cells”. *Med Mycol.* 2012; 50: 835-842.
- Ma D, Li R. “Current understanding of HOG-MAPK pathway in *Aspergillus fumigatus*”. *Mycopathologia.* 2013; 175: 13-23.
- Marr KA, Koudadoust M, Black M, Balajee SA. “Early events in macrophage killing of *Aspergillus fumigatus* conidia: New flow cytometric viability assay”. *Clin Diagn Lab Immunol.* 2001; 8: 1240-1247.
- Miskei M, Karányi Z, Pócsi I. “Annotation of stress-response proteins in the aspergilli”. *Fungal Genet Biol.* 2009; 46: S105-S120.
- Morano KA, Grant CM, Moye-Rowley WS. “The response to heat shock and oxidative stress in *Saccharomyces cerevisiae*”. *Genetics.* 2012; 190: 1157-1195.
- Morgan BA, Ranks GR, Toone WM, Raitt D, Kuge S, Johnston LH. “The Skn7 response regulator controls gene expression in the oxidative stress response of the budding yeast *Saccharomyces cerevisiae*”. *Embo J.* 1997; 16: 1035-1044.
- Moye-Rowley WS. “Regulation of the Transcriptional Response to Oxidative stress in Fungi: similarities and differences”. *Eukaryot Cell.* 2003; 2: 381-389.
- Nelson KE, Supparatpinyo K, Vanittanakom N. Penicilliosis. In: Kauffman CA, Pappas PG, Sobel JD, Dismukes WE, editors. *Essentials of Clinical Mycology* (2nd Ed), Springer Science+Business Media, LLC. 2011; pp. 399-411.

- Papadakis MA, Workman CT. "Oxidative stress response pathways: Fission yeast as archetype". *Crit Rev Microbiol*. 2014 Feb 27; Epub ahead of print.
- Pongpom M, Sirisanthana T, Vanittanakom N. "Application of nested PCR to detect *Penicillium marneffe* in serum sample". *Med Mycol*. 2009; 47: 549-553.
- Pongpom M, Sawatdeechaikul P, Kummasook A, Khanthawong S, Vanittanakom N. "Antioxidative and immunogenic properties of catalase-peroxidase protein in *Penicillium marneffe*". *Med Mycol*. 2013; 51: 835-842.
- Punt PJ, Oliver RP, Dingemans MA, Pouwels PH, Van den Hondel CA. "Transformation of *Aspergillus* based on the hygromycin B resistance marker from *Escherichia coli*". *Genetics*. 1987; 56: 117-124.
- Rispail N, Soanes DM, Ant C, Czajkowski R, Grünler A, Huguet R *et al*. "Comparative genomics of MAP kinase and calcium-calmodulin signaling components in plant and human pathogenic fungi". *Fungal Genet Biol*. 2009; 46: 287-298.
- Román E, Arana DM, Nombela C, Alonso-Monge R, Pla J. "MAP kinase pathways as regulators of fungal virulence". *Trends Microbiol*. 2007; 15: 181-190.
- Romani L. "Immunity to fungal infections". *Nat Rev Immunol*. 2011; 11: 275-288.
- Rongrungruang Y, Levitz SM. "Interactions of *Penicillium marneffe* with human leucocytes *in vitro*". *Infect Immun*. 1999; 67: 4732-4736.
- Sakamoto K, Iwashita K, Yamada O, Kobayashi K, Mizuno A, Akita O, *et al*. "*Aspergillus oryzae* controls conidial germination and stress tolerance". *Fungal Genet Biol*. 2009; 46: 887-897.
- Samson RA, Yilmaz N, Houbraken J, Spierenburg H, Seifert DA, Peterson SW, *et al*. "Phylogeny and nomenclature of the genus *Talaromyces* and taxa accommodated in *Penicillium* subgenus *Biverticillium*". *Stud Mycol*. 2011; 70: 159-183.
- Sanger F, Nicklen S, Coulson AR. "DNA sequencing with chain-terminating inhibitors". *Proc Natl Acad Sci USA*. 1977; 74: 5463-5467.

- Segmüller N, Ellendorf U, Tudzynski B, Tudzynski P. “BcSAK1, a stress-activated mitogen-activated protein kinase, is involved in vegetative differentiation and pathogenicity in *Botrytis cinerea*”. *Eukaryot Cell*. 2007; 6: 211-221.
- Seider K, Heyken A, Lüttich A, Miramón P, Hube B. “Interaction of pathogenic yeasts with phagocytes: survival, persistence and escape”. *Curr Opin Microbiol*. 2010; 13: 1-9.
- Sheppard DC, Doedt T, Chiang LY, Kim HS, Chen D, Nierman WC, *et al*. “The *Aspergillus fumigatus* StuA protein governs the up-regulation of a discrete transcriptional program during the acquisition of developmental competence”. *Mol Biol Cell*. 2005; 16: 5866-5879.
- Smith DA, Morgan BA, Quinn J. “Stress signalling to fungal stress-activated protein kinase pathways”. *FEMS*. 2010; 306: 1-8.
- Tanoue T, Nishida E. “Molecular recognitions in the MAP kinase cascades”. *Cell Signal*. 2003; 15: 455-462.
- Thirach S, Cooper CR Jr, Vanittanakom P, Vanittanakom N. “The copper, zinc superoxide dismutase gene of *Penicillium marneffe*: cloning, characterization, and differential expression during phase transition and macrophage infection”. *Med Mycol*. 2007; 45: 409-417.
- Tomazett PK, Castro Nda S, Lenzi HL, de Almeida Soares CM, Pereira M. “Response of *Paracoccidioides brasiliensis* Pb01 to stressor agents and cell wall osmoregulators”. *Fungal Biol*. 2011; 115: 62-69.
- Tongchusak S, Pongtanalert P, Pongsunk S, Chawengkirttikul R, Chaiyaroj SC. “Distinct immunologic properties of *Penicillium marneffe* yeasts obtained from different in vitro growth conditions”. *Asian Pac J Allergy Immunol*. 2004; 22: 229-235.
- Vanittanakom N, Vanittanakom P, Hay RJ. “Rapid identification of *Penicillium marneffe* by PCR-based detection of specific sequences on the rRNA gene”. *J Clin Microbiol*. 2002; 40: 1739-1742.

- Vanittanakom N, Cooper CR Jr, Fisher MC, Sirisanthana T. “*Penicillium marneffe* infection and recent advances in the epidemiology and molecular biology aspects”. *Clin Microbiol Rev.* 2006; 19: 95-110.
- Vanittanakom N, Pongpom M, Praparattanapan J, Cooper CR Jr, Sirisanthana T. “Isolation and expression of heat shock protein 30 from *Penicillium marneffe*”. *Med Mycol.* 2009; 47: 521-526.
- Vivancos AP, Jara M, Zuin A, Sansó M, Hidalgo E. “Oxidative stress in *Schizosaccharomyces pombe*: different H₂O₂ levels, different response pathways”. *Mol Genet Genomics.* 2006; 276: 495-502.
- Viviani MA, Vanittanakom N. Penicilliosis. In: Merz WG, Hay RJ, editors. Medical Mycology. Topley & Wilson’s Microbiology and Microbial Infections, Cox F, Wakelin D, Murray P, Borriello P, editors (10th Ed). Hodder Arnold, London. 2005; pp: 560-575.
- Wilkinson MG, Samuels M, Takeda T, Toone WM, Shieh JC, Toda T, *et al.* “The Atf1 transcription factor is a target for the Sty1 stress-activated MAP kinase pathway in fission yeast”. *Genes Dev.* 1996; 10: 2289-2301.
- Woo PC, Zhen H, Cai JJ, Yu J, Lau SK, Wang J, *et al.* “The mitochondrial genome of the thermal dimorphic fungus *Penicillium marneffe* is more closely related to those of molds than yeasts”. *FEBS Let.* 2003; 555: 469-477.
- Woo PC, Lam CW, Tam EW, Leung CK, Wong SS, Lau SK, *et al.* “First discovery of two polyketide synthase genes for mitorubrinic acid and mitorubrinol yellow pigment biosynthesis and implication in virulence of *Penicillium marneffe*”. *PLoS Negl Trop Dis.* 2012; 6: e1871.
- Xu JR. “MAP kinases in fungal pathogens”. *Fungal Genet Biol.* 2000; 31: 137-152.
- Yang E, Wang G, Woo PC, Lau SK, Chow WN, Chong KT, *et al.* “Unraveling the molecular basis of temperature-dependent genetic regulation in *P. marneffe*”. *Eukaryot Cell.* 2013; 12: 1214-1224.

- Youngchim S, Vanittanakom N, Hamilton AJ. "Analysis of the enzymatic activities of mycelial and yeast phases of *Penicillium marneffe*". *Med Mycol.* 1999; 37: 455-450.
- Yuen KY, Pascal G, Wong SS, Glaser P, Woo PC, Kunst F, *et al.* "Exploring the *Penicillium marneffe* genome". *Arch Microbiol.* 2003; 179: 339-353.
- Zhang Y, Zhao J, Fang W, Zhang J, Luo Z, Zhang M, *et al.* "Mitogen-activated protein kinase *hog1* in the entomopathogenic *Beauveria bassiana* regulates environmental stress responses and virulence to insects". *Appl Environ Microbiol.* 2009; 75: 3787-3795.
- Zhao X, Mehrabi R, Xu JR. "Mitogen-activated protein kinase pathways and fungal pathogenesis". *Eukaryot Cell.* 2007; 6: 1701-1714.
- Zi Z, Liebermeister W, Klipp E. "A quantitative study of the Hog1 MAPK response to fluctuating osmotic stress in *Saccharomyces cerevisiae*". *PLoS ONE.* 2010; 5: e9522.