

REFERENCES

- Ahmad, B., Khalid, B., Zani, A., Hussain, N., & Quek, K. (2011). Influencing factors of glycemic control in patients attending different types of urban health care settings in Malaysia. *Diabetes Clinical Practice*, 93(1), e12-e14. doi:10.1016/j.diabres.2011.02.020.
- Akbar, D. H. (2001). Low rates of diabetes patients reaching good control targets. *Eastern Mediterranean Health Journal*, 7(4-5), 671-678.
- Alahakoon, A. (2012). *Tea and prepared food should sell with separate sugar and salt*. Divaina e-paper, page 3. Retrieved from <http://epaper.divaina.com/>
- Albarran, N. B., Ballesteros, M. N., Morales, G. G., & Ortega, M. I. (2006). Dietary behavior and type 2 diabetes care. *Patient Education and Counseling*, 61(2), 191-199.
- Ali, M. K., Bullard, K. M., Imperatore, G., Barker, L., & Gregg, E. W. (2012). Characteristics associated with poor glycemic control among adults with self-reported diagnoses diabetes--National Health and Nutrition Examination Survey, United States, 2007-2010. *Morbidity and Mortality Weekly Report*, 61(2), 32-37.
- Aljasem, L. I., Peyrot, M., Wissow, L., & Rubin, R.R. (2001). The impact of barriers and self-efficacy on self-care behaviors in type 2 diabetes. *Diabetes Educator*, 27(3), 393-404.

- Allender, S., Lacey, B., Webster, P., Rayner, M., Deepa, M., Scarborough, P., . . . Mohan, V. (2010). Level of urbanization and noncommunicable disease risk factors in Tamil Nadu, India. *Bulletin of the World Health Organization*, 88, 297-304. doi:10.2471/BLT.09.065847
- Al-Sultan, F. A., & Al-Zanki, N. (2005). Clinical epidemiology of Type 2 diabetes mellitus in Kuwait. *Kuwait Medical Journal*, 37(2), 98-104.
- American Association of Diabetic Educators. (2012). Position statement: Culturally sensitivity and diabetes education. *The Diabetic Educator*, 38, 137-140.
- American Diabetes Association. (2008). Nutrition recommendations and interventions for diabetes: A position statement of the American Diabetes Association. *Diabetes Care*, 31(1), S61-S78. doi:10.2337/dc08-S061
- American Diabetes Association. (2012). Standards of medical care in diabetes-2012. *Diabetes Care*, 35(Suppl. 1), S11-S63. doi:10.2337/dc12-s011
- American Diabetes Association. (2013). Standards of medical care in diabetes-2013. *Diabetes Care*, 36(Suppl. 1), S11- S66. doi:10.2337/dc13-S011
- Arembepola, C., Allender, S., Ekanayake, R., & Fernando, D. (2008). Urban living and obesity: is it independent of its population and lifestyle characteristics? *Tropical Medicine and International Health*, 13(4), 448-457. doi:10.1111/j.1365-3156.2008.02021.x
- Awareness campaign for Diabetes Day-2011*. (2011). Retrieved from <http://www.youtube.com/watch?v=JMC1umXropU>
- Ayele, K., Tesfa, B., Abebe, L., Tilahun, T., & Girma, E. (2012). Self care behavior among patients with diabetes in Harari, Eastern Ethiopia: The health belief model perspective. *PLoS ONE*, 7(4), e35515. doi:10.1371/journal.pone.0035515

- Beverly, E. A., Hultgren, B. A., Brooks, K. M., Ritholz, M. D., Abrahamson, M. J., & Weinger, K. (2011). Understanding physicians' challenges when treating type 2 diabetic patients' social and emotional difficulties: A qualitative study. *Diabetes Care*, *34*(5), 1086-1088. doi:10.2337/dc10-2298
- Block, D. J. (2006). *Healthcare outcomes management: Strategies for planning and evaluation*. London: Jones and Bartlett.
- Bowser, D., Utz, S., Glick, D., & Harmon, R. (2010). A systematic review of the relationship of diabetes mellitus, depression, and missed appointments in a low-income uninsured population. *Archives of Psychiatric Nursing*, *24*(5), 317-329. doi:10.1016/j.apnu.2009.12.004
- Brand-Miller, J., Hayne, S. Petocz, P., & Colagiuri, S. (2003). Low-glycemic index diets in the management of diabetes: A meta-analysis of randomized controlled trials. *Diabetes Care*, *26*(8), 2261-2267. doi:10.2337/diacare.26.8.2261
- Brem, H., & Tomic-Canic, M. (2007). Cellular and molecular basis of wound healing in diabetes. *The Journal of Clinical Investigation*, *117*(5), 1219-1222. doi:10.1172/JCI32169
- Broom, A., Wijewardena, K., Sibbritt, D., Adams, J., & Nayar, K. R. (2010). The use of traditional, complementary and alternative medicine in Sri Lankan cancer care: Results from a survey of 500 cancer patients. *Public Health*, *124*(4), 232-237. doi:10.1016/j.puhe.2010.02.012
- Brown, K., Avis, M., & Hubbard, M. (2007). Health beliefs of African-Caribbean people with type 2 diabetes: A qualitative study. *British Journal of General Practice*, *57*(539), 461-469.

Burns, N., & Grove, S. K. (2009). *The practice of nursing research: Conduct, critique and utilization* (6th ed.). Philadelphia: W.B. Saunders.

Caban, A., & Walker, E. A. (2006). A systematic review of research on culturally relevant issues for hispanics with diabetes. *The Diabetes Educator*, 32(4), 584-595. doi:10.1177/0145721706290435

Caldwell, J., Gajanayaka, I., Caldwell, P., & Pieris, I. (1989). Sensitization to illness and the risk of death: An exploration for Sri Lanka's good health for all. *Social Science & Medicine*, 28(4), 365-379.

Campos, C. (2007). Addressing cultural barriers to the successful use of insulin in Hispanics with type 2 diabetes: Review article. *Southern Medical Association*, 100(8), 812-820.

Carbone, E. T., Rosal, M. C., Torres, M. I., Goins, K. V., & Bermudez, O. I. (2007). Diabetes self-management: Perspectives of Latino patients and their health care providers. *Patient Education & Counseling*, 66(2), 202-210.

Casarez, R. L., Engebretson, J., & Ostwald, S. (2010). Spiritual practices in self-management of diabetes in African Americans. *Holistic Nursing Practice*, 24(4), 227-237. doi:10.1097/HNP.0b013e3181e903c6

Cavanaugh, K., Huizinga, M. M., Wallston, K. A., Gebretsadik, T., Shintani, A., Davis, D., . . . Rothman, R. L. (2008). Association of numeracy and diabetes control. *Annals of Internal Medicine*, 148(10), 737-746.

doi:10.7326/0003-4819-148-10-200805200-00006

- Cha, E., Yang, K., Lee, J., Min, J., Kim, K. H., Dunbar, S. B., & Jennings, B. M. (2012). Understanding cultural issues in the diabetes self-management behaviors of Korean immigrants. *Diabetes Educator*, 38(6), 835-844. doi:10.1177/0145721712460283.
- Chacko, E. (2003). Culture and therapy: Complementary strategies for the treatment of type 2 diabetes in an urban setting in Kerala, India. *Social Science & Medicine*, 56(5), 1087-1098.
- Chang, H. A., Wallis, M., Tiralongo, E., & Wang, H. L. (2012). Decision-making related to complementary and alternative medicine use by people with Type 2 diabetes: a qualitative study. *Journal of Clinical Nursing*, 21(21-22), 3205-3215. doi:10.1111/j.1365-2702.2012.04339.x
- Chaufan, C., Davis, M., & Constantino, S. (2011). The twin epidemics of poverty and diabetes: Understanding diabetes disparities in a low-income Latino and immigrant neighborhood. *Journal of Community Health*, 36(6), 1032-1043. doi:10.1007/s10900-011-9406-2
- Chesla, C., & Chun, K. (2005). Accommodating type 2 diabetes in the Chinese American family. *Qualitative Health Research*, 15(2), 240-255. doi:10.1177/1049732304272050
- Chesla, C., Chun, K., & Kwan, C. M. (2009). Cultural and family challenges to managing type 2 diabetes in immigrant Chinese Americans. *Diabetes Care*, 32(10), 1812-1816. doi:10.2337/dc09-0278

- Chlebowy, D. O., Hood, S., & LaJoie, A. S. (2010). Facilitators and barriers to self-management of type 2 diabetes among urban African American adults: Focus group findings. *The Diabetes Educator*, 36(6), 897-904. doi:10.1177/0145721710385579
- Chowdhury, A. M., Helman, C., & Greenhalgh, T. (2000). Food beliefs and practices among British Bangladeshis with diabetes: Implications for health education. *Anthropology & Medicine*, 7(2), 209-226. doi:10.1080/713650589
- Chun, K. M., & Chesla, C. A. (2004). Cultural issues in disease management for Chinese Americans with type 2 diabetes. *Psychology & Health*, 19(6), 767-785. doi:10.1080/08870440410001722958
- Clark, M., & Hampson, S. E. (2003). Comparison of patients' and healthcare professionals' beliefs about and attitudes towards type 2 diabetes. *Diabetic Medicine*, 20(2), 152-154.
- Clarke, A. (1993). Focus group interview in health care research. *Professional Nurse*, 14, 395-379.
- Colombo South Teaching Hospital. (2010). *Quarterly hospital statistical bulletin-- 2nd quarter*. Sri Lanka: Colombo South Teaching Hospital.
- Creswell, J. W. (2009). *Research design: Quantitative, qualitative and mixed method approaches* (3rd ed). Los Angeles, CA: SAGE.
- Daly, J.M., Hartz, A.J., Xu, Y., Levy, B. T., James, P. A., Merchant, M. L., & Garrett, R. E. (2009). An assessment of attitudes, behaviors, and outcomes of patients with type 2 diabetes. *Journal of the American Board of Family Medicine*, 22(3), 280-290. doi:10.3122/jabfm.2009.03.080114

Danaei, G., Finucane, M. M., Lu, Y., Singh, G. M., Cowen, M. J., Paciorek, C. J., . . .

Ezzati, M. (2011). National, regional, and global trends in fasting plasma glucose and diabetes prevalence since 1980: Systematic analysis of health examination surveys and epidemiological studies with 370 country-years and 2.7 million participants. *Lancet*, 378(9785), 31-40. doi:10.1016/S0140-6736(11)60679-x

Davies, M., Brophy, S., Williams, R., & Taylor, A. (2006). The prevalence, severity and impact of painful diabetic peripheral neuropathy in type 2 diabetes. *Diabetes Care*, 29(7), 1518-1522.

Davila, E. P. (2010). *Glycemic control risk factors, quality of life, workforce participation, and mortality among adults with type 2 diabetes* (Doctoral dissertation, Florida International University). Available from ProQuest ETD Collection for FIU. (Paper No. AAI3470215).

Davila, E. P., Florez, H., Trepka, M. J., Fleming, L. E., Niyonsenga, T., Lee, D. J. & Parkash, J. (2011). Long work hours is associated with suboptimal glycemic control among US workers with diabetes. *American Journal of Industrial Medicine*, 54(4), 375-383. doi:10.1002/ajim.20923

De Silva, A. P., De Silva, S. H. P., Liyanage, I. K., Rajapakse, L. C., Jayasinghe, K. S. A., Katulanda, P., . . . Wijeratne, S. (2012). Social, cultural and economical determinants of diabetes mellitus in Kalutara district, Sri Lanka: A cross sectional descriptive study. *International Journal for Equity in Health*, 11(76), doi:10.1186/1475-9276-11-76

De Silva, K. M. (1981). *A history of Sri Lanka*. Delhi: Oxford University Press.

Denzin, N. K., & Lincoln, Y. S. (1994). *Handbook of qualitative research*. United Kingdom: SAGE.

Department of Census and Statistics. (2007). *Statistics abstract 2007*. Retrieved from [http://www.statistics.gov.lk/Abstract 2006/abstract 2006/index.htm](http://www.statistics.gov.lk/Abstract%202006/abstract%202006/index.htm)

Department of Census and Statistics. (2010a). *Household and income report 2009/10: Final report*. Retrieved from http://www.statistics.gov.lk/PopHouSat/CPH2011/Pages/sm/CPH%202011_R1.pdf

Department of Census and Statistics. (2010b). *Population of Sri Lanka by district*. Retrieved from http://www.statistics.gov.lk/samplesurvey/occ_code.pdf

Department of Census and Statistics. (2012). *Population of Sri Lanka by district*. Retrieved from http://www.statistics.gov.lk/samplesurvey/occ_code.pdf

Dwivedi, G., & Dwivedi, S. (2007). Sushruta-the clinician-teacher par excellence. *Indian Journal of Chest Disease and Allied Science*, 49, 243-244.

Early, K. B., Shultz, J., & Corbett, S. (2009). Assessing diabetes dietary goals and self-management based in-depth interviews with Latino and Caucasian clients with type 2 diabetes. *Journal of Transcultural Nursing*, 20(4), 371-381. doi:10.1177/1043659609334928

Ediriweera, E. R., & Rathnasooriya, W. D. (2009). A review of herbs used in treatment of diabetes mellitus by Sri Lankan ayurvedic and traditional physicians. *Ayurveda*, 30, 373-391.

Ekanayake, S., Welihinda, J., & Jansz, E. R. (2009). *Glycemic indices of Sri Lankan food*. Retrieved from <http://www.nrc.gov.lk/SAG/index.html>

Engelgau, M., Okamoto, K., Navaratne, K. V., & Gopalan, S. (2010). *Prevention and control of selected chronic NCDs in Sri Lanka: Policy options and action.*

HNP Discussion paper, World Bank. Retrieved from http://siteresources.worldbank.org/HEALTH_NUTRITION_AND_POPULATION/Resources/281627-1095698140167/NCDsSriLanka.pdf

Farmer, A., Kinmonth, A. L., & Sutton, S. (2006). Measuring beliefs about taking hypoglycaemic medication among people with Type 2 diabetes. *Diabetic Medicine*, 23(3), 265-270.

Fernando, D., & Dewapura, G. (2000). *Guidelines on management of diabetes mellitus in Sri Lanka.* Sri Lanka: Sri Lanka Medical Association.

Fetterman, D. M. (1998). *Ethnography* (2nd ed.). London: SAGE.

Field, P. A., & Morse, J. M. (1985). *Nursing research: The application of qualitative approaches.* London: Croon Helm.

Filestead, W. J. (1979). Qualitative methods: A needed perspective in evaluation research. In T. D. Cook & C. I. Reichardt (Eds.), *Qualitative and quantitative methods in evaluation research.* Beverly Hills, CA: SAGE.

Fleming, E., Carter, B., & Pettigrew, J. (2008). The influence of culture on diabetes self-management: Perspectives of Gujarati Muslim men who reside in northwest England. *Journal of Clinical Nursing*, 17(5A), 51-59.

doi:10.1111/j.1365-2702.2007.02178.x

Fleming, E., & Gillibrand, W. (2009). An exploration of culture, diabetes and nursing in the South Asian Community: A meta-synthesis of qualitative studies. *Journal of Transcultural Nursing*, 20(2), 146-155.

doi:10.1177/1043659608330058

- Fortmann, A., Gallo, L. C., & Philis-Tsimikas, A. (2011). Glycemic control among Latinos with type 2 diabetes: The role of socio-environmental support resources. *Health Psychology, 30*(3), 251-258. doi:10.1037/a0022850.
- Fox, K. M., Gerber, R. A., Bolinder, B., Chen, J., & Kumara, S. (2006). Prevalence of inadequate glycemic control among patients with type 2 diabetes in the United Kingdom general practice research data base: A series of retrospective analysis of data from 1998 through 2002. *Clinical Therapeutics, 28*(3), 388-395.
- Franz, M. J., Bantle, J. P., Beebe, C. A., Brunzell, J. D., Chiasson, J. L., Garg, A., . . . Wheeler, M. (2004). Nutrition principles and recommendations in diabetes. *Diabetes Care, 27*(Suppl. 1), s36-s46.
- Frykberg, R. G., Armstrong, D. G., Giurini, J., Edwards, A., Kravette, M., Kravitz, S., . . . Vanore, J. (2000). Diabetic foot disorders: A clinical practice guideline. American College of Foot and Ankle Surgeons. *The Journal of Foot and Ankle Surgery, 39*(Suppl. 5), s1-s60.
- Greenhalgh, T., Helman, C., & Chowdhury, A. M. (1998). Health beliefs and folk models of diabetes in British Bangladeshis: A qualitative study. *British Medical Journal, 316*(7136), 978-983. doi:10.1136/bmj.316.7136.978
- Guba, E. G., & Lincoln, Y. S. (1982). Epistemological and methodological basis of naturalistic inquiry. *Educational Communication and Technology, 30*, 233-252.
- Hasseler, M. K., Heide, M., & Indefrey, S. (2011). Resources for and barriers to effective diabetes care management-experiences and perspectives of people with type 2 diabetes. *Journal of Public Health, 19*, 65-71.

- Hatcher, E., & Whittemore, R. (2007). Hispanic adults' beliefs about type 2 diabetes: Clinical implications. *Journal of the American Academy of Nurse Practitioners, 19*(10), 536-545.
- Hawthorne, K., Robles, Y., Cannings-John, R., & Edwards, A. G. (2008). Culturally appropriate health education for type 2 diabetes in ethnic minority groups: A systematic and narrative review of randomized controlled trials. *Diabetic Medicine, 27*, 613-632.
- Heisler, M. (2007). Overview of peer support models to improve diabetes self-management and clinical outcomes. *Diabetes Spectrum, 20*(4), 214-221.
- Heisler, M., Bouknight, R. R., Hayward, R. A., Smith, D. M., & Kerr, E. A. (2002). The relative importance of physician communication, participatory decision making, and patient understanding in diabetes self-management. *Journal of General Internal Medicine, 17*(4), 243-252.
- Heisler, M., Vijan, S., Makki, F., & Piette, J. (2010). Diabetes control with reciprocal peer support versus nurse care management. A randomized control trial. *Annals of Internal Medicine, 153*(8), 507-515. doi:10.7326/0003-4819-153-8-201010190-00007.
- Helman, C. (2001). *Culture health and illness* (4th ed.). London: Arnold.
- Herman, W. H., Kinmonth, A. L., Wareham, N. J., & Williams, R. (2010). *The evidence base for diabetes care* (2nd ed.). United Kingdom: John Wiley & Sons.
- Hettiarachi, U. P., Ekanayaka, S., & Welihinda, J. (2009). Do Sri Lankan meals help decrease blood glucose response? *Ceylon Medical Journal, 54*, 39-45.

- Hjelm, K., & Mufanda, E. (2010). Zimbabwean diabetics 'beliefs about health and illness: an interview study. *BMC International Health and Human Rights*, 10(7), 1-10.
- Horne, M., & Tierney, S. (2012). What are the barriers and facilitators to exercise and physical activity up take and adherence among South Asian Older adults: A systematic review of qualitative studies. *Preventive Medicine*, 55(4), 276-284. doi:10.1016/j.ypmed.2012.07.016
- Hornsten, A., Lundman, B., Stenlund, H., & Sandstrom, H. (2005). Metabolic improvement after intervention focusing on personal understanding in type 2 diabetes. *Diabetes Research and Clinical Practices*, 68, 65-74. doi:10.1016/j.diabres.2004.08.003
- Howard, A. A., Arnsten, J. H., & Gourevitch, M. N. (2004). Effect of alcohol consumption on diabetes mellitus: A systematic review. *Annals of Internal Medicine*, 140(3), 211-219. doi:10.7326/0003-4819-140-6-200403160-00011
- Hsu, C. C., Lee, C. H., Wahlqvist, M. L., Huang, H. L., Chang, H. Y., Chen, L., . . . Cheng, J. S. (2012). Poverty increases type 2 diabetes incidence and inequality of care despite universal health coverage. *Diabetes Care*, 35(11), 2286-2292. doi:10.2337/dc11-2052
- Hu, J., Gruber, K. J., Liu, H., Zhao, H., & Garcia, A. A. (2012). Diabetes knowledge among older adults with diabetes in Beijing, China. *Journal of Clinical Nursing*, 22(1-2), 51-60. doi:10.1111/j.1365-2702.2012.04273.x
- Huber, C., Huber, J. W., & Shaha, M. (2011). Diabetes care dependent older adults: an exploratory study of nurses 'perspectives. *European Diabetes Nursing*, 8(3), 88-92.

- Huebschmann, A. G., Crane, L. A., Belansky, E. S., Scarbro, S., Marshall, S. J., & Regensteiner, J. G. (2011). Fear of injury with physical activity is greater in adults with diabetes than in adults without diabetes. *Diabetes Care*, *34*(8), 1717-1722. doi:10.2337/dc10-1801
- Hunt, L. M., Arar, N. H., & Akana, L. L. (2000). Herbs, prayer, and insulin use of medical and alternative treatments by a group of Mexican American diabetes patients. *The Journal of Family Practice*, *49*, 216-223.
- Hunt, L. M., Pugh, J., & Valenzuela, M. (1998). How patients adapt diabetes self-care recommendations in everyday life. *Journal of Family Practice*, *46*(3), 207-215.
- Idampititya, C., Lenora, D., Lau, S., Thomson, G. A., & Fernando, D. J. (2008). Letter to the editor. *Diabetes Research and Clinical Practice*, *80*, e25-e26.
- Illangsekera, U. (2011). Towards a cost effective delivery of diabetes care in Sri Lanka. *Sri Lanka Journal of Diabetes Endocrinology and Metabolism*, *1*, 55-57.
- International Diabetes Federation. (2013). *International Diabetes Federation: Diabetes atlas* (6th ed.). Retrieved from: http://www.idf.org/sites/default/files/EN_6E_Atlas_Full_0.pdf
- Jacobson, A. M., de Groot, M., & Samson, J. A. (1994). The evaluation of two measures of quality of life in patients with type 1 and type 11 diabetes. *Diabetes Care*, *17*(4), 267-274. doi:10.2337/diacare.17.4.267
- Jallinoja, P., Absetz, P., Kuronen, R., Nissinen, A., Talja, M., Uutela, A., & Patja, K. (2007). The dilemma of patient responsibility for lifestyle change: Perceptions among primary care physicians and nurses. *Scandinavian Journal of Primary Health Care*, *25*(4), 244-249.

- Järvi, A. N., Karlström, B. E., Granfeldt, Y. E., Björck, I. E., Asp, N. L., & Vessby, B. O. (1999). Improved glycemic control and lipid profile and normalized fibrinolytic activity on a low-glycemic index diet in type 2 diabetic patients. *Diabetes Care*, 22, 10-18.
- Jayawardena, R., Byrne, N. M., Soares, M. J., Katulanda, P., & Hills, A. P. (2012). Food consumption of Sri Lankan adults: An appraisal of serving characteristics. *Public Health Nutrition*, 16, 653-658.
doi:10.1017/S1368980012003011
- Jayawardene, M. H., Idampitiya, C., Jayawarna, C., Wanigasuriya, K., Thomson, G. A., & Fernando, D. J. (2007). Letter to the editor. *Diabetes Research and Clinical Practice*, 75, 249-251.
- Jenkins, D. J. A., Wolever, T. M. S., Taylor, R. H., Barker, H., Fielden, H., Baldwin, . . . Goff, D. V. (1981). Glycemic index of foods: A physiological basis for carbohydrate exchange. *American Journal of Clinical Nutrition*, 34(3), 362-366.
- Jinadasa, C. V., & Jeewantha, M. (2011). A study to determine the knowledge and practices of foot care in patients with chronic diabetic ulcer. *International Journal of Collaborative Research on Internal Medicine & Public Health*, 3, 115-122.
- Jones, L., Crabb, S., Turnbull, D., & Oxlad, M. (2013). Barriers and facilitators to effective type two diabetes management in a rural context: A qualitative study with diabetes patients and health professionals. *Journal of Health Psychology*, 19(3), 441-453. doi:10.1177/1359105312473786

Jones, R. A., Utz, S., Wenzel, J., Steeves, R., Hinton, I., Andrews, D., . . . Oliver, N.

(2006). Use of complementary and alternative therapies by rural African Americans with type 2 diabetes. *Alternative Therapies in Health Medicine*, 12(5), 34-38.

Katulanda, P., Constantine, G. R., Mahesh, J. G., Sheriff, R., Senevirathna, R. D.,

Wijerathne, S., . . . Matthews, D. R. (2008). Prevalence and projections of diabetes and pre-diabetes in adults in Sri Lanka--Sri Lanka Diabetes, Cardiovascular Study (SLDCS). *Diabetic Medicine*, 25(9), 1062-1069
doi:10.1111/j.1464-5491.2008.02523.x

Katulanda, P., Constantine, G. R., Weerakkody, M. I., Perera, Y. S., Jayawardena, M.

G., Wijegoonawardena, P., . . . Sheriff, M. H. (2011). Can we bridge the gap? Knowledge and practices related to diabetes mellitus among general practitioners in a developing country: A cross-sectional study. *Asia Pacific Family Medicine*, 10, 1-6.

Katulanda, P., Ranasinghe, P., Jayawardena, R., Constantine, G. R., Sheriff, M. H., &

Matthews, D. R. (2012). The prevalence, patterns and predictors of diabetic peripheral neuropathy in a developing country. *Diabetology & Metabolic Syndrome*, 4, 21. doi:10.1186/1758-5996-4-21

Katulanda, P., Rathnapala, D., Sheriff, R., & Matthews, D. R. (2011). Provinces and

ethnic specific prevalence of diabetes among Sri Lankan adults. *Journal of Diabetes Endocrinology and Metabolism*, 1, 2-7.

- Khabatt, M., Khader, Y. S., Al-Khawaldeh, A., & Ajlouni, K. (2010). Factors associated with poor glycemic control among adults with type 2 diabetes. *Journal of Diabetes and Its Complications*, 24(2), 84-89. doi:10.1016/j.jdiacomp.2008.12.008.
- Kitabchi, A. E., Umpierrez, G. E., Milles, J. M., & Fisher, J. N. (2009). Hyperglycemic crises in adult patients with diabetes. *Diabetes Care*, 32(7), 1335-1343. doi:10.2337/dc09-9032
- Kleinman, A. (1980). *Patients and healers in the context of culture*. Berkeley, CA: University of California Press.
- Kohinor, M. J., Stronks, K., Nicolaou, M., & Haafkens, J. A. (2011). Considerations affecting dietary behavior of immigrants with type 2 diabetes: A qualitative study among Surinamese in the Netherlands. *Ethnicity & Health*, 16(3), 245-258. doi:10.1080/13557858.2011.563557
- Korkiakangas, E. E., Alahuhta, M. A., & Laitinen, J. H. (2009). Barriers to regular exercise among adults at high risk or diagnosed with type 2 diabetes: A systematic review. *Health Promotion International*, 24(4), 416-426. doi:10.1093/heapro/dap031
- Krawinkel, M. B., & Keding, G. B. (2006). Bittergourd: A dietary approach to hyperglycemia. *Nutr Rev*, 64(7), 331-337.
- Kulkarni, K. D. (2004). Food, Culture and diabetes in the United States. *Clinical Diabetes*, 22(4), 190-192. doi:10.2337/diaclin.22.4.190

- Kwan, J., Razzaq, A., Leiter, L. A., Lillie, D., & Hux, J. E. (2008). Low socio economic status and absence of supplemental health insurance as barriers to diabetes care access and utilization. *Canadian Journal of Diabetes*, 32(3), 174-181.
- Lai, W., Chie, W., & Lew-Ting, C. Y. (2007). How diabetic patients' ideas of illness course affect non-adherent behavior: A qualitative study. *British Journal of General Practice*, 57(537), 296-302.
- Lai, W.A., Lew-Ting, C. Y., & Chie, W.-C. (2004). How diabetic patients think about and manage their illness in Taiwan. *Diabetes UK. Diabetic Medicine*, 22(3), 286-292. doi:10.1111/j.1464-5491.2004.01406.x
- Lawton, J., Ahmad, N., Hallowell, N., Hanna, L., & Douglas, M. (2005). Perceptions and experiences of taking oral hypoglycemic agents among people of Pakistan and Indian Origin: Qualitative study. *British Medical Journal*, 330, 1247-1249. doi:10.1136/bmj.38460.642789.E0
- Lawton, J., Peel, E., Parry, O., & Douglas, M. (2008). Patients' perceptions and experiences of taking oral glucose-lowering agents: A longitudinal qualitative study. *Diabetic Medicine*, 25(4), 491-495. doi:10.1111/j.1464-5491.2008.02400.x
- Leininger, M. M. (1985). Ethnography and ethnonursing: Models and modes of qualitative data analysis. In M. M. Leininger (Ed.), *Qualitative research methods in nursing*. New York: Grune & Stratton.
- Lincoln, Y. S., & Guba, E. G. (1985). *Naturalistic inquiry*. London: SAGE.

- Lipsky, B. A., Berendt, A. R., Deery, H. G, Embil, J. M., Joseph, W. S., Karchmer, A. W., . . . Tan, J. S. (2004). Diagnosis and treatment of diabetic foot infections. *Clinical Infectious Disease*, 39(7), 885-910.
- Little, A. W., & Sabates, R. (2008). Economic globalization, youth expectations and social class: The case of Sri Lanka. *The International Journal of Educational Development*, 28, 708-722.
- Lundberg, P. C., & Thrakul, S. (2012). Type 2 diabetes: How do Thai Buddhist people with diabetes practice self- management? *Journal of Advanced Nursing*, 68, 550-558. doi:10.1111/j.1365-2648.2011.05756.x.
- Lundberg, P. C., & Thrakul, S. (2013). Religion and self-management of Thai Buddhist and Muslim women with type 2 diabetes. *Journal of Clinical Nursing*, 22, 1907-1916. doi:10.1111/jocn.12130.
- Lysaght, M. J. (2002). Maintenance dialysis population dynamics: Current trends and long-term implications. *Journal of American Society of Nephrology*, 13, s37-s40.
- Matthews S. M., Peden A. R., & Rowles, G. D. (2007). Patient-provider communication: Understanding diabetes management among adult females. *Patient Education and Counseling*, 76, 31-37. doi:10.1016/j.pec.2008.11.022
- Mayberry, L. S., & Osborn, C. Y. (2012). Family support, medication adherence and glycemic control among adults with type 2 diabetes. *Diabetes Care*, 35, 1239-1245.
- McCloskey, J., & Flenniken, D. (2010). Overcoming cultural barriers to diabetes control: A qualitative study of southwestern New Mexico Hispanics. *Journal of Cultural Diversity*, 17(3), 110-115.

- Meyer, K. A., Kushi, L. H., Jacobs, D. R., Slavin, J., Sellers, T. A., & Folsom, A. R. (2000). Carbohydrates, dietary fiber, and incident type 2 diabetes in older women. *The American Journal of Clinical Nutrition*, *71*, 921-930.
- Miech, R. A., Kim, J., McConnell, C., & Hamman, R. F. (2009). A growing disparity in diabetes related mortality. U.S. trends, 1989-2005. *American Journal of Preventive Medicine*, *36*, 126-132.
- Mier, N., Medina, A. A., & Ory, M. G. (2007). Mexican Americans with type 2 diabetes: Perspectives on definitions, motivators, and programs of physical activity. *Preventing Chronic Disease, Public Health Research, Practice and Policy*, *4*, 1-7.
- Miles, M. B., & Huberman, A. M. (1994). *Qualitative data analysis: A sourcebook of new methods*. Beverly Hills: SAGE.
- Ministry of Health. (2007). *Annual health bulletin*. Sri Lanka: Ministry of Health.
- Morgan, D. L. (1993). *Successful focus groups: Advanced the State of the Art*. London: SAGE..
- Morse, J. M., & Field, P. A. (1995). *Qualitative research methods for health professionals* (2nd ed.). London: SAGE.
- Murata, G. H., Shah, J. H., Hoffman, R. M., Wendel, C. S., Adam, K. D., Solvas, P. A., . . . Duckworth, W. C. (2003). Intensified blood glucose monitoring improves glycemic control in stable, insulin-treated veterans with type 2 diabetes: The Diabetes Outcomes in Veterans Study (DOVES). *Diabetes Care*, *26*, 1759-1763.
- Nagelkerk, J., Reick, K., & Meengs, L. (2006). Perceived barriers and effective strategies to diabetes management. *Journal of Advanced Nursing*, *54*(2), 151-158.

- Najm, W., Reinsch, S., Hoehler, F., & Tohis, J. (2003). Use of complementary medicine and alternative medicine among ethnic elderly. *Alternative Therapies in Health and Medicine*, 9(3), 50-57.
- Nanayakkara, V., & Ekanayaka, L. (2008). Use of traditional medicine for oral conditions in rural Sri Lanka. *International Dental Journal*, 58(2), 86-90. Doi:10.1922/IDJ_1835Ekanayake05.
- Nathan, D. M., Buse, J. B., Davidson, M. B., Heine, R. J., Holman, R. R., Sherwin, R., & Zinman, B. (2006). Management of hyperglycemia in type 2 diabetes: A consensus algorithm for the initiation and adjustment of therapy: A consensus statement from the American Diabetes Association and the European Association for the Study of Diabetes. *Diabetes Care*, 29(8), 1963-1972.
- Nirogi Lanka. (2009). *Nirogi Lanka project diabetes task force*. Retrieved from <http://nirogilanka.org>
- O'Reilly, K. (2005). *Ethnographic methods*. London: Rutledge Taylor & Francis Group.
- Ofteda, B., Karlsen, B., & Bru, E. (2010). Perceived support from health care practitioners among adults with type 2 diabetes. *Journal of Advanced Nursing*, 66(7), 1500-1509. doi:10.1111/j.1365-2648.2010.05329.x
- Okura, T., Heisler, M., & Langa, K. (2010). The association of cognitive function and social support with glycemic control in adults with diabetes. *Journal of American Geriatric Society*, 57, 1816-1824.
- Omery, A. (1988). Ethnography. In B. Sarter (Ed.), *Paths to knowledge: Innovative research methods for nursing*. New York: National League for Nursing.

- Ouyang, C.-M. (2007). *Factors affecting diabetes self-care among patients with type 2 diabetes in Taiwan* (Doctoral dissertation, Tufts University, Boston). Available from ProQuest Dissertation and Theses database. (UMI No. 3300629)
- Oxford Dictionaries. (2011). *Culture*. Retrieved from <http://oxforddictionaries.com/definition/culture>
- Patel, V., Morrissey, J., Goenka, N., James, D., & Shaikh, S. (2001). Diabetes care in the Hindu patient: Cultural and clinical aspects. *British Journal of Diabetes & Vascular Disease, 1*(2), 132-135. doi:10.1177/14746514010010021301
- Peek, M. E., Odoms-Young, A., Quinn, M. T., Gorawara-Bhat, R., Wilson, S. C., & Chin, M. H. (2010). Race and shared decision-making: perspectives of African-Americans with diabetes. *Social Science and Medicine, 71*, 1-9. doi:10.1016/j.socscimed.2010.03.014.
- Peyrot, M., & Rubin, R. (1997). Levels and risks of depression and anxiety symptomatology among diabetic adults. *Diabetes Care, 20*(4), 1904-1909.
- Pieris, I. (1999). *Disease, treatment and health behaviors in Sri Lanka*. New Delhi: Oxford University Press.
- Pinidiyapathirage, M., Warnakulasuriya, T., Kasturiratne, K., Ranawaka, U., Gunasekera, D., Wijekoon, N., . . . Wickremasinghe, A. (2011). Comparison of urban diabetics with optimal and suboptimal control. *Journal Epidemiol Community Health, 65*(Suppl. 1), A312. doi:10.1136/jech.2011.142976k.58
- Poss, J. E., & Jezewski, M. & Stuart, A. G. (2003). Home remedies for type 2 diabetes used by Mexican Americans in El Paso, Texas. *Clinical Nursing Research, 12*, 304-323.

- Rais, A. (2010). *Is it really God's will?* London: Diabetes UK. Retrieved from http://www.diabetes.org.uk/How_we_help/Magazines/Balance/Past-issues/Archived-issues-and-articles/JulyAugust-2010/Is-it-really-Gods-will/
- Rajakumaraswamy, N., Rajapakse, I., & Fernando, D. S. (2008). The frequency of cognitive dysfunction in elderly Sri Lankans with type 2 diabetes mellitus. *International Journal of Geriatric Psychiatry, 23*, 1205-1206. doi:10.1002/gps.2022
- Ramachandran, A., Snehalatha, C., & Ma, R. C. (2013). Diabetes in South-East Asia: An update for 2013 for the IDF Diabetes Atlas. *Diabetes Research and Clinical Practice, 103*(2), 231-237. doi:10.1016/j.diabres.3013.11.011
- Ramachandran, A., Snehalatha, C., Shetty, A., & Nanditha, A. (2012). Trends in prevalence of diabetes in Asian countries. *World Journal of Diabetes, 3*(6), 110-117. doi: 10.4239/wjd.v3.i6.110
- Rasmussen, O. W., Thomsen, C., Hansen, K. W., Vesterlund, M., Winther, E., & Hermansen, K. (1993). Effects on blood pressure, glucose, and lipid levels of a high-monounsaturated fat diet compared with a high-carbohydrate diet in NIDDM subjects. *Diabetes Care, 16*(12), 1565-1571.
- Raymond, N. T., Varadhan, L., Reynold, D. R., Bush, K., Sankaranarayanan, S., Bellary, S., . . . O'hare, J. P. (2009). Higher prevalence of retinopathy in diabetic patients of south asian ethnicity compared with white europeans in the community. *Diabetes Care, 32*(3), 410-415.
- Rhee, M. K., Cook, C. B., Dunbar, V. G., Panayioto, R. M., Berkowitz, K. J., Boyd, B., . . . Phillips, L. S. (2005). Limited health care access impairs glycemic control in low income urban African Americans with type 2 diabetes. *Journal of Health Care for the Poor and Underserved, 16*, 734-746.

- Rivera, J. O., Ortiz, M., Lawsone, M. E., & Verma, K. (2002). Evaluation of the use of complementary and alternative medicine in the largest United States-Mexico border city. *Pharmacotherapy*, 22(2), 256-264.
- Ropper, J. M., & Shapira, J. (2000). *Ethnography in nursing research*. London: Sage.
- Rosal, M. C., Ebbeling, C. B., Lofgren, I., Ockene, J. K., Ockene, I. S., & Hebert, J. R. (2001). Facilitating dietary change: The patient-centered counseling model. *Journal of the American Dietetic Association*, 101(3), 332-341.
- Rothman, R., Malone, R., Bryant, B., Dewalt, D., & Pignone, M. (2002). Health literacy and diabetes control. *Journal American Medical Association*, 288(21), 2687-2688.
- Russell, S. (2005). Treatment seeking behavior in urban Sri Lanka: Trusting the state, trusting private providers. *Social Science & Medicine*, 61(7), 1396-1407.
- Ryan, D. H., Espeland, M. A., Foster, G. D., Haffner, S. M., Hubbard, V. S., Johnson, K. C., . . . Yanovski, S. K. (2003). Look AHEAD (Action for Health in Diabetes): Design and methods for a clinical trial of weight loss for the prevention of cardiovascular disease in type 2 diabetes. *Control Clinical Trials*, 24(5), 610-628.
- Sabate, E. (2003). *Adherence to long-term therapies: Evidence for action*. Geneva: World Health Organization.
- Sachs, L., & Tomson, G. (1992). Medicines and culture--A double perspective on drug utilization in a developing country. *Social Science Medicine*, 34(3), 307-315.

- Samuel-Hodge, C. D., Headen, S. W., Skelly, A. H., Ingram, A. F., Keyserling, T.C., Jackson, E. J., . . . Elasy, T. A. (2000). Influences on day-to-day self-management of type 2 diabetes among African-American women. *Diabetes Care*, 23(7), 928-933.
- Sanz, C., Gautier, J. F., & Hanaire, H. (2010). Physical exercise for the prevention and treatment of type 2 diabetes. *Diabetes & Metabolism*, 36(5), 346-351. doi:10.1016/j.diabet.2010.06.001
- Savoca, M., & Miller, C. (2001). Food selection and eating patterns: Themes found among people with type 2 diabetes mellitus. *Journal of Nutrition Education*, 33(4), 224-233.
- Saydah, S., & Lochner, K. (2010). Socioeconomic status and risk of diabetes related mortality in the U.S. *Public Health Report*, 125(3), 377-388.
- Schillinger, D., Grumbach, K., Piette, J., Wang, F., Osmond, D., Daher, C., . . . Bindman, A. B. (2002). Association of health literacy with diabetes outcomes. *Journal American Medical Association*, 288(4), 475-482. doi:10.1001/jama.288.4.475
- Seidman, I. (2006). *Interviewing as qualitative research* (3rd ed.). New York: Teachers College Press.
- Seligman, H. K., Jacobs, E. A., López, A., Tschann, J., & Fernandez, A. (2012). Food insecurity and glycemic control among low-income patients with type 2 diabetes. *Diabetes Care*, 35, 233-238. doi:10.2337/dc11-1627.

- Selvin, E., Marinopoulos, S., Berkenblit, G., Rami, T., Brancati, F. L., Powe, N. R., & Golden, S. H. (2004). Meta-analysis: glycosylated hemoglobin and cardiovascular disease in diabetes mellitus. *Annals of Internal Medicine*, *141*(6), 421-431.
- Senevirathne, U., Rajapaksha, P., Pathirana, R., & Seetha, T. (2002). Knowledge, attitudes and practices of epilepsy in rural Sri Lanka. *Seizure*, *11*(1), 40-43.
- Sheard, N. F., Clark, N. G., Brand-Miller, J. C., Franz, M. J., Pi-Sunyer, F. X., Mayer-Davis, E., . . . Geil, P. (2004). Dietary carbohydrate (amount and type) in the prevention and management of diabetes: a statement by the American Diabetes Association. *Diabetes Care*, *27*(9), 2266-2271.
- Shiu, A. T., & Wong, R. Y. (2002). Fears and worries associated with hypoglycemia and diabetes complications: perspectives and experience of Hong Kong Chinese clients. *Journal of Advanced Nursing*, *39*(2), 155-163.
- Sicree, R., Shaw, J., & Zimmet, P. (2006). Diabetes and Impaired glucose tolerance. In D. Gan (Ed.), *Diabetes atlas* (3 rd ed., pp. 15-103). Belgium: International Diabetes Federation.
- Smith, C. A. (2012). Living with sugar: Influence of cultural beliefs on type 2 diabetes self-management of English-speaking Afro-Caribbean Women. *Journal of Immigrant and Minority Health*, *14*(4), 640-647.
doi:10.1007/s10903-011-9513-2
- Sowattanagoon, N., Kotchabhakdi, N., & Petrie, K. J. (2009). The influence of Thai culture on diabetes perceptions and management. *Diabetes Research and Clinical Practice*, *84*(3), 245-251. doi:10.1016/j.diabres.2009.02.011
- Spradley, J. P. (1979). *The ethnographic interview*. London: Harcourt College.

Spradley, J. P. (1980). *Participant observation*. New York: Holt, Rinehart & Winston.

Stratton, I. M., Adler, A. I., Neil, H. A., Matthews, D. R., Manley, S. E., Cull, C. A.,

. . . Holman, R. R. (2000). Association of glycaemia with macrovascular and microvascular complications of type 2 diabetes (UKPDS 35):

Prospective observational study. *British Medical Journal*, *321*, 405-412.

doi:10.1136/bmj.321.7258.405

Streubert, H. J., & Carpenter, D. R. (2011). *Qualitative research in nursing: Advancing the humanistic imperative* (5th ed.). London: Lippincott Williams & Wilkins.

Thomas, D. E., Elliott, E. J., & Naughton, G. A. (2006). Exercise for type 2 diabetes mellitus. *Cochrane Database Systematic Review*, *19*(3), CD002968.

Thompson, S. J., Gifford, S. M., & Thorpe, L. (2000). Social and cultural context of risk and prevention: Food and physical activity in an urban Aboriginal community. *Health Education & Behavior*, *27*(6), 725-745.

Umpierre, D., Ribeiro, P. A., Kramer, C. K., Leitão, C. B., Zucatti, A. T., Azevedo, M. J., . . . Schaan, B. D. (2011). Physical activity advice only or structured

exercise training and association with HbA1c levels in type 2 diabetes: A systematic review and meta-analysis. *Journal of American Medical*

Association, *305*(17), 1790-1799. doi:10.1001/jama.2011.576

Unger, J. (2007). *Diabetes management in primary care*. New York: Lippincott, Williams & Wilkins.

United Kingdom Prospective Diabetes Study Group. (1990). UK prospective diabetes study 7: Response of fasting plasma glucose to diet therapy in newly presenting type II diabetic patients. *Metabolism*, *39*(9), 905-912.

- United Kingdom Prospective Diabetes Study Group. (1998). Intensive blood-glucose control with sulphonylureas or insulin compared with conventional treatment with risk complications in patients with Type 2 diabetes. *Lancet*, 352(9144), 837-853.
- Uragoda, C. G. (1987). *History of medicine in Sri Lanka*. Colombo, Sri Lanka: Medical Association, .
- Utz, S. W., Steeves, R. H., Wenzel, J., Hinton, I., Jones, R. A., Andrews, D., . . . Oliver, M. N. (2006). "Working hard wiht it": Self-management of type 2 diabetes by rural African Americans. *Family Community Health*, 29(3), 195-205.
- Valenzuela, G. A., Mata, J. E., Mata, A. S., Gabali, C., Gaona, E., Thom, D., & LeBaron, S. (2010). Knowledge and beliefs regarding type 2 diabetes mellitus in rural Mexico. *Ethnicity and Health*, 89(4), 353-360.
- Vinink, A. J., & Mehrabyan, A. (2004). Diabetic neuropathies. *The Medical Clinics of North America*, 88(4), 947-99, xi.
- Vithanage, P. V., Panapitiya, P. A., Padmakumara, N., Hemantha, S., Kasturiratne, K.T., Wickremasinghe, A. R., . . . Pinidiyapathirage, M. J. (2011). Mortality in urban cohort in Ragama, Sri Lanka. *Journal of Epidemiol Community Health*, 65(Suppl. 1), A398-A399.
- Wallin, A.-M., Lovfvander, M., & Ahlstrom, G. (2007). Diabetes: A cross-cultural interview study of immigrants from Somalia. *Journal of Clinical Nursing*, 16(11), 305-314. doi:10.1111/k.1365-2702.2007.02099.x
- Wanasingha, Y. A. (1995). Spatial variations in the provision of health care facilities in Sri Lanka. *Health & Place*, 1(4), 235-242.

- Weerasingha, M. C., & Fernando, D. N. (2011). Paradox in treatment seeking: An experience from rural Sri Lanka. *Qualitative Health Research, 21*(3), 365-372. doi:10.1177/1049732310385009
- Wellard S. J., Rennie, S., & King, R. (2008). Perception of people with type 2 diabetes about self management and the efficacy of community based service. *Contemporary Nurse: Journal for the Australian Nursing Profession, 29*(2), 218-226.
- Wen, L., Shepherd, M., & Parchman, M. (2005). Family support, diet and exercise among older Mexican Americans with type 2 diabetes. *The Diabetes Educator, 30*, 980-993.
- Whites, K. M., Terry, D. J., Troup, C., & Rempel, L. A. (2007). Behavioral, normative and control beliefs underlying low-fat dietary and regular physical activity behaviors for adults diagnosed with type 2 diabetes and/or cardiovascular disease. *Psychology, Health & Medicine, 12*(4), 485-494.
- Wijewardene, K., Mohideen, M. R., Mendis, S., Fernando, D. S., Kulathilaka, T., Weerasekara, D., & Uluwitta, P. (2005). Prevalence of hypertension, diabetes and obesity: Baseline findings of a population based survey in four provinces in Sri Lanka. *The Ceylon Medical Journal, 50*(2), 62-70.
- Williams, G., & Pickup, J. C. (2004). *Handbook of diabetes* (3rd ed.). USA: Blackwell.
- World Bank Report. (2011). *Sri Lanka economic update. Poverty Reduction and Economic Management*. Retrieved from <http://siteresources.worldbank.org/SRILANKAEXTN/Resources/233046-1298393467535/SLEconupdateNov2011.pdf>

World Health Organization. (2008). *Country health system profile-Sri Lanka*.

Retrieved from [http://www.searo.who.int/en/Section 313/Section 1524.html](http://www.searo.who.int/en/Section%20313/Section%201524.html)

Xie, W., Zhao, Y., & Zhang, Y. (2011). Traditional Chinese medicine in treatment of patients with type 2 diabetes mellitus. *Evidence Bases Complementary and Alternative Medicine*, 2011, 726723. doi:10.1155/2011/726723

Xu, Y., Pan, W., & Liu, H. (2010). Self-management of Chinese Americans with type 2 diabetes. *Nursing and Health Sciences*, 12(2), 228-234. doi:10.1111/j.1442-2018.2010.00524.x

Yusuf, S., Sleight, P., Pogue, J., Bosch, J., Davies, R., & Dagenais, G. (2000). Effects of angiotensin-converting enzyme inhibitor Ramipril on cardiovascular events in high-risk populations. *New England Journal of Medicine*, 342, 145-153.

Zgibor, J. C., & Songer, T. J. (2001). External barriers to diabetes care: Addressing personal and health systems issues. *Diabetes Spectrum*, 14(1), 23-28. doi:10.2337/diaspect.14.1.23

Zhang, X., Norris, S. L., Gregg, E. W., & Beckles, G. (2007). Social support and mortality among older persons with diabetes. *Diabetes Educator*, 33(2), 273-281.