

REFERENCES

- Agudelo-Silva F, Spielman A. Penetration of mosquito midgut wall by sheathed microfilariae. *J Invertebr Pathol* 1985; 45: 117-119.
- Bangs MJ, Ash LR, Barr AR. Susceptibility of various mosquitoes of California to subperiodic *Brugia malayi*. *Acta Trop* 1995; 59: 323-332.
- Barillas-Mury C, Wells MA. Cloning and sequencing of the blood meal-induced late trypsin gene from the mosquito *Aedes aegypti* and characterization of the upstream regulatory region. *Insect Mol Biol* 1993; 2: 7-12.
- Bian G, Raikhel AS, Zhu J. Characterization of a juvenile hormone-regulated chymotrypsin-like serine protease gene in *Aedes aegypti* mosquito. *Insect Biochem Mol Biol* 2008; 38: 190-200.
- Bain O and Babayan S. Behaviour of filariae: morphological and anatomical signatures of their life style within the arthropod and vertebrate hosts. *Filaria Journal* 2003; 2: 16.
- Billingsley PF, Rudin W. The role of the mosquito peritrophic membrane in bloodmeal digestion and infectivity of *Plasmodium* species. *J Parasitol* 1992; 78: 430-440.
- Chaithong U. Study on *Brugia pahangi* infection in *Aedes albopictus*, *Aedes aegypti* and *Aedes togoi*. Thesis (M.Sc. (Tropical Medicine))--Mahidol University 1976.

- Chang MS, Chan KL, Ho BC. Comparative transmission potential of three *Mansonia* mosquitoes (Diptera: Culicidae) for filariasis in Sarawak, Malaysia. Bull Entomol Res 1991; 81: 437-444.
- Chen CC, Shin CM. Exsheathment of microfilariae of *Brugia pahangi* in the susceptible and refractory strains of *Aedes aegypti*. Ann Trop Med Parasitol 1988; 82: 201-206.
- Chomcharn Y, Surathin K, Bunnag D, Sucharit S, Harinasuta T. Effect of a single dose of primaquine on a Thai strain of *Plasmodium falciparum*. Southeast Asian J Trop Med Public Health 1980; 11: 408-412.
- Choochote W, Chaithong U, Somboon P, Pakdicharoen A, Tookyang B, Likitvong K, Siriprasert V, Sukontasan K, Thitasut P. Small laboratory animal model for nocturnally subperiodic *Brugia malayi* (Narathiwat province, southern Thailand strain). J Trop Med Parasitol 1991; 14: 51-58.
- Choochote W, Sukhavat K, Somboon P, Khamboonruang C, Maleewong W, Suwanpanit P. The susceptibility of small laboratory animals to nocturnally superiodic *Brugia malayi* in Thailand. J Parasitol Trop Med Assoc Thailand 1986; 9: 35-37.
- Christensen BM, Sutherland DR. *Brugia pahangi*: exsheathment and midgut penetration in *Aedes aegypti*. Trans Amer Microscop Soc 1984; 4: 423-433.
- Christensen BM, Li J, Chen CC, Nappi A. Melanization immune response in mosquito vectors. Trends Parasitol 21:192-199 Denham DA, McGreevy PB (1977) Brugian filariasis: epidemiological and experimental studies. Adv Parasitol 2005; 15: 243-309.

- Denham DA, McGreevy PB. Brugian filariasis: epidemiological and experimental studies. *Adv Parasitol* 1977; 15: 243-309.
- Devaney E, Howells RE. The exsheathment of *Brugia pahangi* microfilariae under controlled conditions *in vitro*. *Ann Trop Med Parasitol* 1979; 73: 227-333.
- Edwards MJ, Moskalyk LA, Donnelly-Doman M, Vlaskova M, Noriega FG, Walker VK, Jacobs-Lorena M. Characterization of a carboxypeptidase A gene from the mosquito, *Aedes aegypti*. *Insect Mol Biol* 2000; 9: 33-38.
- Erickson SM, Xi Z, Mayhew GF, Ramirez JL, Aliota MT, Christensen BM, Dimopoulos G. Mosquito infection responses to developing filarial worms. *PLoS Negl Trop Dis* 2009; 3: e529.
- Esslinger JH. Behavior of microfilaria of *Brugia pahangi* in *Anopheles quadrimaculatus*. *Am J Trop Med Hyg* 1962; 11: 749-758.
- Ewert A. Comparative migration of microfilariae and development of *Brugia pahangi* in various mosquitoes. *Am J Trop Med Hyg* 1965; 14: 254-259.
- Felix CR, Betschart B, Billingsley PF, Freyvogel TA. Post-feeding induction of trypsin in the midgut of *Aedes aegypti* is separable into two cellular phases. *Insect Bioch* 1991; 21: 197-203.
- Guptavaniij P, Harinasuta C, Vutikes S, Deesin T, Surathin K. The vectors of *Brugia malayi* in southern Thailand. *Southeast Asian J Trop Med Public Health* 1978; 9: 543-548.
- Isoe J, Zamora J, Miesfeld RL. Molecular analysis of the *Aedes aegypti* carboxypeptidase gene family. *Insect Biochem Mol Biol* 2009a; 39: 68-73.

- Isoe J, Rascon AA Jr, Kunz S, Miesfeld RL. Molecular genetic analysis of midgut serine proteases in *Aedes aegypti* mosquitoes. *Insect Biochem Mol Biol* 2009b; 39: 903-912.
- Iyengar MOT. Entry of *Filaria* larvae into the body cavity of the mosquito. *Parasitology* 1936; 28: 190-195.
- Jariyapan N, Saeung A, Intakhan N, Chanmol W, Sor-Suwan S, Phattanawiboon B, Taai K, Choochote W. Peritrophic matrix formation and *Brugia malayi* microfilaria invasion of the midgut of a susceptible vector, *Ochlerotatus togoi* (Diptera: Culicidae). *Parasitol Res* 2013; 112: 2431-2440.
- Jiang Q, Hall M, Noriega FG, Wells M. cDNA cloning and pattern of expression of an adult, female specific chymotrypsin from *Aedes aegypti* midgut. *Insect Biochem Mol Biol* 1997; 27: 283-289.
- Kalhok SE, Tabak LM, Prosser DE, Brook W, Downe AE, White BN. Isolation, sequencing and characterization of two cDNA clones coding for trypsin-like enzymes from the midgut of *Aedes aegypti*. *Insect Mol Biol* 1993; 2: 71-79.
- Kershaw WE, Edeson JF, Johnson M. The intake of *Brugia malayi* by different strains of *Aedes aegypti*. *Ann Soc Belg Med Trop* 1961; 41: 281-282.
- Kumar NP, Sabesan S, Panicker KN. Susceptibility status of *Mansonia annulifera* to *Brugia malayi* parasites in Cherthala, Alappuzha district, Kerala. *Indian J Exp Biol* 1998; 36: 829-831.
- Lek-Uthai U, Tomoen W. Susceptibility of *Mansonia uniformis* to *Brugia malayi* microfilariae from infected domestic cat. *Southeast Asian J Trop Med Public Health* 2005; 36: 434-441.

- Luo H, Qu FY. Experimental infection index of *Anopheles sinensis* and melanization of periodic *Brugia malayi*. *Zhongguo Ji Sheng Chong Xue Yu Ji Sheng Chong Bing Za Zhi* 1990; 8: 260-263.
- Magalhaes T, Oliveira IF, Melo-Santos MAV, Oliveira CMF, Lima CA, Ayres CFJ. Expression of defensin, cecropin, and transferrin in *Aedes aegypti* (Diptera: Culicidae) infected with *Wuchereria bancrofti* (Spirurida: Onchocercidae), and the abnormal development of nematodes in the mosquito. *Exp Parasitol* 2008; 120: 364-371.
- Melrose WD. Lymphatic filariasis: new insights into an old disease. *Int J Parasitol* 2002; 32: 947-960.
- Michalski ML, Erickson SM, Bartholomay LC, Christensen BM. Midgut barrier imparts selective resistance to filarial worm infection in *Culex pipiens pipiens*. *PLoS Negl Trop Dis* 2010; 4: e875.
- Nayar JK, Knight JW. Comparison of migration and encapsulation of *Brugia malayi* microfilariae from the midgut to the hemocoel between *Anopheles quadrimaculatus* and *Aedes aegypti*. *J Invertebr Pathol* 1995; 65: 295-299.
- Nelson RL. Parity in winter populations of *Culex tarsalis* Coquillett in Kern County, California. *Amer J Hyg* 1964; 80: 242-253.
- Noriega FG, Edgar KA, Bechet R, Wells MA. Midgut exopeptidase activities in *Aedes aegypti* are induced by blood feeding. *J Insect Physiol* 2002; 48: 205-212.
- Nutman TB, Kumaraswami V. Regulation of the immune response in lymphatic filariasis: perspectives on acute and chronic infection with *Wuchereria bancrofti* in South India. *Parasite Immunol* 2001; 23: 389-399.

- O'Connor FW, Beatty H. The early migrations of *Wuchereria bancrofti* in *Culex fatigans*. Trans R Soc Trop Med Hyg 1936; 30: 125-127.
- Oda T, Wada Y. Exsheathment and migration of microfilariae of *Brugia malayi* (Cheju Strain) in mosquitoes. Trop Med 1980; 22: 27-33.
- Owen RR. Non-development of *Brugia pahangi* in a refractory mosquito, *Aedes malayensis*. Ann Trop Med Parasit 1979; 73:193-195.
- Pascoa V, Oliveira PL, Dansa-Petretski M, Silva JR, Alvarenga PH, Jacobs-Lorena M, Lemos FJ. *Aedes aegypti* peritrophic matrix and its interaction with heme during blood digestion. Insect Biochem Mol Biol 2002; 2: 517-523.
- Perrone JB, Spielman A. Microfilarial perforation of the midgut of a mosquito. J Parasitol 1986; 72: 723-727.
- Pothikasikorn J, Bangs MJ, Boonplueang R, Chareonviriyaphap T. Susceptibility of various mosquitoes of Thailand to nocturnal subperiodic *Wuchereria bancrofti*. J Vector Ecol 2008; 33: 313-320.
- Rao UR, Vickery AC, Kwa BH, Nayar JK. *Brugia malayi*: ivermectin inhibits the exsheathment of microfilariae. Am J Trop Med Hyg 1992; 46: 183-188.
- Rascón AA Jr, Gearin J, Isoe J, Miesfeld RL. *In vitro* activation and enzyme kinetic analysis of recombinant midgut serine proteases from the Dengue vector mosquito *Aedes aegypti*. BMC Biochem 2011; 12: 43.
- Saboia-Vahia L, Borges-Veloso A, Cuervo P, Junqueira M, Mesquita-Rodrigues C, Britto C, Domont GB, De Jesus JB. Protein expression in the midgut of sugar-fed *Aedes albopictus* females. Parasit Vectors 2012; 5: 290.

- Saeung A, Choochote W. Development of a facile system for mass production of *Brugia malayi* in a small space laboratory. *Parasitol Res* 2013; 112: 3259-3265.
- Saeung A, Hempolchom C, Baimai V, Thongsahuan S, Taai K, Jariyapan N, Chaithong U, Choochote W. Susceptibility of eight species members in the *Anopheles hyrcanus* group to nocturnally subperiodic *Brugia malayi*. *Parasit Vectors* 2013; 6: 5.
- Sakamoto M, Ohta M, Suzuki A, Takase H, Yoshizawa Y, Kitami M, Sato R. Localization of the serine protease homolog BmSPH-1 in nodules of *E. coli*-injected *Bombyx mori* larvae and functional analysis of its role in nodule melanization. *Dev Comp Immunol* 2011; 35: 611-619.
- Santos JN, Lanfredi RM, Pimenta PF. The invasion of the midgut of the mosquito *Culex (Culex) quinquefasciatus* Say, 1823 by the helminth *Litomosoides chagasfilhoi* Moraes Neto, Lanfredi and De Souza, 1997. *J Invertebr Pathol* 2006; 93: 1-10.
- Schacher JF. Morphology of the microfilaria of *Brugia pahangi* and of the larval stages in the mosquito. *J Parasitol* 1962; 48: 679-692.
- Tchankouo-Nguetchou S, Khun H, Pincet L, Roux P, Bahut M, Huerre M, Guette C, Choumet V. Differential protein modulation in midguts of *Aedes aegypti* infected with chikungunya and dengue 2 viruses. *PLoS One* 2010; 5: e13149.
- Tellam RL, Wijffels G, Willadsen P. Peritrophic matrix proteins. *Insect Biochem Mol Biol* 1999; 29: 87-101.

- Terra WR. The origin and functions of the insect peritrophic membrane and peritrophic gel. *Arch Insect Biochem Physiol* 2001; 47: 47-61.
- Thomas P, Kenny N, Eyles D, Moreira LA, O'Neill SL, Asgari S. Infection with the wMel and wMelPop strains of *Wolbachia* leads to higher levels of melanization in the hemolymph of *Drosophila melanogaster*, *Drosophila simulans* and *Aedes aegypti*. *Dev Comp Immunol* 2011; 35: 360-365.
- Tokura A, Fu GS, Sakamoto M, Endo H, Tanaka S, Kikuta S, Tabunoki H, Sato R. Factors functioning in nodule melanization of insects and their mechanisms of accumulation in nodules. *J Insect Physiol* 2014; 60: 40-49.
- Trpis M. Susceptibility of the autogenous group of the *Aedes scutellaris* complex of mosquitoes to infection with *Brugia malayi* and *Brugia pahangi*. *Tropenmed Parasitol* 1981; 32: 184-188.
- Townson H, Chaithong U. Mosquito host influences on development of filariae. *Ann Trop Med Parasitol* 1991; 85: 149-163.
- Wada Y. Vector mosquitoes of filariasis in Japan. *Trop Med Health* 2001; 39: 39-45.
- Weinstein PP. Development *in vitro* of the microfilariae of *Wuchereria bancrofti* and of *Litomosoides carinii* as far as the sausage form. *Trans R Soc Trop Med Hyg* 1963; 57: 236.
- WHO. Control of lymphatic filariasis. A manual for health personnel. Geneva: WHO 1987.
- Yamamoto H, Ogura N, Kobayashi M, Chigusa Y. Studies on filariasis II: exsheathment of the microfilariae of *Brugia pahangi* in *Armigeres subalbatus*. *Jpn J Parasitol* 1983; 32: 287-292.

Zou Z, Shin SW, Alvarez KS, Kokoza V, Raikhel AS. Distinct melanization pathways in the mosquito *Aedes aegypti*. *Immunity* 2010; 32: 41-53.